LOCAL PROPERTIES OF QUOTIENT ANALYTIC SPACES

KUNIO TAKIJIMA AND TADASHI TOMARU

ABSTRACT. Let $T \simeq \mathbb{C}/\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$ be a complex 1-torus and E_n the set of all elliptic functions of order n . Then M. Namba showed that E_n is a $2n$ -dimensional complex manifold. Let $\text{Aut } T$ be the automorphism group of T , then $\text{Aut } T$ is a 1-dimensional compact complex Lie group and the orbit space $E_n/\text{Aut } T$ is an analytic space. In this paper, we shall show that $E_n/\text{Aut } T$ has only rational singularities and if $n > 5$, $E_n/\text{Aut } T$ is rigid.

1. Introduction. Let M be a complex manifold and G a properly discontinuous transformation group on M . Then H. Cartan [2] showed that the quotient space M/G is a normal analytic space and D. Burns [1] has proved that M/G is rational. Moreover M. Schlessinger [7], [8] showed that if $\text{codim } F(G) \geq 3$, M/G has only rigid singularities, where

$$F(G) := \bigcup_{g \in G - \{e\}} F(g), \quad F(g) := \{x \in M; gx = x\}$$

is the set of all fixed points.

In this paper, let M be also a complex manifold and G a complex Lie transformation group whose action is proper on M . Then H. Holmann [4] proved that the quotient space M/G is a normal analytic space. We shall show the following

THEOREM 1. *M/G has only rational singularities. Moreover if $\text{codim } S(M/G) \geq 3$, M/G is rigid, where $S(M/G)$ is the set of all singular points of M/G .*

Let $\omega_1, \omega_2 \in \mathbb{C}$; $\text{Im}(\omega_2/\omega_1) > 0$, $T := \mathbb{C}/\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$ be a complex 1-torus and E_n the set of all elliptic functions of order n . Then E_n is a $2n$ -dimensional complex manifold [5] and the automorphism group $\text{Aut } T$ acts on E_n naturally. Then we have

THEOREM 2. *$E_n/\text{Aut } T$ is rational and if $n \geq 5$, $E_n/\text{Aut } T$ has only rigid singularities.*

2. The proof of Theorem 1. We shall give some definitions. Let $(X, \mathcal{O}_X)$ be an analytic space and $\pi: (\tilde{X}, \mathcal{O}_{\tilde{X}}) \rightarrow (X, \mathcal{O}_X)$ a resolution of singularities of X . Then $x \in X$ is called a rational singularity if $(R^i \pi_* \mathcal{O}_{\tilde{X}})_x = 0$ for any $i > 0$

Received by the editors December 13, 1977 and, in revised form, March 27, 1978.

AMS (MOS) subject classifications (1970). Primary 32M05, 32C40.

Key words and phrases. Quotient analytic space, rational singularity, rigid singularity, elliptic function.

© American Mathematical Society 1978

(cf. [1]). $x \in X$ is called a rigid singularity if any local flat deformation of (X, x) is locally trivial (cf. [7], [8]).

Let R be an equivalence relation on X . We identify the graph of R with R . Then R is called open (resp. proper, finite) if the natural projection $p_1: R \rightarrow X$ is open (resp. proper, finite). And R is called analytic if the graph R is an analytic set in $X \times X$ (cf. [4], [9]).

Let M be a complex manifold and G a complex Lie transformation group on M . Then G is called proper on M if the graph mapping $\Psi: G \times M \rightarrow M \times M$, $\Psi(g, x) := (gx, x)$ is proper. (This is different from the above definition that an equivalence relation R is proper.) Let $G(x) := \{gx; g \in G\}$ be the orbit through x and $G_x := \{g \in G; gx = x\}$ an isotropy group at x .

We shall prove Theorem 1 and its proof is essentially owing to the technique by H. Holmann [3], [4]. Let $\Phi: G \times M \rightarrow M$, $\Phi(g, x) := gx$ be the natural holomorphic mapping. Then $G(x) = \Phi(G \times \{x\})$ is a complex submanifold of M and biholomorphic to G/G_x . Since G is proper on M , M/G is Hausdorff with respect to the quotient topology. Hence from Hilfssatz 7 of [4], for any $x \in M$ there exist an open connected neighborhood $U \subset M$ of x and a submanifold $N \subset U$ such that

$$(1) x \in N,$$

$$(2) U \cong (G(x) \cap U) \times N,$$

(3) $G(y) \cap U \cong (G(x) \cap U) \times N_y$ for any $y \in N$, where $N_y := G(y) \cap N$ is finite and $N_x = \{x\}$. Let R be an equivalence relation on N defined by $R(y) := N_y$ for any $y \in N$. Then R is open proper finite analytic (if we choose U small enough) and M/G is locally isomorphic to N/R (cf. the proof of Satz 15 in [4]).

For any $g \in G_x$, there exists an open connected neighborhood $V \subset N$ of x such that $g' := qg: V \rightarrow g'(V) \subset N$ is biholomorphic, where $q: U \rightarrow N$ is the natural projection and the inverse of g' is

$$(g')^{-1} := qg^{-1}.$$

Let $G'(V)$ be the set of all biholomorphic mappings on V induced from G_x for any connected neighborhood $V \subset N$ of x . Since G is proper on M , there exists an open connected neighborhood $V \subset N$ of x such that $G'(V)$ is a finite transformation group on V and R coincides with the equivalence relation induced by $G'(V)$ on V (cf. the proof of Satz 19 in [4]). Therefore M/G is locally isomorphic to $V/G'(V)$ and so M/G has only rational singularities. If we show the following lemma, the proof of Theorem 1 is complete.

LEMMA 1. *Let Γ be a finite subgroup of $GL(n, \mathbb{C})$. If $\text{codim } S(\mathbb{C}^n/\Gamma) \geq 3$, $\mathbb{C}^n/\Gamma$ is rigid.*

PROOF. Let Γ_0 be the normal subgroup of Γ generated by reflections, where $g \in GL(n, \mathbb{C})$ is called a reflection if g is order-finite and $\text{codim } F(g) = 1$.

Then $\mathbb{C}^n/\Gamma_0$ is biholomorphic to $\mathbb{C}^n$ and Γ/Γ_0 acts on $\mathbb{C}^n/\Gamma_0$ naturally. Moreover $\mathbb{C}^n/\Gamma \cong (\mathbb{C}^n/\Gamma_0)/(\Gamma/\Gamma_0)$ and Γ/Γ_0 has no reflection (cf. [6]). Thus we may assume that Γ has no reflection. Then $p(F(\Gamma)) = S(\mathbb{C}^n/\Gamma)$, where $p: \mathbb{C}^n \rightarrow \mathbb{C}^n/\Gamma$ is the projection.

In fact, $p(F(\Gamma)) \supset S(\mathbb{C}^n/\Gamma)$ is clear. Let $x \in \mathbb{C}^n$ and $p(x) \notin S(\mathbb{C}^n/\Gamma)$, then there exists a nonsingular neighborhood $U \subset \mathbb{C}^n/\Gamma$ of $p(x)$ such that $p: p^{-1}(U) \rightarrow U$ is a finite covering between complex manifolds. We have

$$F(\Gamma) \cap p^{-1}(U) = \{t \in p^{-1}(U); Jp(t) = 0\},$$

where Jp is the Jacobian of p . Since $\text{codim } F(\Gamma) \geq 2$, $F(\Gamma) \cap p^{-1}(U)$ must be empty. Thus $x \notin F(\Gamma)$. Therefore $\text{codim } F(\Gamma) = \text{codim } S(\mathbb{C}^n/\Gamma) \geq 3$ and so $\mathbb{C}^n/\Gamma$ is rigid.

REMARK. If G is not proper on M , locally M/G is not always isomorphic to $V/G'(V)$.

3. The proof of Theorem 2. Since $\text{Aut } T$ is compact, $\text{Aut } T$ is proper on E_n . Hence the first part of Theorem 2 is straightforward from Theorem 1. Thus we must calculate $\text{codim } S(E_n/\text{Aut } T)$.

Let S_n be the n -symmetric group. Then $S_n \times S_n$ acts on $\mathbb{C}^{2n+1}$ as follows:

$$\begin{aligned} (S_n \times S_n) \times \mathbb{C}^{2n+1} &\rightarrow \mathbb{C}^{2n+1}, \\ (g_1, g_2, \alpha, \beta, \gamma) &\rightsquigarrow (g_1\alpha, g_2\beta, \gamma), \end{aligned}$$

where $\alpha = (\alpha_1, \dots, \alpha_n)$, $\beta = (\beta_1, \dots, \beta_n)$. We define

$$\begin{aligned} \phi: \mathbb{C}^{2n+1} &\rightarrow \mathbb{C}^{2n+1}, \\ \phi(\alpha, \beta, \gamma) &:= (\sigma_1(\alpha), \dots, \sigma_n(\alpha), \sigma_1(\beta), \dots, \sigma_n(\beta), \gamma), \end{aligned}$$

where σ_i is the i -fundamental symmetric function ($i = 1, \dots, n$). Then we have $\mathbb{C}^{2n+1}/(S_n \times S_n) \cong \phi(\mathbb{C}^{2n+1}) = \mathbb{C}^{2n+1}$.

For $\omega_1, \omega_2 \in \mathbb{C}$; $\text{Im}(\omega_2/\omega_1) > 0$, we put $\Delta := \mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$, $\Delta^* := \Delta - \{0\}$ and

$$\begin{aligned} \tilde{S}_\Omega := \left\{ (\alpha, \beta, \gamma) \in \mathbb{C}^{2n+1}; \sum_{i=1}^n \alpha_i - \sum_{i=1}^n \beta_i = \Omega, \gamma \neq 0, \alpha_i - \beta_j \notin \Delta, \right. \\ \left. \alpha_i - \alpha_j \notin \Delta^*, \beta_i - \beta_j \notin \Delta^* (i, j = 1, \dots, n) \right\} \end{aligned}$$

for $\Omega \in \Delta$. Since $\tilde{S}_\Omega$ is an $S_n \times S_n$ -invariant complex manifold,

$$\begin{aligned} S_\Omega := \phi(\tilde{S}_\Omega) = \{(x, y, z) \in \mathbb{C}^{2n+1}; x_1 - y_1 = \Omega, z \neq 0, \alpha_i - \beta_j \notin \Delta, \\ \alpha_i - \alpha_j \notin \Delta^*, \beta_i - \beta_j \notin \Delta^* (i, j = 1, \dots, n)\} \end{aligned}$$

is also a $2n$ -dimensional complex manifold, where α_i, β_i are roots of

$$X^n - x_1 X^{n-1} + x_2 X^{n-2} + \dots + (-1)^n x_n = 0, \quad (1)$$

$$X^n - y_1 X^{n-1} + y_2 X^{n-2} + \dots + (-1)^n y_n = 0 \quad (2)$$

respectively. And $\tilde{S} := \bigcup_{\Omega \in \Delta} \tilde{S}_\Omega$, $S := \bigcup_{\Omega \in \Delta} S_\Omega$ are complex manifolds.

By the classical theory of elliptic functions, for any $f(w) \in E_n$, let $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n$ be zeroes and poles of $f(w)$ respectively, then

$$\sum_{i=1}^n \alpha_i - \sum_{i=1}^n \beta_i = \Omega \in \Delta, \quad \Omega = p_0\omega_1 + q_0\omega_2,$$

$$f(w) = \gamma \exp 2(p_0\eta_1 + q_0\eta_2)w \prod_{i=1}^n \frac{\sigma(w - \alpha_i)}{\sigma(w - \beta_i)},$$

where $\eta_j := \zeta(\frac{1}{2}\omega_j)$ ($j = 1, 2$) and σ, ζ are the Weierstrass functions. We define $\tilde{F}: \tilde{S} \rightarrow E_n$

$$\tilde{F}(\alpha, \beta, \gamma) := \gamma \exp 2(p_0\eta_1 + q_0\eta_2)w \prod_{i=1}^n \frac{\sigma(w - \alpha_i)}{\sigma(w - \beta_i)},$$

where

$$\sum_{i=1}^n \alpha_i - \sum_{i=1}^n \beta_i = p_0\omega_1 + q_0\omega_2 \in \Delta.$$

Since $\tilde{F}$ is $S_n \times S_n$ -invariant, $\tilde{F}: \tilde{S} \rightarrow E_n$ induces $F: S \rightarrow E_n$

$$F(x, y, z) := z \exp 2(p_0\eta_1 + q_0\eta_2)w \prod_{i=1}^n \frac{\sigma(w - \alpha_i)}{\sigma(w - \beta_i)},$$

where $x_1 - y_1 = p_0\omega_1 + q_0\omega_2 \in \Delta$ and α_i, β_i are roots of (1), (2) respectively. Then we can show the following lemma (cf. [5, Remark, p. 75]).

LEMMA 2. $F: S \rightarrow E_n$ is surjective open holomorphic and locally biholomorphic.

We shall calculate $\text{codim } S(E_n/\text{Aut } T)$ using this locally biholomorphic mapping F . Here we assume that $\omega_2/\omega_1 \not\cong e^{\pi i/2}, e^{2\pi i/3}$ under the modular group. Then $\text{Aut } T$ has two connected components T_0, T_1 and T_0 is isomorphic to T as Lie group. We put

$$g_1(x_1, x_2, x_3, n) := -x_3 + \frac{n-2}{n}x_1x_2 - \frac{(n-1)(n-2)}{3n^2}x_1^3,$$

$$g_2(x_1, x_2, x_3, x_4, x_5, n) := -x_5 + \frac{n-4}{n}x_1x_4 - \frac{(n-3)(n-4)}{n^2}x_1^2x_3 \\ + \frac{2(n-2)(n-3)(n-4)}{3n^3}x_1^3x_2 - \frac{(n-1)(n-2)(n-3)(n-4)}{5n^4}x_1^5,$$

$$\rho(x, \lambda) := \lambda^n + x_1\lambda^{n-1} + \dots + x_{n-1}\lambda + x_n, \quad x = (x_1, \dots, x_n),$$

$$a_i(x, \lambda) := \frac{1}{(n-i)!} \rho^{(n-i)}(x, \lambda)$$

$$= \binom{n}{i} \lambda^i + \binom{n-1}{i-1} x_1 \lambda^{i-1} + \dots + (n-(i-1)) x_{i-1} \lambda + x_i,$$

$$a'_i(x, \lambda) := (-1)^i a_i(x, \lambda), \quad t := -(\lambda + 2x_1/n).$$

Then we have the following lemma by long elementary calculation.

LEMMA 3.

$$\begin{aligned}
 a'_1(x, \lambda) &= a_1(x, t), \quad a'_2(x, \lambda) = a_2(x, t), \\
 a'_3(x, \lambda) &= a_3(x, t) + 2g_1(x_1, x_2, x_3, n), \\
 a'_4(x, \lambda) &= a_4(x, t) + 2(n-3)(t + x_1/n)g_1(x_1, x_2, x_3, n), \\
 a'_5(x, \lambda) &= a_5(x, t) + (n-3)(n-4)(t^2 + 2x_1t/n)g_1(x_1, x_2, x_3, n) \\
 &\quad + 2g_2(x_1, x_2, x_3, x_4, x_5, n), \\
 a'_6(x, \lambda) &= a_6(x, t) + (n-3)(n-4)\left(\frac{n-5}{3}t^3 + \frac{n-5}{n}x_1t^2 - \frac{1}{3n^2}x_1^2\right)g_1 \\
 &\quad \cdot (x_1, x_2, x_3, n) + 2(n-5)(t + x_1/n)g_2(x_1, x_2, x_3, x_4, x_5, n).
 \end{aligned}$$

In general,

$$\begin{aligned}
 a'_i(x, \lambda) &= a_i(x, t) + \sum_{j=0}^{i-3} \left(\sum_{k=0}^{i-j} (-1)^k \binom{n-k}{i-k} \binom{i-k}{i-j-k} \left(\frac{2}{n}x_1\right)^{i-j-k} \right. \\
 &\quad \left. \cdot x_k - \binom{n-i+j}{j} x_{i-j} \right) t^j \\
 &\quad (i = 1, \dots, n).
 \end{aligned}$$

Let

$$\begin{aligned}
 A_n &:= \{(x, y, z) \in S_0; a'_i(x, -2x_1/n) = x_i, \\
 &\quad a'_i(y, -2y_1/n) = y_i \ (i = 1, \dots, n)\}.
 \end{aligned}$$

(If $i = 1, 2$, $a'_i(x, -2x_1/n) = x_i$ and $a'_i(y, -2y_1/n) = y_i$ are always true.)
 Then $A_n = \{(x, y, z) \in S_0; g_1(x_1, x_2, x_3, n) = g_1(y_1, y_2, y_3, n) = 0\}$ for $n = 3, 4$ and $A_n = \{(x, y, z) \in S_0; g_1(x_1, x_2, x_3, n) = g_2(x_1, x_2, x_3, x_4, x_5, n) = g_1(y_1, y_2, y_3, n) = g_2(y_1, y_2, y_3, y_4, y_5, n) = 0\}$ for $n = 5, 6$. Thus if $n > 5$, we have $\text{codim } A_n > 4$.

Now we shall prove the rest of Theorem 2. For any $f(w) \in E_n$, there exists $(\alpha, \beta, \gamma) \in \tilde{S}_\Omega$ such that

$$f(w) = \gamma \exp 2(p_0\eta_1 + q_0\eta_2)w \prod_{i=1}^n \frac{\sigma(w - \alpha_i)}{\sigma(w - \beta_i)}, \quad \Omega = p_0\omega_1 + q_0\omega_2.$$

Let $B_n := \{(x, y, z) \in S; (1) \text{ and } (2) \text{ have no simple root}\}$, then B_n is an analytic set in S . If $(a, b, \gamma) := \phi(\alpha, \beta, \gamma) \in (S - B_n) \cup S_0$, we may assume that $(a, b, \gamma) \in S_0$. Then

$$f(w) = \gamma \prod_{i=1}^n \frac{\sigma(w - \alpha_i)}{\sigma(w - \beta_i)}, \quad a_1 = b_1.$$

For any $a(w) \in T_0$; $a(w) = w - \lambda$, we have

$$f \circ a(w) = \gamma \prod_{i=1}^n \frac{\sigma(w - \lambda - \alpha_i)}{\sigma(w - \lambda - \beta_i)}.$$

Hence $(\lambda + \alpha, \lambda + \beta, \gamma) \in \tilde{S}_0 \cap \tilde{F}^{-1}(f \circ a(w))$ and $\phi(\lambda + \alpha, \lambda + \beta, \gamma) = (a_1(a, \lambda), \dots, a_n(a, \lambda), a_1(b, \lambda), \dots, a_n(b, \lambda), \gamma) \in S_0 \cap F^{-1}(f \circ a(w))$. For any $a(w) \in T_1$; $a(w) = -w - \lambda$, we have

$$f \circ a(w) = \gamma \prod_{i=1}^n \frac{\sigma(w + \lambda + \alpha_i)}{\sigma(w + \lambda + \beta_i)}.$$

Hence $(-\lambda - \alpha, -\lambda - \beta, \gamma) \in \tilde{S}_0 \cap \tilde{F}^{-1}(f \circ a(w))$ and $\phi(-\lambda - \alpha, -\lambda - \beta, \gamma) = (a'_1(a, \lambda), \dots, a'_n(a, \lambda), a'_1(b, \lambda), \dots, a'_n(b, \lambda), \gamma) \in S_0 \cap F^{-1}(f \circ a(w))$. Namely the orbit $(\text{Aut } T)(f(w))$ is locally described by $\{(a_1(a, \lambda), \dots, a_n(a, \lambda), a_1(b, \lambda), \dots, a_n(b, \lambda), \gamma); \lambda \in T\} \cup \{(a'_1(a, \lambda), \dots, a'_n(a, \lambda), a'_1(b, \lambda), \dots, a'_n(b, \lambda), \gamma); \lambda \in T\}$ in S_0 . Hence using Lemma 3, we can show that $(\text{Aut } T)_{f(w)} = \{w\}$ if $(a, b, \gamma) \notin A_n$. Then $E_n/\text{Aut } T$ is nonsingular at $p(f(w))$ by the proof of Theorem 1. If $n \geq 5$, we have $\text{codim } A_n \geq 4$ and $\text{codim } B_n \geq n \geq 5$. Therefore $\text{codim } S(E_n/\text{Aut } T) \geq 3$ and so $E_n/\text{Aut } T$ is rigid.

In the case $\omega_2/\omega_1 \cong e^{\pi i/2}$ or $e^{2\pi i/3}$, we can also show that if $n \geq 5$, $\text{codim } S(E_n/\text{Aut } T) \geq 3$ by almost the same argument as above.

EXAMPLE. Let (w) be the Weierstrass pe -function. Then

$$\wp(w)^2 = \frac{1}{\sigma(\alpha)^4} \frac{\sigma(w - \alpha)^2 \sigma(w + \alpha)^2}{\sigma(w)^4} \in E_4(\wp(\alpha) = 0)$$

and $(\text{Aut } T)_{\wp(w)^2} = \{w, -w\}$. Thus $(\alpha, \alpha, -\alpha, -\alpha, 0, 0, 0, 0, 1/\sigma(\alpha)^4) \in \tilde{S}_0 \cap \tilde{F}^{-1}(\wp(w)^2)$ and $(0, -2\alpha^2, 0, \alpha^4, 0, 0, 0, 0, 1/\sigma(\alpha)^4) \in S_0 \cap F^{-1}(\wp(w)^2)$. By Lemma 3 and the proof of Theorems 1 and 2, we may take $N := \{x_1 = y_1 = 0\} \subset S_0$ and for any $(x, y, z) = (0, x_2, x_3, x_4, 0, y_2, y_3, y_4, z) \in N$, the orbit $(\text{Aut } T)(F(x, y, z))$ is locally described in S_0 as follows:

$$\begin{aligned} & \{(4\lambda, 6\lambda^2 + x_2, 4\lambda^3 + 2x_2\lambda + x_3, \lambda^4 + x_2\lambda^2 + x_3\lambda + x_4, 4\lambda, 6\lambda^2 + y_2, \\ & \quad 4\lambda^3 + 2y_2\lambda + y_3, \lambda^4 + y_2\lambda^2 + y_3\lambda + y_4, z); \lambda \in T\} \\ & \cup \{(-4\lambda, 6\lambda^2 + x_2, -4\lambda^3 - 2x_2\lambda - x_3, \lambda^4 + x_2\lambda^2 + x_3\lambda + x_4, -4\lambda, \\ & \quad 6\lambda^2 + y_2, -4\lambda^3 - 2y_2\lambda - y_3, \lambda^4 + y_2\lambda^2 + y_3\lambda + y_4, z); \lambda \in T\}. \end{aligned}$$

Hence $N_{(x,y,z)} = \{(0, x_2, x_3, x_4, 0, y_2, y_3, y_4, z), (0, x_2, -x_3, x_4, 0, y_2, -y_3, y_4, z)\}$. Therefore $E_4/\text{Aut } T$ is isomorphic to $\{w^2 - uv = 0\} \times \mathbf{C}^5 \subset \mathbf{C}^8$ at $p(\wp(w)^2)$ and so not rigid.

REFERENCES

1. D. Burns, *On rational singularities in dimension > 2* , Math. Ann. **211** (1974), 237–244.
2. H. Cartan, *Quotient d'un espace analytique par un groupe d'automorphismes*, Algebraic Geometry and Topology. A symposium in honor of S. Lefschetz, Princeton Univ. Press, Princeton, N. J., 1957, pp. 90–102.
3. H. Holmann, *Quotienten komplexer Räume*, Math. Ann. **142** (1961), 407–440.
4. ———, *Komplexe Räume mit komplexen Transformationsgruppen*, Math. Ann. **150** (1963), 327–360.
5. M. Namba, *Moduli of open holomorphic maps of compact complex manifolds*, Math. Ann. **220** (1976), 65–76.
6. D. Prill, *Local classification of quotients of complex manifolds by discontinuous groups*, Duke Math. J. **34** (1967), 375–386.
7. M. Schlessinger, *Rigidity of quotient singularities*, Invent. Math. **14** (1971), 17–26.
8. ———, *On rigid singularities*, Proc. Conf. on Singularities, Rice Univ. Studies, Houston, Tex., 1972, pp. 147–162.
9. K. Takijima and T. Suzuki, *On the trivial extension of equivalence relations on analytic spaces*, Trans. Amer. Math. Soc. **219** (1976), 369–377.

DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, SAITAMA UNIVERSITY, URAWA, JAPAN