PRODUCTS OF REFLECTIONS IN THE UNITARY GROUP

DRAGOMIR Ž. DJOKOVIĆ AND JERRY MALZAN

ABSTRACT. Let $A \in U(n)$, $\det(A) = \pm 1$ and let $\exp(i\alpha_k)$, $1 \leq k \leq n$ be the eigenvalues of A where $0 < \alpha_1 < \alpha_2 < \dots < \alpha_n < 2\pi$. Then $k(A) = (\alpha_1 + \dots + \alpha_n)/\pi$ is an integer and $0 < k(A) \leq 2n - 1$. Denote by $l(A)$ the length of A with respect to the set of all reflections, i.e., $l(A)$ is the smallest integer m such that A is a product of m reflections. A reflection is a matrix conjugate to $\text{diag}(-1, 1, \dots, 1)$. Our main result is the formula $l(A) = \max(k(A), k(A^*))$.

Introduction. By a reflection we mean here an element R of the unitary group $U(n)$ such that $R^2 = I_n$ and $\text{rank}(R - I_n) = 1$. We consider $U(n)$ as operating on the space of column vectors $\mathbb{C}^n$.

The reflections in $U(n)$ generate a normal subgroup $G(n)$ which consists of all matrices $A \in U(n)$ with $\det(A) = \pm 1$. It was shown by H. Radjavi [1] that in fact every element of $G(n)$ is a product of at most $2n - 1$ reflections and he conjectured that this bound is the best possible.

It is a simple consequence of our Theorem below that this conjecture is correct.

If $A = R_1 R_2 \dots R_m$ where R_k are reflections and m is minimal, then we say that m is the *length* of A and we denote it by $l(A)$.

For $A \in U(n)$ there exist uniquely determined real numbers $\alpha_1, \dots, \alpha_n$ satisfying $0 < \alpha_1 < \alpha_2 < \dots < \alpha_n < 2\pi$ and such that A is conjugate to the diagonal matrix

$$\begin{pmatrix} e^{i\alpha_1} & & & 0 \\ & \ddots & & \\ & & \ddots & \\ 0 & & & e^{i\alpha_n} \end{pmatrix} \quad (1)$$

We shall say that $\alpha_1, \dots, \alpha_n$ are the *angles* of A . The sum of the angles plays a very important role and we shall write

$$\alpha_1 + \dots + \alpha_n = k(A) \cdot \pi.$$

Note that $k(A)$ is an integer if and only if $A \in G(n)$.

In this paper we shall prove the following result.

THEOREM. If $A \in G(n)$ then $l(A) = \max(k(A), k(\bar{A}))$.

Received by the editors April 4, 1978.

AMS (MOS) subject classifications (1970). Primary 20G20, 22C05; Secondary 22E15.

Key words and phrases. Unitary group, reflections, length, eigenvalues, angles, characteristic polynomial.

© 1979 American Mathematical Society
0002-9939/79/0000-0053/\$02.00

For instance, if the angles $\alpha_1, \dots, \alpha_n$ of A satisfy $0 < \alpha_k < \pi$ ($1 \leq k \leq n$) and $\alpha_1 + \dots + \alpha_n = \pi$ then $k(A) = 1$, $k(\bar{A}) = 2n - 1$ and consequently $l(A) = 2n - 1$. Thus, Radjavi's conjecture is correct.

Proof of the Theorem. The proof will be given in a sequence of several lemmas.

If $a \in \mathbb{C}^n$ is a unit vector, i.e., $a^*a = 1$, then we denote by R_a the reflection $I_n - 2aa^*$. The hyperplane orthogonal to a is fixed pointwise by R_a and $R_a(a) = -a$.

LEMMA 1. *Let $A \in U(n)$ be diagonal with eigenvalues $\lambda_1, \dots, \lambda_n$. Then the characteristic polynomial $\phi(\lambda)$ of $R_a A$ is*

$$\phi(\lambda) = \left\{ \prod_{k=1}^n (\lambda - \lambda_k) \right\} \cdot \sum_{k=1}^n \frac{\lambda + \lambda_k}{\lambda - \lambda_k} |a_k|^2 \quad (2)$$

where the a_k are the coordinates of the unit vector a .

PROOF. We have

$$\begin{aligned} \phi(\lambda) &= \det(\lambda I_n - R_a A) = -\det(\lambda R_a - A) \\ &= -\det(\lambda I_n - A - 2\lambda a a^*) \\ &= -\det(\lambda I_n - A) \cdot \det(I_n - 2\lambda(\lambda I_n - A)^{-1} a a^*). \end{aligned}$$

Since $(\lambda I_n - A)^{-1} a a^*$ has rank 1, its eigenvalues are 0 (with multiplicity $n - 1$), and $a^*(\lambda I_n - A)^{-1} a$. Therefore

$$\phi(\lambda) = -\det(\lambda I_n - A) \cdot (1 - 2\lambda a^*(\lambda I_n - A)^{-1} a).$$

Since $|a_1|^2 + \dots + |a_n|^2 = 1$, this is equivalent to the formula (2).

LEMMA 2. *Let $A \in U(n)$ have the angles $\alpha_1, \dots, \alpha_n$. If $R \in U(n)$ is a reflection and if $\beta_1, \dots, \beta_n$ are the angles of RA , then*

$$\beta_1 + \dots + \beta_n = \alpha_1 + \dots + \alpha_n \pm \pi.$$

PROOF. Let $\lambda_1, \dots, \lambda_m$ be the distinct eigenvalues of A . Let V_k be the λ_k -eigenspace of A . Finally, let H be the hyperplane of R -fixed vectors and $W_k = V_k \cap H$. It is clear that $\dim(V_k/W_k) = 0$ or 1 for each k . The subspace $W = W_1 + \dots + W_m$ is invariant under both R and A . Hence $L = W^\perp$ is also invariant under R and A . Let R_0 and A_0 be their respective restrictions to L . It is clear that all eigenvalues of A_0 are simple and that it suffices to prove the assertion for R_0 and A_0 instead of R and A . In other words we may assume now that A has only simple eigenvalues.

Without loss of generality we may assume that A is given by (1). Let a be a unit vector such that $R = R_a = I_n - 2aa^*$. By conjugating both A and R by a suitable permutation matrix, we may assume that the first m coordinates $a_1, \dots, a_m$ of a are nonzero while $a_{m+1} = \dots = a_n = 0$. Consequently A and RA have the angles $\alpha_{m+1}, \dots, \alpha_n$ in common. Hence it suffices to prove the lemma in the case when $m = n$, i.e., when all $a_k \neq 0$.

Now let $\lambda_k = e^{i\alpha_k}$, $1 \leq k \leq n$. By Lemma 1, the characteristic polynomial

$\phi(\lambda)$ of RA is given by (2). Putting $\lambda = e^{i\theta}$ the equation $\phi(\lambda) = 0$ becomes

$$\sum_{k=1}^n |a_k|^2 \cotan \frac{1}{2}(\theta - \alpha_k) = 0. \quad (3)$$

Now we may assume that $0 \leq \alpha_1 < \alpha_2 < \cdots < \alpha_n < 2\pi$. It is immediate from (3) that there is at least one root of (3) in each of the intervals (α_k, α_{k+1}) for $1 \leq k \leq n-1$ and another one in $(\alpha_n, \alpha_1 + 2\pi)$. Hence the angles $\beta_1, \dots, \beta_n$ of RA ordered so that $0 \leq \beta_1 < \beta_2 < \cdots < \beta_n < 2\pi$ will satisfy either

$$0 \leq \beta_1 < \alpha_1 < \beta_2 < \alpha_2 < \cdots < \beta_n < \alpha_n \quad (4)$$

or

$$\alpha_1 < \beta_1 < \alpha_2 < \cdots < \beta_{n-1} > \alpha_n < \beta_n < 2\pi. \quad (5)$$

In case (4) we have

$$\alpha_1 + \cdots + \alpha_n > \beta_1 + \cdots + \beta_n < \alpha_1 + \cdots + \alpha_n - 2\pi$$

and in case (5) we have

$$\alpha_1 + \cdots + \alpha_n < \beta_1 + \cdots + \beta_n < \alpha_1 + \cdots + \alpha_n + 2\pi.$$

Since $\det(RA) = -\det A$ the difference

$$(\beta_1 + \cdots + \beta_n) - (\alpha_1 + \cdots + \alpha_n)$$

must be an odd multiple of π and the proof is completed.

For $A \in G(n)$ let $k'(A) = \max(k(A), k(\bar{A}))$.

LEMMA 3. *If $A \in G(n)$ and R is a reflection then $k'(RA) = k'(A) \pm 1$. Consequently, $l(A) \geq k'(A)$.*

PROOF. Let $r = \text{rank}(A - I_n)$ and let $\alpha_1, \dots, \alpha_r$ be the nonzero angles of A . Then the nonzero angles of $\bar{A}$ are $2\pi - \alpha_1, \dots, 2\pi - \alpha_r$. Thus $k(A) + k(\bar{A}) = 2r$, and so the difference $k(A) - k(\bar{A})$ is even. Therefore it suffices to prove that $k(RA) = k(A) \pm 1$. But this last claim is immediate from Lemma 2.

LEMMA 4. *Let α, β be two angles of $A \in U(n)$ such that $0 < \alpha < \pi < \beta < 2\pi$. Then there exists a reflection R such that the angles of RA are the same as those of A except that α and β are replaced by 0 and $\alpha + \beta - \pi$.*

PROOF. Clearly, it suffices to consider the case when $n = 2$. We need only to show that we can choose a reflection $R = R_a$ so that RA has 1 as eigenvalue. The equation (3) will be satisfied by $\theta = 0$ if and only if

$$|a_1|^2 \cotan \frac{\alpha}{2} + |a_2|^2 \cotan \frac{\beta}{2} = 0.$$

One can choose a_1 and a_2 nonzero satisfying this equation because the two cotangents have opposite signs.

LEMMA 5. *If $A \in G(n)$ and $k'(A) \geq 1$, there exists a reflection R such that $k'(RA) = k'(A) - 1$.*

PROOF. We may assume that A is given by (1) where $\alpha_1, \dots, \alpha_n$ are the angles of A in increasing order. If some $\alpha_k = \pi$ it suffices to take $R = I_n - 2e_k e_k^*$, e_k being the standard basic vector of $\mathbb{C}^n$.

If $\alpha_n < \pi$ we can take $R = I_n - 2e_n e_n^*$.

If $\alpha_1 > \pi$ we can take $R = I_n - 2e_1 e_1^*$.

Finally, if $0 < \alpha_k < \pi < \alpha_r < 2\pi$ for some k, r then we can apply Lemma 4.

The assertion of the Theorem follows from Lemmas 3 and 5.

REFERENCES

1. H. Radjavi, *Decomposition of matrices into simple involutions*, Linear Algebra and Appl. **12** (1975), 247–255.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA N2L 3G1 (current address of D. Ž. Djoković)

Current address (Jerry Malzan): Department of Mathematics, University of Toronto, Toronto, Ontario, Canada M5S 1A1