ON A THEOREM OF DUREN, SHAPIRO AND SHIELDS

SHINJI YAMASHITA

ABSTRACT. We shall show that, an extension of the theorem of Duren, Shapiro and Shields on the univalence of a function f holomorphic in the unit disk D , still remains of significance, if the power $\alpha \in (0, 1)$ in

$$\sup_{z \in D} (1 - |z|^2)^\alpha |f''(z)/f'(z)|$$

is small enough.

Let f be a function nonconstant and holomorphic in $D = \{ |z| < 1 \}$, let $\alpha > 0$, and let

$$\|f\|_\alpha = \sup_{z \in D} (1 - |z|^2)^\alpha |f''(z)/f'(z)|.$$

It was proved by P. L. Duren, H. S. Shapiro and A. L. Shields [3, Theorem 2] that if

$$\|f\|_1 \leq 2(\sqrt{5} - 2) = 0.47 \dots,$$

then f is univalent in D . This was improved later by J. Becker [1, Corollary 4.1, p. 36] in the sense that if $\|f\|_1 \leq 1$, then f is univalent in D . Therefore, if $\alpha \in [0, 1]$, and if $\|f\|_\alpha \leq 1$, then f is univalent in D because $\|f\|_\alpha \geq \|f\|_1$.

We shall show, nevertheless, the method found by Duren, Shapiro and Shields still remains available for small $\alpha > 0$.

THEOREM. *Let f be a function nonconstant and holomorphic in D . Suppose that $\alpha \in [0, 1]$, and suppose that*

$$\|f\|_\alpha \leq 2(\sqrt{4^\alpha + 2 - \alpha} - 2^\alpha) \equiv k_\alpha.$$

Then f is univalent in D .

Our theorem is thus significant for $k_\alpha > 1$; this is the case for $0 < \alpha < \alpha_0$, where α_0 is the constant satisfying $2^{\alpha_0} + \alpha_0 = 7/4$. We note that $\alpha_0 = 0.416 \dots$; the detailed calculation is due to the referee. The constant k_1 is the constant found by Duren, Shapiro and Shields.

We remark that the constant $k_0 = 2(\sqrt{3} - 1) = 1.46 \dots$ is not good. In effect, it is known that if $\|f\|_0 \leq 2\sqrt{2} = 2.82 \dots$, then f is univalent in D ; see [4, Problem 2, p. 179] (this Problem 2 is solved on making use of the Schwarz and Pick lemma, together with the theorem of V. V. Pokornyi, cited in [4, Problem 1, p. 179]).

Received by the editors March 15, 1978.

AMS (MOS) subject classifications (1970). Primary 30A36.

© 1979 American Mathematical Society
0002-9939/79/0000-0058/\$01.75

We begin with

LEMMA. Let g be a function holomorphic in D , and let $\alpha > 0$. Then

$$\sup_{z \in D} (1 - |z|^2)^{1+\alpha} |g'(z)| \leq 2^{1+\alpha} \sup_{z \in D} (1 - |z|^2)^\alpha |g(z)|. \quad (1)$$

PROOF. We may assume that the supremum in the right-hand side of (1), denoted by M , is finite, because the case where $M = +\infty$ is obvious. For each $z \in D$, we let

$$\rho = \left[\frac{1}{2} (1 + r^2) \right]^{1/2}, \quad r = |z|.$$

Then

$$(1 - r^2)^{1+\alpha} = 2^{1+\alpha} (\rho^2 - r^2) (1 - \rho^2)^\alpha.$$

Since

$$g'(z) = (2\pi i)^{-1} \int_{|w|=\rho} g(w) (w - z)^{-2} dw,$$

it follows that

$$\begin{aligned} (1 - |z|^2)^{1+\alpha} |g'(z)| &\leq 2^{1+\alpha} (\rho^2 - r^2) (1 - \rho^2)^\alpha (\sup_{|w|=\rho} |g(w)|) \\ &\quad \times \left[(2\pi)^{-1} \int_0^{2\pi} |\rho e^{i\theta} - z|^{-2} \rho d\theta \right] \\ &\leq 2^{1+\alpha} M \left[(2\pi)^{-1} \int_0^{2\pi} (\rho^2 - r^2) |\rho e^{i\theta} - z|^{-2} d\theta \right] = 2^{1+\alpha} M. \end{aligned}$$

We thus obtain (1).

PROOF OF THEOREM. It follows from the Lemma, applied to $g = f''/f'$, that

$$(1 - |z|^2)^{1+\alpha} |(f''(z)/f'(z))'| \leq 2^{1+\alpha} \|f\|_\alpha \leq 2^{1+\alpha} k_\alpha, \quad z \in D.$$

On the other hand, it follows from $1 + \alpha \geq 2\alpha$, that

$$(1 - |z|^2)^{1+\alpha} |f''(z)/f'(z)|^2 \leq \|f\|_\alpha^2 \leq k_\alpha^2, \quad z \in D.$$

Therefore, we obtain

$$(1 - |z|^2)^{1+\alpha} |S_f(z)| \leq 2^{1+\alpha} k_\alpha + \frac{1}{2} k_\alpha^2 = 2(2 - \alpha), \quad z \in D, \quad (2)$$

where $S_f = (f''/f')' - \frac{1}{2} (f''/f')^2$ is the Schwarzian derivative of f .

Now, P. R. Beesack [2, (2.6), p. 217, and (2.8), p. 218] proved that if there exists $\lambda \in [1, 2]$ such that

$$(1 - |z|^2)^\lambda |S_f(z)| \leq 2(3 - \lambda), \quad z \in D,$$

then f is univalent in D . Since $1 \leq 1 + \alpha \leq 2$, it follows from (2) that f is univalent in D .

REFERENCES

1. Jochen Becker, *Löwnersche Differentialgleichung und quasikonform fortsetzbare schlichte Funktionen*, J. Reine Angew. Math. **255** (1972), 23–43.

2. Paul R. Beesack, *Nonoscillation and disconjugacy in the complex domain*, Trans. Amer. Math. Soc. **81** (1956), 211–242.
3. Peter L. Duren, Harold S. Shapiro and Allen L. Shields, *Singular measures and domains not of Smirnov type*, Duke Math. J. **33** (1966), 247–254.
4. Christian Pommerenke, *Univalent functions*, Vandenhoeck und Ruprecht, Göttingen, 1975.

DEPARTMENT OF MATHEMATICS, TOKYO METROPOLITAN UNIVERSITY, FUKAZAWA,
SETAGAYA-KU, TOKYO 158, JAPAN