EMBEDDING OF A LIE ALGEBRA INTO LIE-ADMISSIBLE ALGEBRAS

HYO CHUL MYUNG

ABSTRACT. Let A be a flexible Lie-admissible algebra over a field of characteristic $\neq 2, 3$. Let S be a finite-dimensional classical Lie subalgebra of A^- which is complemented by an ideal R of A^- . It is shown that S is a Lie algebra under the multiplication in A and is an ideal of A if and only if S contains a classical Cartan subalgebra H which is nil in A and such that $HH \subseteq S$ and $[H, R] = 0$. In this case, the multiplication between S and R is determined by linear functionals on R which vanish on $[R, R]$. If A is finite-dimensional and of characteristic 0 then this can be applied to give a condition that a Levi-factor S of A^- be embedded as an ideal into A and to determine the multiplication between S and the solvable radical of A^- .

1. Introduction. For an algebra A , denote by A^- the algebra with multiplication $[x, y] = xy - yx$ defined on the vector space A . If A^- is a Lie algebra then A is said to be Lie-admissible. If A satisfies the flexible law $(xy)x = x(yx)$ for all $x, y \in A$ then A is called a flexible algebra. A linearized form of the flexible law is $(xy)z - x(yz) + (zy)x - z(yx) = 0$. If A is flexible and Lie-admissible then it is well known that the mapping $\text{ad } x: a \rightarrow [a, x]$ is a derivation of A for all $x \in A$; that is, $[a, bc] = [a, b]c + b[a, c]$ for all $a, b, c \in A$. Recent results show that a Cartan subalgebra of A^- plays a major role for the structure of flexible Lie-admissible algebras A [1], [2], [3]. Possible applications of Lie-admissible algebras in physics have been recently pointed out by a number of physicists. For this, the reader is referred to Santilli's recent work [5].

The purpose of this paper is to give a condition in terms of a Cartan subalgebra that a classical Lie algebra S be embedded as an ideal into a flexible Lie-admissible algebra A when the subspace S is complemented by an ideal R of A^- , and then to determine the multiplication between S and R . We make use of the known structure for classical Lie algebras [4] and the result that if A is a flexible algebra and A^- is a classical Lie algebra with a classical Cartan subalgebra which is nil in A , then A is a Lie algebra isomorphic to A^- [2]. A subset M of A said to be nil in A if every element of M is power-associative and nilpotent in A .

Received by the editors March 27, 1978.

AMS (MOS) subject classifications (1970). Primary 17A20, 17B05.

Key words and phrases. Flexible algebra, Lie-admissible algebra, classical Lie algebra, Cartan subalgebra, Levi-factor.

© 1979 American Mathematical Society
0002-9939/79/0000-0103/\$02.25

Let S be the direct sum of Lie algebras $S_1, \dots, S_n$ over a field Φ of characteristic $\neq 2$ and R be a flexible Lie-admissible algebra over Φ . Let $f_1, \dots, f_n$ be linear functionals on R which vanish on $[R, R]$. Define a multiplication in the vector space direct sum $A = S + R$ as follows:

$$\left(\sum_{i=1}^n x_i + r \right) \left(\sum_{i=1}^n y_i + s \right) = \sum_{i=1}^n [x_i y_i + f_i(r) y_i + f_i(s) x_i] + rs,$$

$$x_i, y_i \in S_i, r, s \in R, i = 1, 2, \dots, n. (*)$$

Then one sees that $[x + r, y + s] = 2xy + [r, s]$ for $x, y \in S$ and $r, s \in R$ and so A is Lie-admissible. One also computes

$$[(x + r)(y + s)](x + r) - (x + r)[(y + s)(x + r)] = \sum_i f_i([r, s]) x_i = 0,$$

where $x = \sum_i x_i$, since $f_i([R, R]) = 0$. Thus A is flexible. Clearly, R is a subalgebra of A and is an ideal of A^- . The Lie algebra S is embedded as an ideal in A and if $x \in S_i$ then $xr = rx = f_i(r)x$ for $r \in R$. In fact, we show that this is the essential source for the embedding in question. We state the main result as follows.

THEOREM 1. *Let A be a flexible Lie-admissible algebra over a field Φ of characteristic $\neq 2, 3$ (not necessarily finite-dimensional). Let S be a finite-dimensional classical subalgebra of A^- which is complemented by an ideal R of A^- . Then S is a Lie algebra under the multiplication in A and is an ideal of A if and only if S contains a classical Cartan subalgebra H which is nil in A and such that $HH \subseteq S$ and $[H, R] = 0$. In this case, R is a subalgebra of A and the multiplication in A is given by the rule $(*)$ where $f_1, \dots, f_n$ are linear functionals on R which vanish on $[R, R]$, and n is the number of simple summands in S .*

Recall that a finite-dimensional classical Lie algebra is a direct sum of simple Lie algebras [4].

2. Proof of Theorem 1. We begin with the following lemma.

LEMMA. *Let L be a Lie algebra over an arbitrary field Φ . Let S be a finite-dimensional subalgebra of L and H be a Cartan subalgebra of S . Then, for an ideal R of L , $[SR] = 0$ if and only if $[HR] = 0$.*

PROOF. One may assume that Φ is algebraically closed; if not, one takes the scalar extension of S to the algebraic closure of Φ . Let $S = \sum_{\alpha} S_{\alpha}$ be the Cartan decomposition for S relative to H . For each nonzero root α , choose an $h \in H$ such that $\alpha(h) \neq 0$. Then $\text{ad } h: S_{\alpha} \rightarrow S_{\alpha}$ is surjective; for, if not then there is an element $x \neq 0$ in S_{α} such that $[x, h] = 0$ and this together with $x(\text{ad } h - \alpha(h)I)^n = 0$ implies that $\alpha(h)^n x = 0$. This is absurd. Thus we have that $[S_{\alpha} h] = S_{\alpha}$ for $\alpha \neq 0$ and $\alpha(h) \neq 0$. If $[HR] = 0$ and $\alpha \neq 0$ then, by the Jacobi identity, $[S_{\alpha} R] = [[S_{\alpha} h] R] \subseteq [[Rh] S_{\alpha}] + [[S_{\alpha} R] h] = 0$ since R is an ideal of L , and so $[SR] = 0$.

For the proof of Theorem 1, we first observe that the centralizer $C_{A^-}(M)$ of

a subset M of A in A^- is a subalgebra of A , since $[xy, M] \subseteq x[y, M] + [x, M]y = 0$ for all $x, y \in C(M)$. Suppose that S is a finite-dimensional classical subalgebra of A^- having a classical Cartan subalgebra H which is nil in A and that $HH \subseteq S$ and $[H, R] = 0$. That $HH \subseteq S$ implies that S is a subalgebra of A [3]. It then follows from [2, Corollary 3.4] that S is a Lie algebra under the multiplication in A . Thus we have that $[x, y] = xy - yx = 2xy$ for $x, y \in S$. By the Lemma, we get $[S, R] = 0$ and since S has center 0, $R = C_{A^-}(S)$ and so R is a subalgebra of A .

Let $x, y \in S$ and let $r \in R$. Write $yr = z + s$ for some $z \in S$ and $s \in R$. Then the flexible law $(xy)r - x(yr) + (ry)x - r(yx) = 0$ implies $(xy)r = xz$ and so $(SS)R \subseteq S$. Since $SS = S$, this proves that S is an ideal of A . Since $C_S(H) = H$ and $HH = [H, H] = 0$, from $[h', hr] = [h', h]r + h[h', r] = 0$ for all $h, h' \in H$ and $r \in R$, we have

$$HR = RH \subseteq H. \quad (1)$$

Let $S = \sum_{\alpha} S_{\alpha}$ be the Cartan decomposition for S relative to H . Note that $S_0 = H$ and $\dim S_{\alpha} = 1$ for $\alpha \neq 0$ since S is classical. Then $xh = \alpha(h)x$ for $x \in S_{\alpha}$ and $h \in H$. For each nonzero root α , choose an $h \in H$ with $\alpha(h) \neq 0$. If $x \in S_{\alpha}$ and $r \in R$ then, from $(xh)r - x(hr) + (rh)x - r(hx) = 0$ and (1), one gets $xr = \alpha(h)^{-1}\alpha(hr)x$ and so

$$xr = rx = \lambda x, \quad x \in S_{\alpha}, \quad \alpha \neq 0, \quad (2)$$

where $\lambda \in \Phi$ depends on $r \in R$ and $\alpha \neq 0$. Since $S_{\alpha}S_{-\alpha}$ is one-dimensional for $\alpha \neq 0$, one chooses nonzero elements $x \in S_{\alpha}, y \in S_{-\alpha}, h \in H$ such that

$$xh = x, \quad yh = -y, \quad xy = h. \quad (3)$$

Then by (2) we have that $xr = rx = \lambda x$ and $yr = ry = \mu y$ for $\mu \in \Phi$ and $r \in R$. If $H = \Phi x + B$ is a vector space direct sum then by (1) we can let $hr = rh = \nu h + b$, $\nu \in \Phi$, $b \in B$. From (3) and $(xy)r - x(yr) + (ry)x - r(yx) = 0$, one gets $2(\nu h + b) = 2\mu h$, so $b = 0$ and $\nu = \mu$, and by symmetry $\nu = \lambda$. Therefore

$$xr = rx = f_{\alpha}(r)x, \quad yr = ry = f_{\alpha}(r)y, \quad hr = rh = f_{\alpha}(r)h \quad (4)$$

for $\alpha \neq 0$ and $r \in R$, where f_{α} is a linear functional on R and $\{x, y, h\}$ is the canonical basis as in (3). In particular, we have that $f_{\alpha} = f_{-\alpha}$.

Recall that S is the direct sum of simple classical Lie algebras, so that each simple summand has a fundamental system of roots relative to a classical Cartan subalgebra which is connected. Thus we may assume that S is simple. If α, β ($\beta \neq 0$) is an ordered pair of roots then recall the Cartan integer $A_{\alpha, \beta} = r - q$ where r and q are the least nonnegative integers such that $\alpha - (r + 1)\beta$ and $\alpha + (q + 1)\beta$ are not roots. Let $\Pi = \{\alpha_1, \dots, \alpha_m\}$ be a fundamental system of roots which is connected; that is, for any two roots $\alpha, \beta \in \Pi$, there are roots $\mu_1, \dots, \mu_r \in \Pi$ such that $\alpha = \mu_1, \beta = \mu_r$ and $A_{\mu_i, \mu_{i+1}} \neq 0$, $1 \leq i \leq r$. For brevity, denote $A_{\alpha, \alpha_j} = A_{ij}$ and $f_{\alpha_i} = f_i$. We first show that if $A_{ij} \neq 0$ then $f_i = f_j$. If $A_{ij} < 0$ then $S_{\alpha_i}S_{\alpha_j} \neq 0$ and so choose elements

$x \in S_{\alpha}, y \in S_{\alpha}$ with $xy \neq 0$. Then the flexible law $(xy)r - x(yr) + x(yr) - r(yx) = 0$ with (4) implies that $(xy)r = f_j(r)(xy)$ for all $r \in R$. By symmetry we have that $(xy)r = f_i(r)(xy)$ and so $f_i = f_j$. If $A_{ij} > 0$ then, using the relation $A_{\alpha, -\beta} = -A_{\alpha, \beta}$ and $f_{\alpha} = f_{-\alpha}$, we argue that $f_i = f_j$. Since Π is connected, this proves that $f_i = f_j$ for all $i, j = 1, 2, \dots, m$. Denote one of these linear functionals f_i by f .

Let x_i, y_i, h_i be the canonical basis corresponding to the root $\alpha_i \in \Pi$. We have then shown

$$x_i r = r x_i = f(r) x_i, \quad y_i r = r y_i = f(r) y_i, \quad h_i r = r h_i = f(r) h_i, \\ r \in R, i = 1, 2, \dots, m. \quad (5)$$

Since, for any root $\alpha \neq 0$, α or $-\alpha$ is a sum of roots in Π , there is a basis for S which consists of elements of the form

$$h_i, \quad (\dots (x_{i_1} x_{i_2}) \dots x_{i_k}), \quad (\dots (y_{i_1} y_{i_2}) \dots y_{i_k}), \\ i = 1, 2, \dots, m, \{i_1, \dots, i_k\} \subseteq \{1, 2, \dots, m\}.$$

The flexible law $(x_i x_j) r - x_i (x_j r) + (r x_j) x_i - r (x_j x_i) = 0$ gives $(x_i x_j) r = f(r)(x_i x_j)$ by (5). Therefore, by induction, we have

$$x r = r x = f(r) x, \quad x \in S, r \in R.$$

Finally, let $r, s \in R$ and let x be a nonzero element of S . Then $(rs)x - r(sx) + (xs)r - x(sr) = 0$ gives $f([r, s]) = 0$ and so f vanishes on $[R, R]$. Thus the multiplication in A is given by (*) and this completes the proof.

3. Application. Let A be a finite-dimensional flexible Lie-admissible algebra over a field of characteristic 0. Then, by Levi's theorem, every Levi-factor S (a maximal semisimple subalgebra of A^-) of A^- is complemented by the solvable radical of A^- . It then follows from Corollary of [3] that if S is power-associative in A and contains a Cartan subalgebra H with $HH \subseteq S$ then S is a subalgebra of A and is a Lie algebra under the multiplication in A . Therefore, in view of Theorem 1, we have

THEOREM 2. *Let A be a finite-dimensional flexible Lie-admissible algebra over a field of characteristic 0. Let R be the solvable radical of A^- and S be a Levi-factor of A^- which is power-associative in A . Then S is a Lie algebra under the multiplication in A and is an ideal of A if and only if S contains a split Cartan subalgebra H such that $HH \subseteq S$ and $[H, R] = 0$. In this case, the multiplication in A is given by (*).*

Theorem 2 strengthens Theorem 4.1 in [2] which requires the additional assumptions that the radical R of A^- is nilpotent in A^- and H is nil in A . If, in Theorem 1, the complementary subalgebra R of A satisfies $[R, R] = R$, then the linear functionals f_i are 0. Thus we have

COROLLARY 1. *Let A, S, R be the same as in Theorem 1 and let S be embedded as in Theorem 1. If $[R, R] = R$ then R is an ideal of A . In particular, if R^- is a simple Lie algebra then R is an ideal of A .*

COROLLARY 2. *Let A, S, R be the same as in Corollary 1 and of characteristic 0. If R^- is nilpotent and R is nil in A then R is an ideal of A .*

If R^- is nilpotent then $\text{ad } x$ is nilpotent in A for all $x \in R$. Since R is a nilalgebra, it follows from [2, Lemma 4.4] that the right multiplication $R(x)$ in A by $x \in R$ is nilpotent. Hence the linear functionals f_i are 0 and this proves Corollary 2.

REFERENCES

1. P. J. Laufer and M. L. Tomber, *Some Lie admissible algebras*, Canad. J. Math. **14** (1962), 287–292.
2. H. C. Myung, *Some classes of flexible Lie-admissible algebras*, Trans. Amer. Math. Soc. **167** (1972), 79–88.
3. ———, *A subalgebra condition in Lie-admissible algebras*, Proc. Amer. Math. Soc. **59** (1976), 6–8.
4. G. B. Seligman, *Modular Lie algebras*, Ergebnisse der Math. und ihrer Grenzgebiete, Band 40, Springer-Verlag, New York, 1967.
5. R. M. Santilli, *Lie-admissible approach to the hadronic structure*, Volumes I, II, and III, Hadronic Press, Inc., Nonantum, Mass. (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTHERN IOWA, CEDAR FALLS, IOWA 50613