AN ALGEBRA OF FUNCTIONS

BJÖRN E. J. DAHLBERG

ABSTRACT. It is established that the space of logarithmic potentials of H^1 -functions is closed under multiplication.

In this note we shall establish that the product of two functions, both of which are logarithmic potentials of H^1 -functions, is the logarithmic potential of a function in H^1 . Here $H^1 = H^1(R^n)$ is defined as the class of those L^1 -functions f on R^n for which it is possible to find L^1 -functions $g_1, \dots, g_n$ such that $\hat{g}_j(\xi) = \xi_j |\xi|^{-1} \hat{f}(\xi)$, where $\hat{f}$ denotes the Fourier transform of f . The H^1 -norm of f is then

$$\|f\| = \|f\|_1 + \sum_1^n \|g_j\|_1.$$

For properties of H^1 see Stein [5] and Fefferman and Stein [3].

We define the logarithmic potential of a function f as

$$Kf(x) = \int_{R^n} f(y) \log|x - y| dy.$$

We can now state our main result.

THEOREM. *If $f, g \in H^1$ then there is a function $h \in H^1$ such that $KfKg = Kh$. Moreover, there is a constant C independent of f and g such that $\|h\| \leq C\|f\| \|g\|$.*

We remark that in the case $n = 1$ this type of algebra has been studied by Coifman and Weiss [2, p. 601] in connection with the algebra of Calderon-Zygmund operators on R^2 with characteristic in $H^1(T)$.

For other results when spaces of potentials are closed under multiplication see Strichartz [6], [7], Bagby [1].

The proof of Theorem 1 is based on the atomic decomposition of H^1 -functions. We recall that a function $a \in L^\infty(R^n)$ is called an atom if there is a cube I with its sides parallel to the coordinate axis such that a is supported on I , $\int_{R^n} a dx = 0$, and $\|a\|_\infty \leq |I|^{-1}$. Here $|E|$ denotes the Lebesgue measure of a set $E \subset R^n$.

It is now known that given $f \in H^1$ there is a sequence of atoms a_j and

Received by the editors April 21, 1978.

AMS (MOS) subject classifications (1970). Primary 31B99.

Key words and phrases. Logarithmic potentials, H^1 -functions, atoms, Sobolev inequalities.

© 1979 American Mathematical Society
0002-9939/79/0000-0157/\$02.25

numbers λ_j such that

$$f = \sum \lambda_j a_j \quad (1)$$

and $\sum |\lambda_j| \leq C \|f\|$ for some fixed constant C . Also, whenever the decomposition (1) holds $\|f\| \leq C \sum |\lambda_j|$ for some fixed constant C . For a full discussion of these results see Coifman and Weiss [2]. From these results it follows that in order to prove the theorem it is sufficient to show that if a and b are two atoms then $KaKb = Kh$ for some $h \in H^1$ such that $\|h\| \leq C$, where C only depends on n .

We recall that a function ϕ is said to have bounded mean oscillation if for the family of cubes I we have

$$\sup |I|^{-1} \int_I |\phi - \phi_I| dx = \|\phi\|_* < \infty,$$

where $\phi_I = |I|^{-1} \int_I \phi dx$. The space of all such functions is denoted by BMO. An important fact is that the dual of H^1 is BMO; see Fefferman and Stein [3]. We shall need the following estimate, whose proof is a straightforward modification of inequality (1.2) in Fefferman and Stein [3]. If $\phi \in \text{BMO}$ and $1 \leq p < \infty$ then for any cube I centered at 0 we have

$$\left(\int \alpha(\alpha + |x|)^{-1-n} |\phi(x) - \phi_I|^p dx \right)^{1/p} \leq C \|\phi\|_*, \quad (2)$$

where α is the side of I and C is independent of ϕ .

For a cube I we denote by $2I$ the cube with the same center as I but with twice the side. With the support cube of an atom a we mean the smallest (axis parallel) cube which contains the support of a .

LEMMA. *Suppose a is an atom, the support cube of which is centered at 0. There is a constant C only depending on n such that if $x \notin 2I$ and $0 \leq j \leq n$ then*

$$|\nabla_j Ka(x)| \leq C\alpha(|x| + \alpha)^{-1-j}.$$

If $x \in 2I$ and $0 \leq j \leq n-1$ then

$$|\nabla_j Ka(x)| \leq C\alpha^{-j}.$$

Here α is the side of I and $\nabla_j h$ denotes the partial derivatives of h of order j arranged in some order.

PROOF. Let $F_j(x) = \nabla_j \log|x|$. Since $|F_j(x)| \leq C_j|x|^{-j}$, $1 \leq j$, it follows that if $x \notin 2I$, then

$$\begin{aligned} \nabla_j Ka(x) &= \int_I a(y) F_j(x-y) dy \\ &= \int_I a(y) (F_j(x-y) - F_j(x)) dy. \end{aligned}$$

Hence $|a_j Ka(x)| \leq C_j|x|^{-j-1}$ which shows the first part of the lemma. Since

$\|a\| \leq C$ and $\log |x| \in \text{BMO}$ (see John and Nirenberg [4]) it follows from the duality between H^1 and BMO that

$$\|Ka\|_\infty \leq C. \quad (3)$$

If $1 \leq j \leq n-1$ and $x \in 2I$ then

$$\begin{aligned} |\nabla_j Ka(x)| &\leq \int |F_j(x-y)| |a(y)| dy \\ &\leq C_j \int_{|x| \leq \alpha} |y|^{-j} |I|^{-1} dy \leq C_j \alpha^{-j}, \end{aligned}$$

which completes the proof of the lemma.

We shall now assume that a and b are atoms with support cubes I and J respectively. We shall assume that a and b are C^∞ and we shall next show that there is a constant C , only depending on n , such that for all $\phi \in \text{BMO}$ we have

$$\left| \int_{R^n} \phi D^n (KaKb) dx \right| \leq C \|\phi\|_*. \quad (4)$$

Here D denotes differentiation in the direction of a unit vector e .

We let α and β denote the sides of I and J respectively. It is no loss in generality in assuming $\alpha \leq \beta$ and I is centered at the origin. Fix j , $0 \leq j \leq n$, and put

$$A_j = \int_{R^{n-2I}} (\phi - \phi_{2I}) D^j Ka D^{n-j} Kb.$$

Defining $p = p_j = (n+1)(j+1)^{-1}$ we have

$$1 - p^{-1} = n(p^{-1} - kn^{-1}). \quad (5)$$

From the lemma, (2) and Hölder's inequality we get

$$\begin{aligned} |A_j| &\leq C \int_{R^{n-2I}} \alpha(|x| + \alpha)^{-j-1} |\phi(x) - \phi_{2I}| |D^{n-j} Kb(x)| dx \\ &\leq C \alpha^{1-1/p} \|\phi\|_* \|D^{n-j} Kb\|_q, \end{aligned}$$

where $q = p/p-1$. If $j = 0$ then $1 < q < \infty$ and since

$$\widehat{D^n Kb}(\xi) = \langle e, \xi \rangle^n |\xi|^{-n} \hat{b}(\xi)$$

it follows from well-known results on multipliers (see Stein [5]) that $\|D^n Kb\|_q \leq C \|b\|_q \leq C \beta^{-n(1-q^{-1})}$. Here $\langle e, \xi \rangle$ denotes the scalar product. Remembering (5) and the assumption $\alpha \leq \beta$ we see that $|A_0| \leq C$. If $1 \leq j \leq n-1$ we observe that

$$|D^{n-j} Kb| \leq C I_j |b|,$$

where $I_j f(x) = \int f(y) |x-y|^{-n} dy$. It is known (see Stein [5, Chapter V]) that if r is defined by $q^{-1} = r^{-1} - jn^{-1}$ then $\|I_j f\|_q \leq C \|f\|_r$. Hence $\|D^{n-j} Kb\|_q \leq C \beta^{-n(1-r^{-1})}$. From (5) and the assumption $\alpha \leq \beta$ it follows that $|A_j| \leq C$.

If $j = n$ it follows from (3) that $|A_n| \leq C \|\phi\|_*$ so we have in all cases

$$|A_j| \leq C \|\phi\|_*, \quad 0 \leq j \leq n. \quad (6)$$

We shall now put

$$B_j = \int_{2I} (\phi - \phi_{2I}) D^j K a D^{n-j} K b \, dx.$$

If $1 \leq j \leq n-1$ it follows from the lemma that $|D^j K a D^{n-j} K b| \leq C \alpha^{-j} \beta^{j-n} \leq C |I|^{-1}$. Hence

$$|B_j| \leq C |I|^{-1} \int_{2I} |\phi - \phi_{2I}| \, dx \leq C \|\phi\|_*.$$

From (3) it follows that

$$|B_0| + |B_n| \leq C \left(\int_{2I} |\phi - \phi_{2I}|^2 \, dx \right)^{1/2} (\|D^n K b\|_2 + \|D^n K a\|_2).$$

It is known that $(|I|^{-1} \int_I |\phi - \phi_I|^2)^{1/2} \leq C \|\phi\|_*$, see John and Nirenberg [4]. Since

$$\begin{aligned} \|D^n K a\|_2 + \|D^n K b\|_2 &\leq C (\|a\|_2 + \|b\|_2) \\ &\leq C (|I|^{-1/2} + |J|^{-1/2}) \leq C |I|^{-1/2} \end{aligned}$$

it follows that

$$|B_j| \leq C \|\Phi\|_*, \quad 0 \leq j \leq n.$$

Since $\int_{R^n} D^n (K a K b) \, dx = 0$ we have

$$\int_{R^n} \phi D^n (K a K b) \, dx = \sum_0^n \binom{n}{j} (A_j + B_j),$$

which yields (4).

To prove the theorem it is now sufficient to show that if $F = K a K b$ then there is a function $h \in H^1$ such that $\hat{h}(\xi) = |\xi|^n \hat{F}(\xi)$ and $\|h\| \leq C$.

Let f_i , $1 \leq i \leq n$, be defined by $\hat{f}_i(\xi) = \xi_i^n \hat{F}(\xi)$. From (4) follows $\|f_i\| \leq C$. If T_i denotes the operator defined by $\widehat{T_i(f)}(\xi) = \xi_i^n |\xi|^{-n} \hat{f}(\xi)$ it follows from theory of multipliers on H^1 (see Stein [5, Chapter VII]) that $\|T_i f\| \leq C \|f\|$. Hence $\|g\| \leq C$, where $g = \sum_1^n T_i f_i$. If T denotes the operator defined by $\widehat{Tf}(\xi) = |\xi|^{2n} (\sum_1^n \xi_i^{2n})^{-1} \hat{f}(\xi)$, then T is bounded on H^1 . Finally, observing that $\widehat{Tg}(\xi) = |\xi|^n \hat{F}(\xi)$ yields the theorem.

REFERENCES

1. R. Bagby, *Lebesgue spaces of parabolic potentials*, Illinois J. Math. **15** (1971), 610–635.
2. R. R. Coifman and G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. **83** (1977), 569–645.
3. C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), 137–193.
4. F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), 415–426.

5. E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Math. Series, No. 30, Princeton Univ. Press, Princeton, N. J., 1970.
6. R. S. Strichartz, *Multipliers on fractional Sobolev spaces*, J. Math. Mech. **16** (1967), 1031–1060.
7. ———, *A note on Sobolev algebras*, Proc. Amer. Math. Soc. **29** (1971), 205–207.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GÖTEBORG AND CHALMERS UNIVERSITY OF TECHNOLOGY, FACK, S-40220 GÖTEBORG, SWEDEN