ON THE NORM OF THE CANONICAL PROJECTION OF E^{***} ONTO $E^\perp$

JERRY JOHNSON AND JOHN WOLFE¹

ABSTRACT. For each number t with $1 < t \leq 2$, a renorming of c_0 is given for which $\|I - P\| = t$, where P is the canonical projection of c_0^{***} on c_0^* .

If E is a Banach space, then its dual E^* resides canonically in the third dual E^{***} . The mapping $P: E^{***} \rightarrow E^*$ that restricts a functional in E^{***} to E defines a projection of E^{***} onto E^* with norm one. Thus, if E is nonreflexive of course, $1 \leq \|I - P\| \leq 2$. It is not hard to prove that if $E = c_0$, then $\|I - P\| = 1$. The paper [1] is devoted to proving $\|I - P\| = 2$ for $E = l_1$. Furthermore, it seems that $\|I - P\|$ is either 1 or 2 in the (very few) examples in which the problem seems tractable. Fran Sullivan thus posed the question whether there are examples for which $1 < \|I - P\| < 2$. We answer this by constructing, for each number t with $1 < t \leq 2$, a renorming of c_0 so that $\|I - P\| = t$.

We proceed as follows:

Let $0 < r \leq 1$. For $x = (x_1, x_2, \dots) \in c_0$, define

(*) $\|x\| = \max(r^{-1}|x_1|, \sup_n |x_1 - x_n|)$. Then we have the following:

(1) $(1 + r)^{-1}\|x\|_\infty \leq \|x\| \leq \max(2, r^{-1})\|x\|_\infty$, where $\|x\|_\infty = \sup_n |x_n|$.

(2) By virtue of (1), the dual of c_0 is l_1 and for $\lambda = (\lambda_1, \lambda_2, \dots) \in l_1$ the dual norm is

$$\|\lambda\| = r \left| \sum_{n=1}^{\infty} \lambda_n \right| + \sum_{n=2}^{\infty} |\lambda_n|.$$

(3) The second dual norm on l_∞ is given by the same formula as (*).

(4) If P is the canonical projection of l_∞^* onto l_1 , then $\|I - P\| = 1 + r$.

PROOFS. (1) If $\|x\| \leq 1$, then $|x_1| \leq r$ and $|x_1 - x_n| \leq 1$. Hence, $|x_n| \leq 1 + r$ for $n \geq 1$. The other half is easier.

(2) Notice that for $x \in c_0$ we have

$$\begin{aligned} \sum_{n=1}^{\infty} \lambda_n x_n &= \left(r \sum_{n=1}^{\infty} \lambda_n \right) \left(\frac{1}{r} x_1 \right) + \sum_{n=2}^{\infty} \lambda_n (x_n - x_1) \\ &\leq \left(r \left| \sum_{n=1}^{\infty} \lambda_n \right| + \sum_{n=2}^{\infty} |\lambda_n| \right) \|x\|. \end{aligned}$$

Hence the dual of norm (*) on l_1 is no larger than the norm (2). To see that

Received by the editors April 30, 1978.

AMS (MOS) subject classifications (1970). Primary 46B10.

¹The second author was partially supported by NSF MCS 77-01690

we get equality, assume (without loss of generality) $\sum_{n=1}^\infty \lambda_n \geq 0$. Given n , let $x_j = 0$ for $j > n$; $x_1 = r$; $x_j = r + 1$ if $1 < j \leq n$ and $\lambda_j > 0$; and $x_j = r - 1$ if $1 < j \leq n$ and $\lambda_j < 0$. Then

$$\sum_{j=1}^\infty \lambda_j x_j = r \left| \sum_{j=1}^\infty \lambda_j \right| + \sum_{j=2}^\infty |\lambda_j|.$$

Hence the norm in (2) is the dual of norm (*).

(3) This can be obtained using the same formula,

$$\sum_{n=1}^\infty \lambda_n x_n = \left(r \sum_{n=1}^\infty \lambda_n \right) \left(\frac{1}{r} x_1 \right) + \sum_{n=2}^\infty \lambda_n (x_n - x_1),$$

as in (2). The details are omitted.

(4) We begin by defining $Q_n: l_\infty \rightarrow l_\infty$ by letting the first n coordinates of $Q_n x$ be zero and for $j > n$ let the j th coordinate be x_j . Now, $\|x\| \leq 1$ implies $|x_1| \leq r$ and $|x_j - x_1| \leq 1$, so $|x_j| \leq 1 + r$ for $j > 1$. Hence $\|Q_n x\|_\infty \leq 1 + r$. But if the first term of an element of l_∞ is zero, then its norm is the usual sup-norm. Thus $\|Q_n\| \leq 1 + r$.

Now we use the fact that l_∞^* can be considered as the space of Radon measures on the Stone-Čech compactification βN of the positive integers. (We use μ interchangeably as a measure and linear functional.) The canonical projection P of l_∞^* onto l_1 sends μ to the sequence $\{\mu\{n\}\}$ and so $(I - P)\mu$ is the restriction of μ to $\beta N \setminus N$. Let $\lambda = P\mu$ and $\tilde{\mu} = (I - P)\mu$. Since $\tilde{\mu}(N) = 0$, it follows that $\tilde{\mu}(Q_n x) = \tilde{\mu}(x)$ for each n and each $x \in l_\infty$. Thus

$$\begin{aligned} |\tilde{\mu}(x)| &= |\tilde{\mu}(Q_n x)| = |\mu(Q_n x) - \lambda(Q_n x)| \\ &\leq \|\mu\| \|Q_n x\| + \sum_{j=n}^\infty |\lambda_j x_j| \\ &\leq \|\mu\| (1 + r) \|x\| + \|x\|_\infty \sum_{j=n}^\infty |\lambda_j|. \end{aligned}$$

Taking the limit on n , we get $\|\tilde{\mu}\| \leq (1 + r) \|\mu\|$. Hence $\|I - P\| \leq 1 + r$.

Now, let $s \in \beta N \setminus N$. Let ϵ_s be the point mass at s and ϵ_1 be the point mass at $1 \in N$. Take a net $\{n_\gamma\}$ in N converging to s . If $\|x\| \leq 1$, then $|x_{n_\gamma}| \leq 1 + r$ for each $n_\gamma > 1$, so

$$|\epsilon_s(x)| = \lim_\gamma |x_{n_\gamma}| \leq 1 + r.$$

Also $|x_{n_\gamma} - x_1| \leq 1$ for every γ and therefore

$$|(\epsilon_s - \epsilon_1)(x)| = \lim_\gamma |x_{n_\gamma} - x_1| \leq 1.$$

Thus $\|\epsilon_s\| \leq 1 + r$ and $\|\epsilon_s - \epsilon_1\| \leq 1$. But if $y = (r, 1 + r, 1 + r, \dots)$, then $\|y\| = 1$, $\epsilon_s(y) = 1 + r$ and $(\epsilon_s - \epsilon_1)(y) = 1$. Hence $\|\epsilon_s\| = 1 + r$, $\|\epsilon_s - \epsilon_1\| = 1$, and $(I - P)(\epsilon_s - \epsilon_1) = \epsilon_s$ so $\|I - P\| = 1 + r$. This completes the proof.

A. L. Brown has very recently communicated to us another way of renorming c_0 to achieve the results given above as follows: Let T be an

isomorphism of c_0 onto the space of convergent sequences c . For each $t \in [0, 1]$, $\|x\|_t = t\|x\|_\infty + (1 - t)\|Tx\|_\infty$ defines an equivalent norm on c_0 , and as t ranges from 0 to 1 $\|I - P\|$ ranges from 2 to 1.

The referee has pointed out that all renormings of c_0 presented here have duals isometric to l_1 . Thus $\|I - P\|$ is not an isometric invariant of E^* . He also points out that examples isometric to these were introduced for another purpose in [2].

REFERENCES

1. A. L. Brown, *On the canonical projection of the third dual of a Banach space onto the first dual*, Bull. Austral. Math. Soc. **15** (1976), 351–354.
2. Y. Benyamini, *Near isometries in the class of L^1 -preduals*, Israel J. Math. **20** (1975), 275–281.

DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, STILLWATER, OKLAHOMA 74074