A THEOREM ON EXTENDING REPRESENTATIONS

RAYMOND C. FABEC¹

ABSTRACT. We obtain sufficient conditions for a unitary representation of a closed subgroup M of a separable locally compact group G to have a unique unitary extension to G . The conditions depend on the behavior of the representation on a closed normal subgroup N of G contained in M . We then discuss an application of the theorem in particle physics.

Let N be a closed normal subgroup of a separable locally compact group G , and let M be a closed subgroup of G containing N . Let $H = \{m \in M: gmg^{-1} \in M \text{ for } g \in G\}$. Assume L is a representation of M such that $L|_N$ is type I, and $L|_N = \int_{\hat{N}}^{\oplus} n(\alpha)\alpha \, d\mu(\alpha)$, μ a standard measure on $\hat{N}$; i.e. there is a μ conull set in $\hat{N}$ which is standard. By the von Neumann selection theorem, there is a cross-section c from $\hat{N}$ into the concrete irreducible representations of N such that c is Borel on a conull set in $\hat{N}$. Then

$$\int_{\hat{N}}^{\oplus} n(\alpha)\alpha \, d\mu(\alpha) = \int_{\hat{N}}^{\oplus} n(\alpha)c(\alpha) \, d\mu(\alpha),$$

and this gives the primary decomposition of $L|_N$. G acts on $\hat{N}$ by $\alpha \cdot g = \{R \cdot g: R \in \alpha\}$ where $(R \cdot g)(n) = R(gng^{-1})$. We do the transitive case first.

THEOREM 1. Assume μ is supported on $\alpha_0 \cdot G$. Assume the stabilizer K of α_0 is contained in H and $L|_H \cdot g$ is unitarily equivalent to $L|_H$ for each $g \in G$. Then L has a unique unitary extension to G . In fact, there is a representation W of K such that $\text{ind}_K^G W$ is the extension. This extension is irreducible precisely when W is irreducible.

REMARK. $L = \int_{G/M}^{\oplus} \text{ind}_{x^{-1}Kx}^M W \cdot x \, d\bar{\mu}(x)$ where $\bar{\mu}$ is a quasi-invariant measure on G/M .

PROOF. Let P be the projection valued measure based on the Borel subsets of $\alpha_0 \cdot G \simeq K \setminus G$ giving the direct integral decomposition $L|_N = \int^{\oplus} n(\alpha)c(\alpha) \, d\mu(\alpha)$. Since $L|_H \cdot g \simeq L$ for $g \in G$, $L|_N \cdot g \simeq L|_N$ for $g \in G$; and the argument given by Mackey [4, p. 296], shows μ is quasi-invariant and $n(\alpha \cdot g^{-1}) = n(\alpha)$ for μ -a.e. α for each g . Hence, since G acts transitively, n is essentially constant. Therefore

$$L|_N \simeq n \int^{\oplus} c(\alpha) \, d\mu(\alpha) = n \int^{\oplus} c(\alpha) \, d\mu(\alpha) = n \int^{\oplus} \omega \cdot \gamma(Kx) \, d\mu(Kx)$$

Received by the editors February 27, 1978 and, in revised form, July 6, 1978.

AMS (MOS) subject classifications (1970). Primary 22D10, 22D30; Secondary 22E70.

¹Research supported in part by a Louisiana State Summer Faculty Research Award.

© 1979 American Mathematical Society
0002-9939/79/0000-0283/\$02.50

where $\omega \in \alpha_0$ and $\gamma: K \setminus G \rightarrow G$ is a Borel cross-section.

Since $L_m L|_N L_m^{-1} = L|_N \cdot m \simeq L|_N$ and $n \int^\oplus \omega \cdot \gamma(Kx) d\mu(Kx)$ is the central decomposition of $L|_N$, $L_m P(E) L_m^{-1} = P(E \cdot m^{-1})$ for Borel subsets E of $K \setminus G$ and $m \in M$. Furthermore, the orbit space $(K \setminus G)^M$ of $K \setminus G$ under M is isomorphic to G/M . In fact $KxM = x x^{-1} KxM = xM$, for $x^{-1} Kx \subset x^{-1} Hx = H$. Hence the orbit space is standard; and if $\bar{\mu}$ is the image of μ under the map $Kx \mapsto xM$, μ decomposes uniquely $\bar{\mu}$ -a.e. into an "integral of measures", $\mu = \int_{G/M} \mu_{xM} d\bar{\mu}(xM)$, where each μ_{xM} is a Borel measure on $K \setminus G$ with $\mu_{xM}(K \setminus G - KxM) = 0$ (see [2, Theorem 11]). Since μ is quasi-invariant under M , μ_{xM} is quasi-invariant under M for $\bar{\mu}$ -a.e. xM .

The decomposition $\mu = \int_{G/M} \mu_{xM} d\bar{\mu}(xM)$ yields a decomposition of P into $\int_{G/M}^\oplus P(xM) d\bar{\mu}(xM)$ where each $P(xM)$ is a projection valued measure defined on the Borel subsets of xM satisfying $P(xM)(E) = 0$ iff $\mu_{xM}(E) = 0$. Define a projection valued measure Q on the Borel subsets of G/M by $Q(F) = P(\cup F)$. Then

$$L_m^{-1} Q(F) L_m = P\left(\bigcup F \cdot m\right) = P\left(\bigcup F\right) = Q(F).$$

Therefore L decomposes into a direct integral over G/M ; namely, $L = \int^\oplus L(xM) d\bar{\mu}(xM)$. Furthermore,

$$L(xM)|_N = n \int_{KxM}^\oplus \omega \cdot \gamma(Kz) d\mu_{xM}(Kz)$$

and

$$L(xM)(m^{-1})P(xM)(E)L(xM)(m) = P(xM)(E \cdot m),$$

E any Borel subset of KxM . Hence the pair $(L(xM), P(xM))$ forms a transitive system of imprimitivity based on the transitive M space KxM . Let $\sigma: G/M \rightarrow G$ be a Borel cross-section. By Mackey's imprimitivity theorem [3, Theorem 6.6], $L(xM) = \text{ind}_{\sigma(xM)^{-1}K\sigma(xM)}^M W(xM)$, where $W(xM)$ is a unitary representation of $\sigma(xM)^{-1}K\sigma(xM)$. One can show $W(xM)|_N \simeq n\omega \cdot \sigma(xM)$.

We consider the representation $L|_H$. The orbit space $(K \setminus G)^H$ of $K \setminus G$ under H is isomorphic to $H \setminus G$ since $KxH = KxHx^{-1}x = Hx$. Hence $\mu = \int \nu_{Hx} d\bar{\nu}(Hx)$ where $\bar{\nu}$ is the image of μ under the map $Kx \rightarrow Hx$ and ν_{Hx} is an H quasi-invariant measure on Hx . This decomposition, as before, gives a decomposition of P into $\int^\oplus P(Hx) d\bar{\nu}(Hx)$.

$\bar{\mu}$ is the image of $\bar{\nu}$ under the map $Hx \rightarrow xM$. Therefore $\bar{\nu}$ decomposes relative to $\bar{\mu}$: $\bar{\nu} = \int_{G/M} \bar{\nu}_{xM} d\bar{\mu}(xM)$, $\bar{\nu}_{xM}$ a measure on $H \setminus G$ with

$$\bar{\nu}_{xM}(H \setminus G - xM) = 0.$$

Hence

$$\begin{aligned} \mu &= \int_{H \setminus G} \nu_{Hx} d\bar{\nu}(Hx) = \int_{G/M} \int_{H \setminus G} \nu_{Hy} d\bar{\nu}_{xM}(Hy) d\bar{\mu}(xM) \\ &= \int_{G/M} \mu_{xM} d\bar{\mu}(xM). \end{aligned}$$

By uniqueness of decompositions, $\mu_{xM} = \int_{H \setminus G} \nu_{Hy} d\bar{\nu}_{xM}(Hy)$ for $\bar{\mu}$ -a.e. xM . Furthermore, one has $P(xM) = \int_{H \setminus G} P(Hy) d\bar{\nu}_{xM}(Hy)$.

Define a projection valued measure Q^* on the Borel subsets of $H \setminus G$ by $Q^*(F) = P(\cup F)$. Then $L_h Q^*(F) L_h^{-1} = Q^*(F)$ for $h \in H$. Therefore $L|_H$ decomposes into a direct integral:

$$L|_H = \int_{H \setminus G}^{\oplus} L(Hy) d\bar{\nu}(Hy),$$

where $L(Hy)|_N = n \int_{Hy} \omega \cdot \gamma(Kz) d\nu_{Hy}(Kz)$. Furthermore, since $P(xM) = \int_{xM} P(Hy) d\bar{\nu}_{xM}(Hy)$, one has

$$L(xM)|_H = \bigind_{\sigma(xM)^{-1}K\sigma(xM)}^M W(xM)|_H = \int^{\oplus} L(Hy) d\bar{\nu}_{xM}(Hy).$$

This decomposition of $(L(xM), P(xM))$ is precisely the one given by Mackey's subgroup theorem. Note KxM as an M space is isomorphic to $\sigma(xM)^{-1}K\sigma(xM) \setminus M$ and the orbit space of KxM under H is isomorphic to the double coset space of $\sigma(xM)^{-1}K\sigma(xM) - H$ double cosets in M . The correspondence with KxM carries the double coset $\sigma(xM)^{-1}K\sigma(xM)mH$ to $K\sigma(xM)mH = H\sigma(xM)m$. Hence the double cosets of KxM are the H cosets in KxM . Now let $\rho: H \setminus G \rightarrow G$ be a Borel cross-section. Then $\rho|_{KxM} \rightarrow G$ is a cross-section of the $K - H$ double cosets in KxM and $Hy = K\sigma(xM)\sigma(xM)^{-1}\rho(Hy)H$. Therefore

$$L(Hy) \simeq \bigind_{\rho(Hy)^{-1}K\rho(Hy)}^H W(xM) \cdot \sigma(xM)^{-1}\rho(Hy)$$

$\bar{\nu}_{xM}$ -a.e. Hy $\bar{\mu}$ -a.e. xM . For more details, see the proof of the subgroup theorem [4, p. 227].

We now have

$$L|_H \simeq \int_{H \setminus G}^{\oplus} \bigind_{\rho(Hy)^{-1}K\rho(Hy)}^H W(yM) \cdot \sigma(yM)^{-1}\rho(Hy) d\bar{\nu}(Hy)$$

and $P = \int^{\oplus} P(Hy) d\bar{\nu}(Hy)$ where $P(Hy)$ is the canonical projection valued measure associated with the induced representation $L(Hy)$. Therefore

$$L|_H \simeq L|_H \cdot g \simeq \int^{\oplus} \bigind_{g^{-1}\rho(Hy)^{-1}K\rho(Hy)g}^H W(yM) \cdot \sigma(yM)^{-1}\rho(Hy) g d\bar{\nu}(Hy).$$

Since $\rho(Hy)g = \rho(Hyg)h$ for some h ,

$$L|_H \simeq \int^{\oplus} \bigind_{\rho(Hyg)^{-1}K\rho(Hyg)}^H W(yM) \cdot \sigma(yM)^{-1}\rho(Hyg) d\bar{\nu}(Hy).$$

In this last direct integral decomposition, P decomposes into $P = \int^{\oplus} P(Hy) \cdot g d\bar{\nu}(Hy)$ where $P(Hy) \cdot g(E) = P(Hy)(E \cdot g^{-1})$. Changing $y \rightarrow yg^{-1}$ in the last decomposition yields

$$L|_H \simeq \int^{\oplus} \bigind_{\rho(Hy)^{-1}K\rho(Hy)}^H W(yg^{-1}M) \cdot \sigma(yg^{-1}M)^{-1}\rho(Hy) d\bar{\nu}(Hy)$$

and

$$P = \int^{\oplus} P(Hyg^{-1}) \cdot g \, d\bar{\nu}(Hy).$$

But

$$P(Hyg^{-1}) \cdot g = P(Hy).$$

Hence we have two decompositions of $(L|_H, P)$ relative to Q^* . Therefore

$$\begin{aligned} & \left(\text{ind } W(yg^{-1}M) \cdot \sigma(yg^{-1}M)^{-1} \rho(Hy), P(Hy) \right) \\ & \simeq \left(\text{ind } W(yM) \cdot \sigma(yM)^{-1} \rho(Hy), P(Hy) \right) \end{aligned}$$

for $\bar{\nu}$ -a.e. Hy . It follows that

$$W(yg^{-1}M) \cdot \sigma(yg^{-1}M)^{-1} \rho(Hy) \simeq W(yM) \cdot \sigma(yM)^{-1} \rho(Hy)$$

for $\bar{\nu}$ -a.e. Hy for each $g \in G$. Hence there is a representation W of K such that $W(xM)$ is unitarily equivalent to the representation $\sigma(xM)^{-1} k \sigma(xM) \rightarrow W(k)$ for $\bar{\mu}$ -a.e. xM . We then have

$$L \simeq \int_{G/M}^{\oplus} \text{ind}_{\sigma(xM)^{-1} k \sigma(xM)}^M W \cdot \sigma(xM) \, d\bar{\mu}(xM).$$

Let $U = \text{ind}_K^G W$. By Mackey's subgroup theorem, U extends L . U is unique for if U is an extension of L , (U, P) is a transitive system of imprimitivity based on $K \setminus G$. Hence $U = \text{ind}_K^G W'$. But then

$$\begin{aligned} & \left(\text{ind}_{\sigma(xM)^{-1} k \sigma(xM)}^M W \cdot \sigma(xM), P(xM) \right) \\ & \simeq \left(\text{ind}_{\sigma(xM)^{-1} k \sigma(xM)}^M W' \cdot \sigma(xM), P(xM) \right) \end{aligned}$$

for $\bar{\mu}$ -a.e. xM . Hence $W \simeq W'$.

The commuting rings of U and W are isomorphic. Hence U is irreducible when W is. Q.E.D.

REMARK. The stabilizer K of α_0 is contained in H if the stabilizer of μ -a.e. α is contained in M .

We next assume μ is based on a standard G -invariant subset S of $\hat{N}$.

THEOREM 2. Assume μ is based on S and G acts regularly on S and $L|_H \cdot g \simeq L|_H$ for $g \in G$. Then L has a unique extension from M to G provided that the stabilizer of α is contained in M for μ -a.a. α .

PROOF. Let S^G be the orbit space for S under G . Let $\bar{\mu}$ be the quotient measure. The action is regular if there exists a $\bar{\mu}$ conull set $\bar{S}_0$ in S^G such that $\bar{S}_0$ is standard. Replacing S by the inverse image S_0 of $\bar{S}_0$ in S , we may

assume S^G is standard. Let P be the projection valued measure defined in the proof of Theorem 1. The measure μ decomposes relative to $\bar{\mu}$; $\mu = \int \mu_{\bar{s}} d\bar{\mu}(\bar{s})$ where each $\mu_{\bar{s}}$ is a G quasi-invariant measure based on $\bar{s}$. This leads to a decomposition of both P and L : $P = \int^{\oplus} P^{\bar{s}} d\bar{\mu}(\bar{s})$ and $L = \int^{\oplus} L^{\bar{s}} d\bar{\mu}(\bar{s})$. Since the stabilizer of α is contained in M for μ -a.a. $\alpha \in S$, the stabilizer of $\alpha \in \bar{s}$ is contained in M for $\mu_{\bar{s}}$ -a.a. α for $\bar{\mu}$ -a.a. $\bar{s}$. Since $L|_H \cdot g \simeq L|_H$, $L^{\bar{s}}|_H \cdot g \simeq L^{\bar{s}}|_H$ for $\bar{\mu}$ -a.e. $\bar{s}$. By Fubini's theorem, for $\bar{\mu}$ -a.e. $\bar{s}$, $L^{\bar{s}}|_H \cdot g \simeq L^{\bar{s}}|_H$ a.e. g . But $\{g: L^{\bar{s}}|_H \cdot g \simeq L^{\bar{s}}|_H\}$ is a subgroup of G . Hence for $\bar{\mu}$ -a.e. $\bar{s}$, $L^{\bar{s}}|_H$ is invariant under G . Hence we may apply Theorem 1 and obtain an extension $R^{\bar{s}}$ of $L^{\bar{s}}$ to G for $\bar{\mu}$ -a.a. $\bar{s}$. The extension is a Borel function of $\bar{s}$ and hence $\int^{\oplus} R^{\bar{s}} d\bar{\mu}(\bar{s})$ is an extension of L . Since $R^{\bar{s}}$ is unique a.e. $\bar{s}$, this extension is unique. Q.E.D.

REMARK. These theorems remain valid for σ -representations. One can either work in the group extension or modify the above proofs.

As an application of the theorem we show how it is possible to determine the dynamics of a free particle system having time translations which do not commute with spatial translations. For instance the spatial transformations of $\mathbf{R}^3$ may be the Euclidean group $E_3 = \{(x, R): x \in \mathbf{R}^3 \text{ and } R \in \text{SO}(3)\}$ with multiplication defined by $(x, R)(y, S) = (x + Ry, RS)$. Time translation by t , however, will have the property $t(x, R) = (e^t x, R)t$. This may occur, for instance, if physical space expands exponentially with time. In any case, for this group a free particle of spin j , j a half integer, is a projective representation U such that $U_{(x,R)} f(y) = e^{2\pi i x \cdot y} D^j(R) f(R^{-1}y)$ for $f \in L^2(\mathbf{R}^3, C^{2j+1})$ where D^j is the projective irreducible representation of $\text{SO}(3)$ with dimension $2j + 1$. We shall, for convenience, assume j is an integer, for in this case D^j is an ordinary representation. Next note one has $M = E_3$ and $N = \mathbf{R}^3$. Then

$$U|_{\mathbf{R}^3} = (2j + 1) \int_{\mathbf{R}^3}^{\oplus} e^{2\pi i x \cdot ()} dx$$

and since

$$\exp(2\pi i x \cdot ()) \cdot t(0, R) = \exp(2\pi i e^{-t} R^{-1} x \cdot ()),$$

the orbit of $e_1 = (1, 0, 0)$ is almost all of $\mathbf{R}^3$. The stabilizer K of e_1 is $\{(y, R): Re_1 = e_1, y \in \mathbf{R}^3\}$ while $H = M = E_3$. Hence Theorem 1 applies and we need only find W . Note $U|_{E_3} = \int_{\mathbf{R}}^{\oplus} W(r) dr$ where $W(r)$ is the representation of E_3 on $L^2(S_r, m_r, C^{2j+1})$, m_r the Lebesgue measure on the sphere of radius e^r , defined by

$$W(r)(t, R) f(s) = e^{2\pi i x \cdot s} D^j(R) f(R^{-1}y).$$

One can show $W(r) = \text{ind}_K^M (W \cdot t) = \text{ind}_{i_K t^{-1}}^M W \cdot t$ where $W(x, R) = e^{2\pi i x_1} D^j(R)$ for $(x_1, x_2, x_3, R) \in K$. Hence $U = \text{ind}_K^G W$.

For more information on particle systems having these symmetries, see [1].

REFERENCES

1. R. Fabec, *Free particle systems and their dynamics in de Sitter space*, J. Mathematical Phys. **18** (1977), 2087–2091.
2. G. W. Mackey, *Induced representations of a locally compact group*. I, Ann. of Math. **2** (1952), 101–139.
3. ———, *The theory of unitary group representations*, 1st ed., Univ. of Chicago Press, Chicago, Ill., 1976, 372 pp.
4. ———, *Unitary representations of group extensions*. I, Acta Math. **99** (1958), 265–311.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA 70803