SOME INEQUALITIES OF ALGEBRAIC POLYNOMIALS HAVING REAL ZEROS

A. K. VARMA

Dedicated to Professor A. Zygmund

ABSTRACT. Let $P_n(x)$ be an algebraic polynomial of degree n having all real zeros. We set

$$I_n = \frac{\|P'_n(x)\omega(x)\|_{L_2[a,b]}}{\|P_n(x)\omega(x)\|_{L_2[a,b]}}.$$

In this work the lower and upper bounds of I_n are investigated under the assumptions that all the zeros of $P_n(x)$ are inside $[a, b]$ and outside $[a, b]$, respectively. We restrict ourselves here with two cases, (1) $\omega(x) = (1 - x^2)^{1/2}$, $[a, b] = [-1, 1]$; (2) $\omega(x) = e^{-x/2}$, $[a, b] = [0, \infty)$. Results are shown to be best possible.

In this work we are concerned with the following theorems of P. Turán [3], P. Erdős [1], and A. K. Varma [5].

THEOREM A. Let $P_n(x)$ be an algebraic polynomial of degree n having all its zeros inside $[-1, +1]$; then we have

$$\max_{-1 < x < +1} |P'_n(x)| > \frac{n^{1/2}}{6} \max_{-1 < x < +1} |P_n(x)|. \quad (1.1)$$

($P'_n(x)$ stands for the derivative of $P_n(x)$.)

THEOREM B. Let $P_n(x)$ be a polynomial of degree n satisfying the inequality $|P_n(x)| < 1$ for $-1 < x < +1$. Suppose $P_n(x)$ has only real roots and no root inside $[-1, +1]$; then for $-1 < x < +1$, $|P'_n(x)| < \frac{1}{2}en$. This is the best possible result.

THEOREM C. Let $P_n(x)$ be an algebraic polynomial of degree n having all zeros real and inside $[-1, +1]$; then we have

$$\|P'_n\|_{L_2[-1, +1]}^2 > \left(\frac{n}{2} + \frac{3}{4} + \frac{3}{4n} \right) \|P_n\|_{L_2[-1, +1]}^2, \quad n > 13, \quad (1.2)$$

where

$$\|P_n\|_{L_2[-1, +1]}^2 = \int_{-1}^1 P_n^2(x) dx.$$

Further, if $P_n(+1) = P_n(-1) = 0$ and $n > 2$ then under the above conditions

$$\|P'_n\|_{L_2[-1, +1]}^2 > \left(\frac{n}{2} + \frac{3}{4} + \frac{3}{4(n-1)} \right) \|P_n\|_{L_2[-1, +1]}^2$$

Received by the editors July 26, 1978.

AMS (MOS) subject classifications (1970). Primary 26A60, 30A40.

© 1979 American Mathematical Society
0002-9939/79/0000-0311/\$03.00

with equality holding for $P_n(x) = (1 - x^2)^m$, $n = 2m$.

The above Theorem C is analogous to Theorem A in the L_∞ norm. In view of the above theorems it is natural to ask about lower bound (upper bound) of the expression

$$\|\omega(x)P'_n(x)\|_{L_2[a,b]}/\|\omega(x)P_n(x)\|_{L_2[a,b]}$$

under the assumption that all the zeros of $P_n(x)$ are all inside $[a, b]$ (no roots inside $[a, b]$, respectively). We limit ourselves here in this work to the above question when $[a, b] = [0, \infty)$, $\omega(x) = e^{-x/2}$ and $[a, b] = [-1, +1]$, $\omega(x) = (1 - x^2)^{1/2}$.

Throughout x_k 's denote the zeros of $P_n(x)$. Here we prove the following theorems.

THEOREM 1. *Let $P_n(x)$ be an algebraic polynomial of degree n having all zeros inside $[0, \infty)$. Let $P_n(0) = 0$ or*

$$\sum_{k=1}^n \frac{1}{x_k} > \frac{1}{2};$$

then

$$\|e^{-x/2}P'_n\|_{L_2[0,\infty)}^2 > \frac{n}{2(2n-1)} \|e^{-x/2}P_n\|_{L_2[0,\infty)}^2 \quad (1.3)$$

with equality for $P_n(x) = x^n$.

The following remark is needed concerning the above theorem. Let us choose $P_n(x) = (x - x_1)^n$, $x_1 > 0$. It can be shown easily that the ratio of

$$\int_0^\infty (P'_n(x))^2 e^{-x} dx / \int_0^\infty (P_n(x))^2 e^{-x} dx$$

can be made arbitrary small for such polynomials by choosing x_1 arbitrary large. Thus some condition on the roots such as $\sum_{k=1}^n (x_k)^{-1} > \frac{1}{2}$ is natural.

Next, we turn to the case when $\omega(x) = (1 - x^2)^{1/2}$, $[a, b] = [-1, +1]$. Here we prove

THEOREM 2. *Let $P_n(x)$ be an algebraic polynomial of degree n having all zeros real and inside $[-1, +1]$. Then for $n \geq 2$ we have*

$$\|(1 - x^2)^{1/2}P'_n\|_{L_2[-1,+1]}^2 > \left(\frac{n}{2} + \frac{1}{4} - \frac{1}{4(n+1)}\right) \|(1 - x^2)^{1/2}P_n\|_{L_2[-1,+1]}^2 \quad (1.4)$$

with equality for $P_n(x) = (1 - x^2)^m$, $n = 2m$.

The next two theorems are analogous to Theorem B of P. Erdős [1].

THEOREM 3. Let $P_n(x)$ be an algebraic polynomial of degree $\leq n$ having all real roots and no root inside the interval $[-1, +1]$; then we have

$$\begin{aligned} \|(1-x^2)^{1/2}P'_n(x)\|_{L_2[-1,+1]}^2 \\ \leq \frac{n(n+1)(2n+3)}{4(2n+1)} \|(1-x^2)^{1/2}P_n(x)\|_{L_2[-1,+1]}^2 \end{aligned} \quad (1.5)$$

with equality for $P_n(x) = (1+x)^n$ or $P_n(x) = (1-x)^n$.

THEOREM 4. Let $x_1, x_2, \dots, x_n$ be real zeros of an algebraic polynomial $P_n(x)$ of degree $\leq n$. Further, $-\infty < x_i \leq 0, i = 1, 2, \dots, n$. Then

$$\int_0^\infty x e^{-x} (P'_n(x))^2 dx \leq \frac{n}{2(2n+1)} \int_0^\infty x e^{-x} (P_n(x))^2 dx$$

with equality for $P_n(x) = x^n$.

2. Proof of Theorem 1. For the proof of this theorem we will need the following facts:

$$P'_n(x) = P_n(x) \sum_{k=1}^n \frac{1}{x - x_k}, \quad (2.1)$$

$$P_n'^2(x) - P_n(x)P_n''(x) = P_n^2(x) \sum_{k=1}^n \frac{1}{(x - x_k)^2}, \quad (2.2)$$

$$\frac{(x - x_1)^2}{(x - x_k)^2} = -1 + \frac{2(x - x_1)}{x - x_k} + \frac{(x_1 - x_k)^2}{(x - x_k)^2}, \quad (2.3)$$

$$P'_n(x) - n \frac{P_n(x)}{x - x_1} = \frac{P_n(x)}{x - x_1} \sum_{k=1}^n \frac{x_k - x_1}{x - x_k}. \quad (2.4)$$

From (2.4) and the Cauchy-Schwarz inequality we have

$$\left(P'_n(x) - n \frac{P_n(x)}{x - x_1} \right)^2 \leq n \frac{P_n^2(x)}{(x - x_1)^2} \sum_{k=1}^n \frac{(x_k - x_1)^2}{(x - x_k)^2}. \quad (2.5)$$

From (2.5) we immediately have

$$\begin{aligned} \int_0^\infty \left(P'_n(x) - n \frac{P_n(x)}{x - x_1} \right)^2 e^{-x} dx \\ \leq n \int_0^\infty \frac{P_n^2(x)}{(x - x_1)^2} \sum_{k=1}^n \frac{(x_k - x_1)^2}{(x - x_k)^2} e^{-x} dx. \end{aligned} \quad (2.6)$$

Also, integrating by parts, we have

$$\int_0^\infty e^{-x} P_n(x) P'_n(x) dx = -\frac{1}{2} P_n^2(0) + \frac{1}{2} \int_0^\infty e^{-x} P_n^2(x) dx. \quad (2.7)$$

Now, we turn to the proof of Theorem 1. On integrating by parts, we have

$$\int_0^\infty e^{-x} P_n'^2(x) dx = -P_n(0)P_n'(0) - \int_0^\infty P_n(x)(P_n''(x) - P_n'(x))e^{-x} dx, \quad (2.8)$$

and an equivalent form

$$2 \int_0^\infty e^{-x} P_n'^2(x) dx = P_n^2(0) \sum_{k=1}^n \frac{1}{x_k} + \int_0^\infty e^{-x} (P_n'^2(x) - P_n(x)P_n''(x)) dx \\ + \int_0^\infty e^{-x} P_n(x)P_n'(x) dx. \quad (2.9)$$

On using (2.2) and (2.7) we obtain

$$2 \int_0^\infty e^{-x} P_n'^2(x) dx = P_n^2(0) \left(-\frac{1}{2} + \sum_{k=1}^n \frac{1}{x_k} \right) + \frac{1}{2} \int_0^\infty e^{-x} P_n^2(x) dx \\ + \int_0^\infty e^{-x} P_n^2(x) \sum_{k=1}^n \frac{1}{(x - x_k)^2} dx. \quad (2.10)$$

Now, let us denote by x_1 the first nonnegative zero of $P_n(x)$. Then we can write

$$P_n(x) = (x - x_1)q_{n-1}(x), \quad x_1 \geq 0, \quad (2.11)$$

where $q_{n-1}(x)$ is a polynomial of degree $\leq n - 1$ having real zeros lying inside $[0, \infty)$.

We set

$$I_1 = \int_0^\infty e^{-x} P_n^2(x) \sum_{k=1}^n \frac{1}{(x - x_k)^2} dx = \int_0^\infty e^{-x} q_{n-1}^2(x) \sum_{k=1}^n \frac{(x - x_1)^2}{(x - x_k)^2} dx, \quad (2.12)$$

and on using (2.3) we have

$$I_1 = -n \int_0^\infty e^{-x} q_{n-1}^2(x) dx + 2 \int_0^\infty (x - x_1) q_{n-1}^2(x) \sum_{k=1}^n \frac{1}{(x - x_k)} e^{-x} dx \\ + \int_0^\infty e^{-x} q_{n-1}^2(x) \sum_{k=1}^n \frac{(x_1 - x_k)^2}{(x - x_k)^2} dx \\ = -n \int_0^\infty e^{-x} q_{n-1}^2(x) dx + 2 \int_0^\infty P_n'(x) q_{n-1}(x) e^{-x} dx \\ + \int_0^\infty e^{-x} q_{n-1}^2(x) \sum_{k=1}^n \frac{(x_1 - x_k)^2}{(x - x_k)^2} dx, \\ I_1 = \frac{1}{n} \int_0^\infty P_n^2(x) e^{-x} dx - \frac{1}{n} \int_0^\infty (P_n'(x) - n q_{n-1}(x))^2 e^{-x} dx \\ + \int_0^\infty e^{-x} q_{n-1}^2(x) \sum_{k=1}^n \frac{(x_1 - x_k)^2}{(x - x_k)^2} dx; \quad (2.13)$$

on putting the values of I_1 from (2.13) into (2.10) we obtain

$$\begin{aligned} \left(2 - \frac{1}{n}\right) \int_0^\infty e^{-x} P_n'^2(x) dx &= P_n^2(0) \left(-\frac{1}{2} + \sum_{k=1}^n \frac{1}{x_k}\right) + \frac{1}{2} \int_0^\infty e^{-x} P_n^2(x) dx \\ &\quad + \int_0^\infty e^{-x} q_{n-1}^2(x) \sum_{k=1}^n \frac{(x_1 - x_k)^2}{(x - x_k)^2} dx \\ &\quad - \frac{1}{n} \int_0^\infty (P_n'(x) - n q_{n-1}(x))^2 e^{-x} dx. \end{aligned} \quad (2.14)$$

Now if $P_n(0) = 0$ or $\sum_{k=1}^n (x_k)^{-1} > \frac{1}{2}$ then on using (2.6) we immediately obtain

$$\left(2 - \frac{1}{n}\right) \int_0^\infty e^{-x} P_n'^2(x) dx > \frac{1}{2} \int_0^\infty e^{-x} P_n^2(x) dx,$$

from which the proof of Theorem 1 is complete. We would like to remark that the ratio of

$$\int_0^\infty e^{-x} P_n'^2(x) dx / \int_0^\infty e^{-x} P_n^2(x) dx$$

does indeed get smaller than $n/2(2n-1)$ for polynomials $P_0(x) = (x - x_1)^n$ for $x_1 > 2n$. For such polynomials the last two terms in (2.14) are zero and $-\frac{1}{2} + \sum_{k=1}^n (x_k)^{-1} < 0$.

3. Proof of Theorem 2. In order to prove Theorem 2, we need some results proved in [5]. Let $P_n(x)$ be any polynomial of degree n ; then

$$2 \int_{-1}^1 (1 - x^2) P_n'^2(x) dx = n \int_{-1}^1 P_n^2(x) dx + S_0, \quad (3.1)$$

$$\begin{aligned} (n+1)(2n+3) \int_{-1}^1 x^2 P_n^2(x) dx &= S_2 - S_0 + (n+1) \int_{-1}^1 P_n^2(x) dx \\ &\quad + 2 \int_{-1}^1 ((1 - x^2) P_n'(x) + n x P_n(x))^2 dx \end{aligned} \quad (3.2)$$

where

$$S_{2i} = \int_{-1}^1 x^{2i} P_n^2(x) \sum_{k=1}^n \frac{(1 - x_k^2)}{(x - x_k)^2} dx, \quad i = 0, 1, \dots \quad (3.3)$$

For the proof of (3.1), (3.2) see (2.1) and (2.10) (with $r = 1$) in [5]. Equation (3.2) can be rewritten in the form

$$\begin{aligned} (n+1)(2n+3) \int_{-1}^1 (1 - x^2) P_n^2(x) dx \\ = S_0 - S_2 + 2(n+1)^2 \int_{-1}^1 P_n^2(x) dx \\ - 2 \int_{-1}^1 ((1 - x^2) P_n'(x) + n x P_n(x))^2 dx. \end{aligned}$$

Therefore

$$(n+1)(2n+3) \int_{-1}^1 (1-x^2) P_n^2(x) dx < S_0 - S_2 + 2(n+1)^2 \int_{-1}^1 P_n^2(x) dx. \quad (3.4)$$

From (3.1) and (3.4) we obtain

$$\frac{2 \int_{-1}^1 (1-x^2) P_n^2(x) dx}{(n+1)(2n+3) \int_{-1}^1 (1-x^2) P_n^2(x) dx} > \frac{n \int_{-1}^1 P_n^2(x) dx + S_0}{S_0 - S_2 + 2(n+1)^2 \int_{-1}^1 P_n^2(x) dx}. \quad (3.5)$$

It is given that $|x_k| < 1$, $S_0 > 0$, $S_2 > 0$ and $S_0 - S_2 > 0$ (here we use (3.3)). Hence it can be verified immediately that

$$\begin{aligned} & \left(S_0 + n \int_{-1}^1 P_n^2(x) dx \right) / \left(S_0 - S_2 + 2(n+1)^2 \int_{-1}^1 P_n^2(x) dx \right) \\ & > n / 2(n+1)^2. \end{aligned} \quad (3.6)$$

From (3.5) and (3.6) we obtain

$$\int_{-1}^1 (1-x^2) P_n^2(x) dx > \frac{n(2n+3)}{4(n+1)} \int_{-1}^1 (1-x^2) P_n^2(x) dx.$$

This proves Theorem 2 as well.

4. Proof of Theorem 3. For the proof of Theorem 3 we need the following lemma.

LEMMA 4.1. *Let $P_n(x)$ be an algebraic polynomial of degree $< n$ having real roots and no root in the interval $[-1, +1]$. Then we have*

$$\int_{-1}^1 (1-x^2) P_n^2(x) dx > \frac{2(2n+1)}{(n+1)(2n+3)} \int_{-1}^1 P_n^2(x) dx. \quad (4.1)$$

PROOF It follows from the well-known theorem [2] that for such $P_n(x)$ we have

$$P_n^2(x) = \sum_{p+q=2n} a_{pq} (1+x)^p (1-x)^q, \quad a_{pq} > 0. \quad (4.2)$$

Hence we have

$$\int_{-1}^1 (1-x^2) P_n^2(x) dx = \sum_{p+q=2n} a_{pq} \int_{-1}^1 (1+x)^{p+1} (1-x)^{q+1} dx.$$

But

$$\begin{aligned} \int_{-1}^1 (1+x)^{p+1}(1-x)^{q+1} dx \\ = \frac{4(p+1)(q+1)}{(p+q+3)(p+q+2)} \int_{-1}^1 (1+x)^p(1-x)^q dx. \end{aligned}$$

Hence

$$\begin{aligned} \int_{-1}^1 (1-x^2)P_n^2(x) dx \\ = 4 \sum_{p+q=2n} \frac{q_{pq}(p+1)(q+1)}{(p+q+3)(p+q+2)} \int_{-1}^1 (1+x)^p(1-x)^q dx \\ > \frac{4(2n+1)}{(2n+3)(2n+2)} \sum_{p+q=2n} q_{pq} \int_{-1}^1 (1+x)^p(1-x)^q dx \\ = \frac{2(2n+1)}{(n+1)(2n+3)} \int_{-1}^1 P_n^2(x) dx. \end{aligned}$$

This proves the lemma. Now we turn to the proof of the theorem. Since $|x_k| > 1$, $k = 1, 2, \dots, n$, it follows from (3.1) and (3.3) that

$$2 \int_{-1}^1 (1-x^2)P_n'^2(x) dx < n \int_{-1}^1 P_n^2(x) dx, \quad (4.3)$$

with equality holding for $P_n(x) = (1+x)^p(1-x)^q$, $p+q=n$. Now, on using (4.1) and (4.3) we obtain

$$\begin{aligned} \frac{\int_{-1}^1 (1-x^2)P_n'^2(x) dx}{\int_{-1}^1 (1-x^2)P_n^2(x) dx} &= \frac{\int_{-1}^1 (1-x^2)P_n'^2(x) dx}{\int_{-1}^1 P_n^2(x) dx} \cdot \frac{\int_{-1}^1 P_n^2(x) dx}{\int_{-1}^1 (1-x^2)P_n^2(x) dx} \\ &< \frac{n}{2} \cdot \frac{(n+1)(2n+3)}{2(2n+1)}. \end{aligned}$$

This completes the proof of Theorem 3.

5. Proof of Theorem 4. On integrating by parts

$$\begin{aligned} \int_0^\infty P_n''(x)P_n(x)e^{-x} dx &= - \int_0^\infty P_n'(x)\{P_n'(x)x + P_n(x) - xP_n(x)\}e^{-x} dx \\ &= - \int_0^\infty xe^{-x}(P_n'(x))^2 dx + \frac{1}{2}P_n^2(0) \\ &\quad + \frac{1}{2} \int_0^\infty xe^{-x}P_n^2(x) dx - \int_0^\infty e^{-x}P_n^2(x) dx. \quad (5.1) \end{aligned}$$

Since $x_k < 0$, $k = 1, 2, \dots, n$, we have

$$\begin{aligned} \int_0^\infty x e^{-x} P_n^2(x) \sum_{k=1}^n \frac{1}{(x - x_k)^2} dx &< \int_0^\infty e^{-x} P_n^2(x) \sum_{k=1}^n \frac{1}{x - x_k} dx \\ &= \int_0^\infty e^{-x} P_n(x) P_n'(x) dx \\ &= -\frac{1}{2} P_n^2(0) + \frac{1}{2} \int_0^\infty P_n^2(x) e^{-x} dx. \end{aligned} \quad (5.2)$$

On using (5.1) and (5.2) and the identity

$$P_n'^2(x) = P_n(x) P_n''(x) + P_n^2(x) \sum_{k=1}^n \frac{1}{(x - x_k)^2}$$

we obtain

$$2 \int_0^\infty P_n'^2(x) x e^{-x} dx < \frac{1}{2} \int_0^\infty x e^{-x} P_n^2(x) dx - \frac{1}{2} \int_0^\infty e^{-x} P_n^2(x) dx. \quad (5.3)$$

It is easy to see that under the conditions of our theorem

$$P_n^2(x) = \sum_{k=0}^{2n} a_k x^k, \quad a_k > 0.$$

Hence

$$\int_0^\infty e^{-x} P_n^2(x) dx = \sum_{k=0}^{2n} a_k \int_0^\infty e^{-x} x^k dx = \sum_{k=0}^{2n} a_k k!,$$

similarly

$$\begin{aligned} \int_0^\infty e^{-x} x P_n^2(x) dx &= \sum_{k=0}^{2n} a_k (k+1)! \\ &< (2n+1) \sum_{k=0}^{2n} a_k k! = (2n+1) \int_0^\infty e^{-x} P_n^2(x) dx. \end{aligned}$$

On using (5.3) and the above result it follows that

$$2 \int_0^\infty (P_n'(x))^2 x e^{-x} dx < \left(\frac{1}{2} - \frac{1}{(2n+1)2} \right) \int_0^\infty P_n^2(x) x e^{-x} dx.$$

From this Theorem 4 now follows.

REFERENCES

1. P. Erdős, *On extremal properties of the derivatives of polynomials*, Ann. of Math. (2) **41** (1941), 310–313.
2. G. G. Lorentz, *The degree of approximation by polynomials with positive coefficients*, Math. Ann. **151** (1963) 239–251.
3. P. Turán, *Über die Ableitung von Polynomen*, Compositio Math. **7** (1939), 89–95.
4. A. K. Varma, *An analogue of some inequalities of P. Turán*, Proc. Amer. Math. Soc. **65** (1976), 305–309.
5. ———, *An analogue of some inequalities of P. Turán. II*, Proc. Amer. Math. Soc. **69** (1978), 25–33.