MODULAR GROUP ALGEBRAS OF TOTALLY PROJECTIVE p -PRIMARY GROUPS

WARREN MAY

ABSTRACT. Let F be a field of characteristic $p > 0$ and let G be a totally projective abelian p -group of countable p -length. If $FG \cong FH$ for some group H , then it is shown that $G \cong H$.

Let F be a field of characteristic $p > 0$ and let G be an (infinite) abelian p -group. The question of whether $FG \cong FH$ implies that $G \cong H$ has been considered by Berman and Mollov [1] and May [5]. (For the finite non-abelian case, see Passman [8].) It is shown in [1], [5] and by Dubois and Sehgal [2] that the Ulm-Kaplansky invariants of G and H must be equal. Thus one can easily settle the case when G is countable. Since the totally projective p -groups form the largest natural class of abelian p -groups that are determined by their Ulm-Kaplansky invariants, one would like to know in general whether H must be totally projective if G is. In this direction, Berman and Mollov give an affirmative answer if G is a direct sum of cyclic groups. By utilizing a theorem of Hill [4], we shall extend these results to p -groups of countable p -length. Precisely, we shall prove:

THEOREM 1. *Let R be a commutative ring with identity in which p is not invertible, and let Ω be the first uncountable ordinal. Let G be an abelian p -group of p -length $< \Omega$ whose reduced part is totally projective. If H is a group such that RG and RH are isomorphic algebras, then H is isomorphic to G .*

To deduce Theorem 1, we shall prove several facts about the group of units of RG . For any commutative ring R with identity and group G , let $U(RG)$ denote the group of units of RG that have augmentation 1 (i.e., the coefficients sum to 1). We prove:

THEOREM 2. *Let R be a commutative ring with identity of characteristic p that is perfect (i.e., every element is a p th power). Let G be an abelian p -group of p -length $\leq \Omega$ whose reduced part is totally projective. Then $U(RG)$ is an abelian p -group of the same p -length as G , and whose reduced part is totally projective. Moreover, G is a direct factor of $U(RG)$.*

We remark that Mollov [6], [7] has investigated the structure of unit groups under certain restrictions.

Received by the editors August 19, 1978.

AMS (MOS) subject classifications (1970). Primary 20K10; Secondary 16A26.

Key words and phrases. Totally projective p -groups, modular group algebras.

© 1979 American Mathematical Society
0002-9939/79/0000-0357/\$02.00

Before proving two preliminary lemmas, we must briefly discuss p -heights. Groups will be written multiplicatively, thus we shall use G^{p^σ} (σ an ordinal) to denote the subgroup of G consisting of elements with p -heights $\geq \sigma$. Suppose that R is a commutative ring with identity of characteristic p , and let G be an abelian p -group. Then $U(RG)$ is a p -group and consists precisely of the elements of RG of augmentation 1. Suppose further now that R is perfect. Then $U(RG)^p = U(R(G^p))$, thus it follows that $U(RG)^{p^\sigma} = U(R(G^{p^\sigma}))$ for every ordinal σ . Let $\alpha \in U(RG)$ and consider the support of α , i.e. those $g \in G$ that appear nontrivially in α . Clearly the p -height of α in $U(RG)$ is the minimum of the p -heights in G of the elements in the support of α . In particular, if we regard G as a subgroup of $U(RG)$, then it is an isotype subgroup. We also observe that the p -length of $U(RG)$ is clearly equal to the p -length of G .

LEMMA 1. *Let R be a perfect commutative ring with identity of characteristic p , and let G be an abelian p -group.*

(a) *Let A be a finite subgroup of G , and let N be a subgroup of $U(RA)$. Then N is a nice subgroup of $U(RG)$.*

(b) *If G is countable, then the reduced part of $U(RG)$ is totally projective.*

PROOF. Let $\alpha \in U(RG)$ and let $S = \{ga \mid g \in (\text{support } \alpha), a \in A\}$. If $\beta \in N$, then the p -height of $\alpha\beta$ equals the p -height of some element in the finite set S . Therefore β can be chosen to maximize this p -height. Hence N is nice in $U(RG)$.

Now suppose that G is countable, and let $A_1 \subseteq A_2 \subseteq \cdots$ be finite subgroups of G with $\bigcup_{i < \omega} A_i = G$. For each i , we can choose a family $\{N_{i\alpha}\}$ of subgroups of $U(RG)$ indexed by an initial segment of ordinals such that $U(RA_i) \subseteq N_{i\beta} \subseteq N_{i\alpha} \subseteq U(RA_{i+1})$ if $\beta < \alpha$, $N_{i\alpha} = \bigcup_{\beta < \alpha} N_{i\beta}$ if α is a limit ordinal,

$$N_{i0} = U(RA_i), \quad \bigcup_{\alpha} N_{i\alpha} = U(RA_{i+1}), \quad \text{and} \quad (N_{i,\alpha+1} : N_{i\alpha}) = p.$$

Each $N_{i\alpha}$ is nice in $U(RG)$ by (a), therefore $\bigcup_{i < \omega} \{N_{i\alpha}\}$ is a nice decomposition series for $U(RG)$. It follows that the reduced part of $U(RG)$ is totally projective.

LEMMA 2. *Let R be a commutative ring with identity and let G be an abelian group. Suppose that there exists a group B such that G is isomorphic to a direct factor of $U(RB)$. Then G is a direct factor of $U(RG)$.*

PROOF. Let G be isomorphic to the direct factor V of $U(RB)$. Regarding V as a subgroup of $U(RB)$, the isomorphism $G \rightarrow V$ induces a ring homomorphism $RG \rightarrow RB$, and hence a homomorphism $U(RG) \rightarrow U(RB)$. By composing with a projection of $U(RB)$ onto V , we obtain a homomorphism $U(RG) \rightarrow V$ that maps G isomorphically to V . Thus G is a direct factor of $U(RG)$.

PROOF OF THEOREM 2. We have already observed that $U(RG)$ is an abelian

p -group of the same p -length as G . We shall now show that the reduced part of $U(RG)$ is totally projective by using induction on $|G|$. The countable case is done by Lemma 1, therefore we may suppose that G is uncountable. The assumption that G has totally projective reduced part of p -length $\leq \Omega$ means precisely that G is isomorphic to a direct sum of countable p -groups (see [3, Theorem 82.4]). Let τ be the first ordinal with $|\tau| = |G|$. We may assume that $G = \prod_{\beta < \tau} A_\beta$, where each A_β is countable. For each $\alpha < \tau$, put $G_\alpha = \prod_{\beta < \alpha} A_\beta$ and $U_\alpha = U(RG_\alpha)$. (Note that $G_0 = 1$.) The projection $G_{\alpha+1} \rightarrow G_\alpha$ with kernel A_α induces a surjective map $U_{\alpha+1} \rightarrow U_\alpha$ that is idempotent. Thus we obtain an inner direct splitting $U_{\alpha+1} = U_\alpha \times K_\alpha$. It is easily seen that $U(RG) = \prod_{\alpha < \tau} K_\alpha$. But $|\alpha + 1| < |G|$, thus $|U_{\alpha+1}| < |G|$, and consequently the reduced part of $U_{\alpha+1}$ is totally projective by induction. It now follows that $U(RG)$ has totally projective reduced part.

To show that G is a direct factor of $U(RG)$ let $B = \prod_{i < \omega} G$. Then the reduced part of $U(RB)$ is totally projective by what we have just shown. Let $f_\sigma(G)$ denote the σ th Ulm-Kaplansky invariant of G . We know that $f_\sigma(B) < f_\sigma(U(RB))$ since B is isotype in $U(RB)$, therefore $f_\sigma(U(RB)) = f_\sigma(G) + f_\sigma(U(RB))$ for every ordinal σ . Moreover, if D_G and D_U are the maximal divisible subgroups of G and $U(RB)$, respectively, then $D_U \cong D_G \times D_U$. Hence, $U(RB) \cong G \times U(RB)$ since we are dealing with totally projective reduced parts. By Lemma 2, we conclude that G is a direct factor of $U(RG)$.

Since the complement of G in $U(RG)$ has totally projective reduced part, one could determine it up to isomorphism by computing the rank of its maximal divisible subgroup and its Ulm-Kaplansky invariants in terms of R and the Ulm-Kaplansky invariants of G . We remark that this is a feasible computation, but we shall not discuss it here.

PROOF OF THEOREM 1. Our assumptions on R guarantee that there is a homomorphism (not necessarily surjective) $R \rightarrow F$, where F is an algebraically closed field of characteristic p . It follows that $FG \cong FH$. One easily sees that H must be an abelian p -group since a torsion-free element in H would be transcendental over F , while an element of order relatively prime to p would yield a nontrivial idempotent in FH . Moreover, the maximal divisible subgroups of G and H are isomorphic by reference to [5, Corollary 7].

We may assume that the isomorphism $FG \cong FH$ preserves augmentations, hence $U(FG) \cong U(FH)$. Theorem 2 allows us to conclude that the reduced part of $U(FH)$ is totally projective of countable p -length. Thus the reduced part of H is isomorphic to an isotype subgroup of countable p -length in the reduced part of $U(FH)$. It now follows immediately from a result of Hill [4, Theorem 1] that the reduced part of H is totally projective. As was mentioned in the introduction, the Ulm-Kaplansky invariants of G and H are known to be equal; therefore $G \cong H$.

We conclude with several questions and remarks. Does Theorem 1 hold if G has p -length $\geq \Omega$? In this case, we no longer have Hill's theorem to apply.

Even more, there is uncertainty whether Theorem 2 holds if the p -length is $> \Omega$ since G would not be a direct sum of countable groups. If one tries to prove Theorem 2 by induction on the p -length of G , the difficult step seems to occur at isolated ordinals. As a final question, if G is a torsion-complete abelian p -group, then is H torsion-complete? If so, then one could conclude G and H would be isomorphic since the Ulm-Kaplansky invariants serve to classify torsion-complete groups.

REFERENCES

1. S. D. Berman and T. Zh. Mollov, *The group rings of abelian p -groups of arbitrary power* (Russian), *Mat. Zametki* **6** (1969), 381–392; *Math. Notes* **6** (1969), 686–692. MR **40** #7365.
2. P. F. Dubois and S. K. Sehgal, *Another proof of the invariance of Ulm's functions in commutative modular group rings*, *Math. J. Okayama Univ.* **15** (1972), 137–139. MR **47** #6892.
3. L. Fuchs, *Infinite abelian groups*, vol. II, Academic Press, New York, 1973.
4. P. Hill, *Isotype subgroups of direct sums of countable groups*, *Illinois J. Math.* **13** (1969), 281–290. MR **39** #1550.
5. W. May, *Commutative group algebras*, *Trans. Amer. Math. Soc.* **136** (1969), 139–149. MR **38** #2224.
6. T. Zh. Mollov, *The multiplicative groups of modular group algebras of primary abelian groups of arbitrary cardinality*. I (Russian), *Publ. Math. Debrecen* **18** (1971), 9–21 (1972). MR **47** #341; II (Russian), *ibid.* **19** (1972), 87–96 (1973). MR **49** #2980.
7. ———, *Sylow p -subgroups of group algebras of countable abelian groups* (Russian), *C. R. Acad. Bulgare Sci.* **27** (1974), 733–735. MR **51** #5805.
8. D. S. Passman, *Advances in group rings*, *Israel J. Math.* **19** (1974), 67–107. MR **50** #9945.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ARIZONA, TUCSON, ARIZONA 85721