NEAR-DERIVATIONS AND INFORMATION FUNCTIONS¹

JOHN LAWRENCE², GEOFF MESS AND FRANK ZORZITTO³

ABSTRACT. Near-derivations γ satisfy the conditions $\gamma(xy) = x\gamma(y) + y\gamma(x)$ for $x, y \in R$, $\gamma(x + y) \geq \gamma(x) + \gamma(y)$ for $x, y \geq 0$, $\gamma(x) = 0$ for $x \in \mathbb{Q}$. Existence of near-derivations other than derivations is tied in with that of nonnegative information functions and an example of Daróczy and Maksa. Conditions for near-derivations to be derivations are discussed.

1. Introduction. A *near-derivation* is a self-mapping γ of the reals R such that:

$$\gamma(xy) = x\gamma(y) + y\gamma(x) \quad \text{for all } x, y \text{ in } R, \quad (1)$$

$$\gamma(x + y) \geq \gamma(x) + \gamma(y) \quad \text{for all } x, y \geq 0, \quad (2)$$

$$\gamma(x) = 0 \quad \text{for all rational } x. \quad (3)$$

The question immediately arises as to whether there are any near-derivations that are not already derivations.

An *information function* is a mapping $f: [0, 1] \rightarrow R$ satisfying the equation

$$f(x) + (1 - x)f(y/(1 - x)) = f(y) + (1 - y)f(x/(1 - y)),$$

and the normalizing conditions

$$f(0) = f(1) = 0, \quad f\left(\frac{1}{2}\right) = 1.$$

The most important example of an information function is the Shannon function s given by

$$s(x) = -x \log x - (1 - x) \log(1 - x) \quad \text{for } 0 < x < 1.$$

(Here and throughout we adopt the conventions that $\log \equiv \log_2$ and $0 \log 0 = 0$.) The motivation for the above definition of an information function is developed by Aczél and Daróczy in [1], where such functions are studied extensively as part of an analysis of measures of information.

The present research was motivated by a problem cited in [1]. Namely, is the Shannon function s the only nonnegative information function? This was recently solved by Daróczy and Maksa [2]. They showed that for any

Received by the editors October 2, 1978.

AMS (MOS) subject classifications (1970). Primary 39A15; Secondary 94A15.

¹This research was carried out partly in the NRC Summer Research workshops held at the University of Waterloo in 1978.

²Research supported by NRC grant A4540 and University of Waterloo research grant 131-7052.

³Research supported by NRC grant A4655 and University of Waterloo research grant 131-7067.

derivation δ on R , the formula

$$f(x) = s(x) + \delta(x)^2/x(1-x)$$

provides a nonnegative information function. In our paper we show that f is a nonnegative information function if and only if

$$f(x) = s(x) - \gamma(x) - \gamma(1-x),$$

where γ is a near-derivation. It follows from this that s is the minimal nonnegative information function. The latter fact is also proved in [2] by a different approach. In addition the Daróczy-Maksa example leads to examples of near-derivations which are not derivations. We conclude with some properties of near-derivations.

2. Connection of near-derivations to information functions. A theorem in [1] leads to the connection between information functions and near-derivations.

REPRESENTATION THEOREM (ACZÉL-DARÓCZY). *A function $f: [0, 1] \rightarrow R$ is an information function if and only if there is a self-mapping δ of R such that*

$$f(x) = -\delta(x) - \delta(1-x) \quad \text{for all } x \text{ in } [0, 1], \quad (4)$$

$$\delta(xy) = x\delta(y) + y\delta(x) \quad \text{for all } x, y \text{ in } R, \quad (5)$$

$$\delta(2) = 2. \quad (6)$$

In addition it can be readily checked that f is nonnegative if and only if the representing function δ satisfies

$$\delta(x+y) \geq \delta(x) + \delta(y) \quad \text{for all } x, y \geq 0. \quad (7)$$

Our first result is a refinement of the representation theorem for nonnegative f .

THEOREM 1. *A function $f: [0, 1] \rightarrow R$ is a nonnegative information function if and only if*

$$f(x) = s(x) - \gamma(x) - \gamma(1-x), \quad (8)$$

where γ is a near-derivation.

PROOF. Let γ be a near-derivation. Let $\sigma: R \rightarrow R$ be defined by $\sigma(x) = x \log|x|$, and let $\delta = \sigma + \gamma$. Clearly σ satisfies conditions (5)–(7). Since γ is a near-derivation, δ satisfies (5)–(7) also. By the representation theorem, the formula $f(x) = -\delta(x) - \delta(1-x)$, $0 \leq x \leq 1$, provides a nonnegative information function f .

Conversely, let f be a nonnegative information function. By the representation theorem let $\delta: R \rightarrow R$ satisfy (4)–(7) for f . Let $\sigma(x) = x \log|x|$ as before, and let $\gamma = \delta - \sigma$. We shall prove γ is a near-derivation. It will then follow that

$$f(x) = -(\sigma + \gamma)(x) - (\sigma + \gamma)(1-x) = s(x) - \gamma(x) - \gamma(1-x)$$

as desired.

Now, property (1) of near-derivations holds for γ because δ and σ have it. Property (3) for γ follows from [3, Theorem 2], where it is shown that $\delta = \sigma$ on the rationals. Thus property (2) remains, namely that

$$\begin{aligned} \delta(x+y) - (x+y)\log(x+y) \\ \geq (\delta(x) - x \log x) + (\delta(y) - y \log y) \end{aligned} \quad (9)$$

for all $x, y \geq 0$. This inequality will follow from the subsequent lemmas.

LEMMA 2. Suppose $\delta: R \rightarrow R$ satisfies conditions (5)–(7). Then

$$\delta(1+x) \geq (1+x)\log(1+x) + \delta(x) - x \log x$$

for any $x \geq 0$.

PROOF. For any positive integer n and $x \geq 0$ we apply (5) and (7) to get

$$n(1+x)^{n-1}\delta(1+x) = \delta((1+x)^n) = \delta\left(\sum_{k=0}^n \binom{n}{k} x^k\right) \geq \sum_{k=0}^n \delta\left(\binom{n}{k} x^k\right).$$

Thus

$$\delta(1+x) \geq \frac{1}{n(1+x)^{n-1}} \sum_{k=0}^n \left(\binom{n}{k} \delta(x^k) + x^k \delta\left(\binom{n}{k}\right) \right).$$

In [3] it is proved that $\delta(r) = r \log r$ for any positive rational r . Hence

$$\begin{aligned} \delta(1+x) &\geq \frac{1}{n(1+x)^{n-1}} \sum_{k=0}^n \binom{n}{k} k x^{k-1} \delta(x) \\ &\quad + \frac{1}{n(1+x)^{n-1}} \sum_{k=0}^n x^k \binom{n}{k} \log\left(\binom{n}{k}\right) \\ &= \delta(x) + \frac{1+x}{n} \sum_{k=0}^n \binom{n}{k} \frac{x^k}{(1+x)^n} \log\left(\binom{n}{k}\right) \\ &= \delta(x) + a_n - b_n + c_n, \end{aligned}$$

where

$$\begin{aligned} a_n &= \frac{1+x}{n} \sum_{k=0}^n \binom{n}{k} \frac{x^k}{(1+x)^n} \log\left(\binom{n}{k} \frac{x^k}{(1+x)^n}\right), \\ b_n &= \frac{1+x}{n} \sum_{k=0}^n \binom{n}{k} \frac{x^k}{(1+x)^n} \log(x^k) = x \log x, \\ c_n &= \frac{1+x}{n} \sum_{k=0}^n \binom{n}{k} \frac{x^k}{(1+x)^n} \log((1+x)^n) = (1+x)\log(1+x). \end{aligned}$$

Since

$$\sum_{k=0}^n \binom{n}{k} \frac{x^k}{(1+x)^n} = 1$$

it follows from [1, Theorem 1.3.7] that

$$|a_n| \leq \frac{1+x}{n} \log(n+1).$$

Thus $a_n \rightarrow 0$ as $n \rightarrow \infty$. We therefore get in the limit

$$\delta(1+x) \geq \delta(x) - x \log x + (1+x) \log(1+x).$$

LEMMA 3. *For δ satisfying (5)–(7) and any $x, y \geq 0$, inequality (9) must hold.*

PROOF. By using property (5) and Lemma 2 we get

$$\begin{aligned} \delta(x+y) &= \delta(x(1+y/x)) = x\delta(1+y/x) + (1+y/x)\delta(x) \\ &\geq x((1+y/x)\log(1+y/x)) + \delta(y/x) \\ &\quad - (y/x)\log(y/x) + (1+y/x)\delta(x). \end{aligned}$$

Use of (5) to expand $\delta(y/x)$ and some logarithmic properties reduces the latter expression to

$$(x+y)\log(x+y) + \delta(x) - x \log x + \delta(y) - y \log y,$$

whence (9) follows.

THEOREM 4. *If f is any nonnegative information function and s is the Shannon function, then $f \geq s$.*

PROOF. Let δ represent f according to the representation theorem. By Lemma 3, inequality (9) holds for δ . With $0 \leq x \leq 1$ and $y = 1 - x$, (9) specializes to

$$\delta(x) + \delta(1-x) \leq x \log x + (1-x) \log(1-x),$$

which says that $f \geq s$.

THEOREM 5. *Not all near-derivations must be derivations.*

PROOF. Formula (8) yields an example of such near-derivations for every nonnegative information function f different from s .

3. Some properties of near-derivations. We are interested in conditions for the additivity of near-derivations. For example, it can be seen from [3] that if F, K are fields of real numbers and K is quadratic over F and γ is a near-derivation vanishing on F , then γ is additive on K . By Theorem 5 this need not happen if K is not algebraic over F . If, for a transcendental x , a near-derivation γ vanishing on F were linear on the field $F(x)$, then γ would be constant on the coset $x + F$. We are therefore led to examine γ restricted to $x + F$.

THEOREM 6. *Let F be a real field, x a real number and γ a near-derivation vanishing on F . Then for a, b in F :*

- (a) $0 \leq x + a \leq x + b$ implies that $\gamma(x + a) \leq \gamma(x + b)$;
- (b) $x + a \leq x + b \leq 0$ implies that $\gamma(x + a) \leq \gamma(x + b)$;
- (c) $x + a \leq 0 \leq x + b$ implies that $\gamma(x + b) \leq \gamma(x + a)$;

(d) $\gamma(x + a)$ and $\gamma(x - a)$ converge to the same finite value as a tends to $+\infty$ inside F .

PROOF. To prove (a) we use (2) and the vanishing of γ on F . This yields:

$$\gamma(x + b) = \gamma(x + a + b - a) \geq \gamma(x + a) + \gamma(b - a) = \gamma(x + a).$$

Items (b) and (c) similarly use (2) and the fact that $\gamma = 0$ on F , plus that near-derivations are odd functions. The existence of the finite limits

$$\beta_l(x) = \lim_{\substack{a \rightarrow +\infty \\ a \in F}} \gamma(x - a), \quad \beta_r(x) = \lim_{\substack{a \rightarrow +\infty \\ a \in F}} \gamma(x + a)$$

follows from the monotonicity conditions (a) and (b), and the bounding condition (c). It also follows from (c) that $\beta_r(x) \leq \beta_l(x)$.

To get equality for β_r and β_l consider the finite quantity $\beta_l(x^2)$ and apply (1).

$$\begin{aligned} \beta_l(x^2) &= \lim_{\substack{a \rightarrow +\infty \\ a \in F}} \gamma(x^2 - a) = \lim_{\substack{a \rightarrow +\infty \\ a \in F}} \gamma(x^2 - a^2) \\ &= \lim_{\substack{a \rightarrow +\infty \\ a \in F}} ((x + a)\gamma(x - a) + (x - a)\gamma(x + a)) \\ &= x\beta_l(x) + x\beta_r(x) + \lim_{\substack{a \rightarrow +\infty \\ a \in F}} a(\gamma(x - a) - \gamma(x + a)). \end{aligned}$$

The limit of $a(\gamma(x - a) - \gamma(x + a))$ is now finite. This means $\gamma(x - a) - \gamma(x + a)$ tends to 0, and hence $\beta_l(x) = \beta_r(x)$.

The next result gives a condition for our γ to be a derivation.

THEOREM 7. *Let γ be a near-derivation, and let the function β be defined by $\beta(x) = \lim_{a \rightarrow +\infty; a \in Q} \gamma(x + a)$, x real. Then β is Q -linear, and, for all real x , $\beta(x^2) \geq 2x\beta(x)$. Furthermore, β is a derivation if and only if $\beta = \gamma$.*

PROOF. The superadditivity (2) of γ implies the same for β as follows:

$$\begin{aligned} \beta(x + y) &= \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} \gamma(x + y + a) = \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} \gamma(x + a/2 + y + a/2) \\ &\geq \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} \gamma(x + a/2) + \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} \gamma(y + a/2) = \beta(x) + \beta(y). \end{aligned}$$

The subadditivity of β comes from the equality of β and β_l in Theorem 6(d), and the fact that, for $u > v > 0$, $\gamma(u - v) \leq \gamma(u) + \gamma(-v)$. Indeed,

$$\begin{aligned} \beta(x + y) &= \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} \gamma((x + 2a) - (-y + a)) \\ &\leq \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} \gamma(x + 2a) + \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} \gamma(y - a) = \beta(x) + \beta(y). \end{aligned}$$

Next, for x real use the equality in the proof of Theorem 6(d) to obtain:

$$\beta(x^2) = 2x\beta(x) + \lim_{\substack{a \rightarrow +\infty \\ a \in Q}} a(\gamma(x - a) - \gamma(x + a)). \quad (10)$$

Theorem 6(c) implies that $\gamma(x - a) \geq \gamma(x + a)$ when a is large. Thus $\beta(x^2) > 2x\beta(x)$.

Now if $\beta = \gamma$, then both are derivations because β is additive and γ satisfies (1). Conversely suppose β is a derivation. The difference $\gamma_1 = \gamma - \beta$ is then a near-derivation for which Theorem 6 applies. In particular, for any $x \geq 0$, $\gamma_1(x + a)$ increases to a finite value $\beta_1(x)$ as $a \rightarrow +\infty$ through the rationals. By the definition of γ_1 it follows that $\beta_1(x) = 0$. Thus $\gamma_1(x) \leq 0$ for all $x \geq 0$. However, one can check that any function satisfying (1) and (3) must have a dense graph in R^2 or else vanish everywhere. We must conclude that $\gamma_1 = 0$, or $\beta = \gamma$.

A corollary of Theorem 7 seems worth noting.

THEOREM 8. A near-derivation γ is a derivation if and only if, for all x , $a(\gamma(x - a) - \gamma(x + a)) \rightarrow 0$ as $a \rightarrow +\infty$ through the rationals.

PROOF. By formula (10), $\beta(x^2) = 2x\beta(x)$. This makes β a derivation, and hence γ .

REFERENCES

1. J. Aczél and Z. Daróczy, *On measures of information and their characterizations*, Academic Press, New York, 1975.
2. Z. Daróczy and Gy. Maksa, Nonnegative information functions (to appear).
3. J. Lawrence, *The uniqueness of the nonnegative information function on algebraic extensions*, (to appear).

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA N2L 3G1