A SCHRÖDER-BERNSTEIN THEOREM IN BAER*-RINGS WITH LATTICE-THEORETIC PROOF

JOSUKE HAKEDA

ABSTRACT. The Schröder-Bernstein theorem (SB-theorem) for *-equivalence of projections of a Baer*-ring is known.

Here, we will prove an SB-theorem for algebraic equivalence (Theorem A) as a consequence of a lattice-theoretic SB-theorem (Theorem B). Theorem A and the known result about *-equivalence will be derived from Theorem B.

An element e of an involutive ring A is called a *projection* if it is idempotent ($e^2 = e$) and selfadjoint ($e^* = e$). The relation $e = ef$ defines a partial ordering of projections, denoted $e \leq f$. Projections e, f of A are said to be *algebraically equivalent* if there exist x, y in A such that $yx = e$ and $xy = f$, and **-equivalent* if there exists x in A such that $x^*x = e$ and $xx^* = f$. We write $e \sim f$ for either of these relations, indicating in the context which of the relations is intended. The notation $e \lesssim f$ means that $e \sim f_0$ for some projection $f_0 \leq f$.

An involutive ring A is called a *Baer*-ring* if, for every subset S of A , the right annihilator of S has the form eA , e a projection [3].

THEOREM A. *Let e, f be projections in a Baer*-ring A . If $e \lesssim f$ and $f \lesssim e$ for algebraic equivalence, then $e \sim f$.*

The analogue of Theorem A for *-equivalence is known [4, Theorem]. We shall give a lattice-theoretic proof of Theorem A.

DEFINITION. An orthocomplemented lattice L is called *orthomodular* if its orthocomplementation $x \mapsto \theta(x)$ ($x \in L$) satisfies the identity

$$b = a \vee (b \wedge \theta(a))$$

for $a \leq b$ ($a, b \in L$).

LEMMA [5, Lemma 29.15]. *Let L be an orthomodular lattice with orthocomplementation $x \mapsto \theta(x)$ ($x \in L$). For $a \in L$, $[0, a] = \{x \in L: x \leq a\}$ is an orthomodular lattice for the orthocomplementation*

$$x \mapsto \theta_a(x) = a \wedge \theta(x)$$

for $x \in [0, a]$. If $a \leq \theta(b)$, we write $a \perp b$.

Received by the editors March 28, 1978 and, in revised form, August 7, 1978.

AMS (MOS) subject classifications (1970). Primary 46K99, 46L99, 06A25, 06A23; Secondary 16A28, 16A32.

Key words and phrases. Schröder-Bernstein theorem, Baer*-ring, orthomodular lattice, complete lattice.

THEOREM B. Suppose the orthomodular lattice L is complete and possesses an equivalence relation $\sim$ satisfying the following conditions: (1) if $a_1 \perp a_2$, $b_1 \perp b_2$ and $a_i \sim b_i$ ($i = 1, 2$), then $a_1 \vee a_2 \sim b_1 \vee b_2$;

(2) If $a \lesssim b$, then there exists an order-preserving mapping φ of $[0, a]$ into $[0, b]$ such that $x \sim \varphi(x)$ for all $x \in [0, a]$.

Then $a \lesssim b$ and $b \lesssim a$ imply $a \sim b$ for $a, b \in L$.

PROOF. From condition (2), there exist order-preserving mappings φ_1 and φ_2 such that φ_1 maps $[0, a]$ into $[0, b]$ and φ_2 maps $[0, b]$ into $[0, a]$. Define a mapping φ of $[0, b]$ into $[0, b]$ by the formula

$$\varphi = \theta_b \circ \varphi_1 \circ \theta_a \circ \varphi_2.$$

By the lemma, φ is an order-preserving mapping in the complete lattice $[0, b]$; therefore, by [7, Theorem 1], there exists $c_0 \in [0, b]$ such that $\varphi(c_0) = c_0$. Since $\theta_b \circ \theta_b = 1$,

$$\theta_b(c_0) = \theta_b(\varphi(c_0)) = \varphi_1(\theta_a(\varphi_2(c_0))).$$

By condition (2), one has

$$\theta_b(c_0) = \varphi_1(\theta_a(\varphi_2(c_0))) \sim \theta_a(\varphi_2(c_0))$$

and $c_0 \sim \varphi_2(c_0)$. Therefore, by the condition (1),

$$b = c_0 \vee \theta_b(c_0) \sim \varphi_2(c_0) \vee \theta_a(\varphi_2(c_0)) = a. \quad \text{Q.E.D.}$$

PROOF OF THEOREM A. The projections of A form a complete orthomodular lattice with the orthocomplementation $e \mapsto 1 - e$. Algebraic equivalence satisfies condition (1) [3, Theorem 17], and condition (2) as a special case of [6, Lemma 2.2]. Quote Theorem B. Q.E.D.

REMARK 1. *-equivalence satisfies condition (1) [3, Theorem 25] and condition (2) [3, Theorem 24]. Therefore the SB-theorem for *-equivalence [4, Theorem] holds by Theorem B.

REMARK 2. By the order of set inclusion and orthocomplementation of usual complement, the family of all subsets of a set forms a complete orthomodular lattice. With slight modification, Theorem B gives a proof of the Bernstein theorem of set theory.

The author thanks Professor S. Maeda for his kind advice.

REFERENCES

1. S. K. Berberian, *On the projection geometry of a finite AW*-algebra*, Trans. Amer. Math. Soc. **83** (1956), 493–508.
2. ———, *Baer*-rings*, Springer-Verlag, Berlin, 1972.
3. I. Kaplansky, *Rings of operators*, Benjamin, New York, 1968.
4. A. Lebow, *A Schroder-Bernstein theorem for projections*, Proc. Amer. Math. Soc. **19** (1968), 144–145.
5. F. Maeda and S. Maeda, *Theory of symmetric lattices*, Springer-Verlag, Berlin, 1970.
6. S. Maeda, *On the lattice of projections of a Baer*-ring*, J. Sci. Hiroshima Univ. Ser. A **22** (1958), 75–88.
7. A. Tarsky, *A lattice-theoretical fixpoint theorem and its applications*, Pacific J. Math. **5** (1955), 285–309.

DEPARTMENT OF BASIC TECHNOLOGY, YAMAGATA UNIVERSITY, YONEZAWA, JAPAN