SURFACES IN THE GRASSMANN VARIETY $G(1, 3)$

AIGLI PAPANTONOPOULOU

ABSTRACT. Let G be the Grassmannian $G(1, 3)$, Y a nonsingular subvariety of G and $N_{Y/G}$ the normal bundle of Y in G . Then $N_{Y/G}$ is not an ample bundle if and only if Y is a Schubert cycle.

1. Introduction. In [8] we gave a complete characterization of nonsingular curves in any Grassmannian $G(r, n)$ with ample normal bundle. In the case of $G(1, 3)$ we showed that the Schubert cycle of dimension 1, which is a line, is the only nonsingular curve with normal bundle not ample. In this paper, using the intersection theories developed in [1] and in [9] we show that in dimension 2 also the Schubert cycles, which are isomorphic to $\mathbf{P}^2$, are the only nonsingular surfaces with normal bundle not ample. In the case of codimension 1 the normal bundle is always ample because the $\text{Pic}(G)$ is trivial. Knowing that a nonsingular subvariety of $G(1, 3)$ has an ample normal bundle we can apply on it several well-known theorems: (a) A vanishing theorem on formal schemes [4, Theorem 4.1], (b) A theorem on meromorphic functions [5, Chapter 6], (c) Results on the cohomological dimension of a projective variety minus a subvariety [5, Chapter 7].

Throughout the paper we are working over an algebraically closed field of characteristic 0.

2. Preliminaries. Let $G = G(1, 3)$ be the Grassmannian parametrizing $\mathbf{P}^1$ spaces in a fixed $\mathbf{P}^3$. Then $\dim G = 4$ and G is a quadric in $\mathbf{P}^5$. The Schubert cycles of G can be described as follows:

(a) $Z_0 = G \cap H$, H a hyperplane in $\mathbf{P}^5$, $\dim Z_0 = 3$,

$$Z_0 = \{\mathbf{P}^1 \subset \mathbf{P}^3 \mid \mathbf{P}^1 \cap \text{some fixed } \mathbf{P}^1 \neq \emptyset\}.$$

(b) $Z_1 = \{\mathbf{P}^1 \subset \mathbf{P}^3 \mid \mathbf{P}^1 \subset \text{some fixed } \mathbf{P}^1 \subset \mathbf{P}^3\} \cong \mathbf{P}^2$

$$Z_2 = \{\mathbf{P}^1 \subset \mathbf{P}^3 \mid \text{some fixed point } p \in \mathbf{P}^1 \subset \mathbf{P}^3\} \cong \mathbf{P}^1.$$

(c) $Z_3 = \{\mathbf{P}^1 \subset \mathbf{P}^3 \mid \text{some fixed point } p \in \mathbf{P}^1 \subset \text{some fixed } \mathbf{P}^2 \subset \mathbf{P}^3\} \cong \mathbf{P}^1$, and for any given Z_3 there are exactly two cycles Z_1 and Z_2 such that $Z_3 = Z_1 \cap Z_2 \subset G$.

Recall that on G we have the short exact sequence

$$0 \rightarrow E \rightarrow \mathcal{O}_G^4 \rightarrow Q \rightarrow 0,$$

where Q and E are the canonical bundles on G . The tangent bundle

Received by the editors July 12, 1978 and, in revised form, November 23, 1978.

AMS (MOS) subject classifications (1970). Primary 14M15; Secondary 14J99, 14F05.

© 1979 American Mathematical Society
0002-9939/79/0000-0453/\$02.00

$T_G = Q \otimes \check{E}$, therefore for all nonsingular subvarieties of G the normal bundle $N_{Y/G}$ is generated by global sections. Hence by [2 Proposition 2.1] and [8, Chapter 2] if $N_{Y/G}$ is not ample then there exists a curve C in Y and a surjective map

$$N_{Y/G} \rightarrow 0_C \rightarrow 0.$$

We recall that for Y locally a complete intersection $N_{Y/G}$ is defined as $\text{Hom}_{0_Y}(I_Y/I_Y^2, 0_Y)$, where I_Y is the sheaf of ideals defining Y in G , and when Y is nonsingular $N_{Y/G}$ is simply the usual geometric normal bundle of Y in G .

Finally one more property of G that we need is the fact that the Chow ring $A(G)$ of cycles mod rational equivalence is generated by the Schubert cycles. For example a surface $Y \neq Z_i$, $Y \sim aZ_1 + bZ_2$ where a and b some positive integers, and $Z_1 \cdot Z_2 = 0$, $Z_i^2 = 1$ in $A(G)$.

We use in the proof the definition and properties of ample vector bundles which are shown in [3]. We also use repeatedly the following proposition from [8]:

PROPOSITION 2.4. *Let Y be a subvariety of $G(r, n)$. Then $T_{G|Y}$ is not ample if and only if there is a curve C in Y which lies in some Z_3 .*

In the proof of the theorem we will be using the following notation and diagrams:

Y = a nonsingular surface in $G(1, 3) = G$, Z = a Schubert cycle in G of dim 2, and $D = Y \cap Z$.

G' = the blow up of G with center Y , $p: G' \rightarrow G$ the projection,

$Y' = \mathbf{P}(\check{N}_{Y/G})$, Z' = the strict transform of Z , i.e. the blow up of Z with center D , and $D' = Y' \cap Z'$.

$$\begin{array}{ccccc}
 & & D' & \longrightarrow & Z' \\
 & \swarrow & \downarrow & \swarrow & \downarrow \\
 Y' & \xrightarrow{\tau} & G' & & \\
 \downarrow p & & \downarrow p & & \downarrow \\
 Y & \xrightarrow{\quad} & D & \longrightarrow & Z \\
 & \swarrow & \downarrow & \swarrow & \downarrow \\
 & & G & &
 \end{array} \tag{2.1}$$

G'' = the blow up of G with center Z , $q: G'' \rightarrow G$ the projection,

$Z'' = \mathbf{P}(\check{N}_{Z/G})$, Y'' = the strict transform of Y , i.e. the blow up of Y with center D , and $D'' = Y'' \cap Z'' \cong D'$.

$$\begin{array}{ccccc}
 & & D'' & \longrightarrow & Z'' \\
 & \swarrow & \downarrow & \swarrow & \downarrow \\
 Y'' & \xrightarrow{\tau} & G'' & & \\
 \downarrow q & & \downarrow q & & \downarrow \\
 Y & \xrightarrow{\quad} & D & \longrightarrow & Z \\
 & \swarrow & \downarrow & \swarrow & \downarrow \\
 & & G & &
 \end{array} \tag{2.2}$$

On Y' we have the canonical sequence

$$0 \rightarrow 0_{Y'}(-1) \rightarrow p^*(N_{Y/G}) \rightarrow F \rightarrow 0$$

and restricted to D' we get

$$0 \rightarrow N_{D'/Z'} \rightarrow p^*(N_{Y/G})|_{D'} \rightarrow F|_{D'} \rightarrow 0. \quad (1)$$

Also from the canonical sequence on Z'' restricted to D'' we get

$$0 \rightarrow N_{D''/Y''} \rightarrow q^*(N_{Z/G})|_{D''} \rightarrow E|_{D''} \rightarrow 0. \quad (2)$$

From [1] we have that in the Chow ring $A(D)$ the intersection cycle is given by $Y \cdot Z = \tau_* c_1(F|_{D'}) = \tau_* c_1(D|_{D''})$.

3. Nonsingular surfaces in $G(1, 3)$.

THEOREM. *Let Y be a nonsingular surface in $G = G(1, 3)$. $N_{Y/G}$ the normal bundle of Y in G is not ample if and only if $Y = Z_i$ (a Schubert cycle of dim 2).*

PROOF. If $Y = Z_i$ then $T_{G|Y} = T_Y \oplus Q|_Y$ or $T_{G|Y} = T_Y \oplus \check{E}|_Y$. By [8, 2.2 and 2.3]

$$Q|_{Z_3} \cong \check{E}|_{Z_3} \cong 0_{Z_3}(1) \oplus 0_{Z_3},$$

where $Z_3 \cong \mathbf{P}^1$. Hence $N_{Y/G}$ restricted to a cycle of type Z_3 is not ample, therefore $N_{Y/G}$ is not ample.

Now assume that $Y \neq Z_i$. $N_{Y/G}$ is a vector bundle on Y which is generated by its global sections, hence by [2, Proposition 2.1] it is sufficient to show that $N_{Y/G|C}$ is ample for every curve C in Y . By [8, Proposition 2.4] $T_{G|C}$ is not ample if and only if $C = Z_3 \cong \mathbf{P}^1$. Since $N_{Y/G|C}$ is a quotient bundle of $T_{G|C}$, it will be sufficient to show that for any cycle $Z \subset G$, $N_{Y/G|D}$ is ample, where $D = Y \cap Z$. Let I_D be the sheaf of ideals defining $D = Y \cap Z$ in Y . Then

$$\tau^*(I_D) \oplus 0_{D''} = \check{N}_{D''/Y''}.$$

For any curve C in D'' not in any fiber the map τ is of degree 1 and $\tau_*(\check{N}_{D''/Y''|C}) = \check{N}_{C/Y}$. For the rest of the proof let C be a cycle of type Z_3 . Since $Y \neq Z_i$, $\deg_G Y > 1$ and $(C \cdot C)_Y < 1$. Hence $\tau_*(N_{D''/Y''|C}) = 0_C(v)$ where $v < 1$, and the exact sequence (2) restricted to C becomes

$$0 \rightarrow 0_C(v) \rightarrow q^*(N_{Z/G})|_C \rightarrow E|_C \rightarrow 0.$$

The vector bundle

$$q^*(N_{Z/G})|_C = 0_C(1) \oplus 0_C,$$

therefore $E|_C$ is an ample line bundle on C . Hence the line bundle $E|_{D''}$ has a global section with zeroes on C . Since $\tau_* c_1(F|_{D'}) = \tau_* c_1(E|_{D''})$ we get a global section of the line bundle $F|_{D'}$ which is not constant on C . Hence $F|_C$ is an ample line bundle on C . Consider now the exact sequence (1). $\tau_*(N_{D'/Z'|C}) = N_{C/Z}$ where $C \cong \mathbf{P}^1$ and $Z \cong \mathbf{P}^2$. Hence $\tau_*(N_{D'/Z'|C}) = 0_C(1)$ and $N_{Y/G|C} = 0_C(u) \oplus 0_C(w)$ where $u > 0$ and $w > 0$. Hence $N_{Y/G|D}$ is ample. Q.E.D.

BIBLIOGRAPHY

1. W. Fulton and R. MacPherson, *Intersecting cycles on an algebraic variety*, Aarhus Univ., Aarhus, 1976 (preprint).
2. D. Gieseker, *p-ample bundles and their Chern classes*, Nagoya Math. J. **43** (1971), 91–116.
3. R. Harshorne, *Ample vector bundles*, Inst. Hautes Études Sci. Publ. Math., No. 29 (1966), 63–94.
4. ———, *Ample subvarieties of algebraic varieties*, Lecture Notes in Math., vol. 156, Springer-Verlag, Berlin and New York, 1970.
5. ———, *Cohomological dimension of algebraic varieties*, Ann. of Math. (2) **88** (1968), 403–450.
6. ———, *Ample vector bundles on curves*, Nagoya Math. J. **43** (1971), 71–89.
7. V. D. Hodge and D. Pedoe, *Methods of algebraic geometry*. Vols. I, II, III, Cambridge Univ. Press, 1947, 1952, 1954.
8. A. Papantonopoulou, *Curves in Grassmann varieties*, Nagoya Math. J. **66** (1977), 121–137.
9. ———, *Non-proper intersection of surfaces and applications*, Deltion Greek Math. Soc. **18** (1977), 80–93.

DEPARTMENT OF MATHEMATICS, BRYN MAWR COLLEGE, BRYN MAWR, PENNSYLVANIA 19010