TRIANGLES IN ARRANGEMENTS OF LINES. II

GEORGE PURDY

ABSTRACT. We show that given n lines in the real projective plane, no $n - 1$ of which are concurrent, the number p_3 of triangular regions formed is at most $\frac{2}{3}n(n - 1)$, equality being possible.

We also show that if $n > 6$ then $p_3 < \frac{7}{18}n(n - 1) + \frac{1}{3}$. Grünbaum has conjectured $p_3 < \frac{1}{3}n(n - 1)$.

1. Introduction. Let A be an arrangement of n lines in the real projective plane P_2 , and let p_k denote the number of k -sided regions determined by the lines. It is assumed that A is not a pencil. In [1] Grünbaum raises the question of finding an upper bound for p_3 and conjectures that

$$p_3 < \left[\frac{1}{3}n(2\lfloor n/2 \rfloor - 1) \right] < \frac{1}{3}n(n - 1)$$

for $n \geq 16$, and gives R. J. Canham's proof of this when A is simple (no three lines concurrent). The first nontrivial results on the general problem were found by T. O. Strommer [3] who proved among other things that

$$p_3 \leq \frac{1}{3}n(n - 1) + 2 + \frac{1}{3} \sum_{k \geq 4} (k - 6)p_k.$$

In [2] we extended these results and obtained some inequalities that implied $p_3 \leq \frac{5}{12}n(n - 1)$. We now extend these results further, proving two results.

THEOREM 1. $p_3 \leq \frac{2}{3}n(n - 1)$ if A is not a near pencil ($n - 1$ lines concurrent), and the result is best possible since equality occurs when $n = 6$.

THEOREM 2. $p_3 \leq \frac{7}{18}n(n - 1) + \frac{1}{3}$ if A is not a near pencil.

REMARK. If A is a near pencil, then $p_3 = 2n - 2$.

2. The proofs.

PROOF OF THEOREM 1. Let A be an arrangement of n lines in P_2 , not a pencil or near pencil. Let t_k be the number of points incident with precisely k lines. For a fixed line l of the arrangement, let $t_k(l)$ denote the number of points on l incident with precisely k lines, including l . Let us call a point incident with precisely two lines a simple point, and the other points multiple points. Let $P_1, \dots, P_m$ be the multiple points, in order, on the line l , with the understanding that $m = 0$ if there are no multiple points on l . We let $I_i = (P_i, P_{i+1})$ called the i th block be the set of simple points on l between P_i and P_{i+1} on l , with the understanding that

Received by the editors April 2, 1979.

AMS (MOS) subject classifications (1970). Primary 52A40; Secondary 05A20.

© 1980 American Mathematical Society
0002-9939/80/0000-0215/\$02.25

$(P_m, P_{m+1}) = (P_m, P_1)$. In the event that $m = 1$ we have the single block (P_1, P_1) . We must therefore allow for the possibility that the endpoints of $I_i = (P_i, P_{i+1})$ are the same point. If $m = 0$ there are no blocks. Each block I_i determines $|I_i| + 1$ line segments $\overline{P_i Q_i}, \overline{Q_1 Q_2}, \dots, \overline{Q_{k-1} Q_k}, \overline{Q_k P_{i+1}}$, where $I_i = \{Q_1, \dots, Q_k\}$, where $|S|$ denotes the cardinality of the set S . If $I_i = \emptyset$, then there is only the line segment $\overline{P_i P_{i+1}}$. Let $1 \leq j \leq k$. The line segment $\overline{Q_j Q_{j+1}}$ is the side of at most one triangle, for suppose $\overline{Q_j Q_{j+1}}$ is incident with two triangles. Then the removal of l would result in a digon (a two-sided polygon), and digons only occur in pencils, contradicting the assumption that A is not a near pencil. Thus, if $p_3(l)$ denotes the number of triangles having a side lying on l , then the contribution of I_i to $p_3(l)$ is at most $k + 3 = |I_i| + 3$. If in fact the contribution is less and if $I_i \neq \emptyset$, then we say that I_i is a deficient block. Let D^l denote the number of deficient blocks on l and E^l the number of empty blocks on l . If $I_i = \emptyset$, then the contribution to $p_3(l)$ is only $|I_i| + 2 = 2$. Hence, if $m \geq 1$,

$$\begin{aligned} p_3(l) &\leq \sum_i \{|I_i| + 3\} - D^l - E^l \\ &= t_2(l) + 3 \sum_{k \geq 3} t_k(l) - D^l - E^l. \end{aligned}$$

If $m = 0$,

$$p_3(l) \leq t_2(l) = t_2(l) + 3 \sum_{k \geq 3} t_k(l) - D^l - E^l,$$

and so, in any case,

$$p_3(l) \leq t_2(l) + 3 \sum_{k \geq 3} t_k(l) - D^l - E^l.$$

If we let E and D denote respectively the number of empty and deficient blocks on all lines, then taking into account that triangles get counted three times and points on k lines get counted k times, we obtain on summation over l ,

$$3p_3 \leq 2t_2 + 3 \sum_{k \geq 3} kt_k - E - D. \quad (1)$$

We need to consider so-called F -blocks. Let $|I_i| = 2$, $I_i = \{Q_1, Q_2\}$, and suppose that I_i is not deficient. Then we have the situation depicted in Figure 1, with $|I_i| + 3 = 5$ triangles numbered from one to five. We call such a block a type F -block and it forces the existence of a point P incident with at least four lines. Let us give the segments $\overline{Q_1 P}$ and $\overline{Q_2 P}$ weight one to indicate that they cannot occur in any other block, and give $\overline{P_i P}$ and $\overline{P_{i+1} P}$ weight $\frac{1}{2}$ to indicate that they can each occur in at most two blocks (or possibly $P_i = P_{i+1}$). The sum of these weights is $3 \sum_i F^l = 3F$, where F^l is the number of F -blocks on l , and F is the total number of F -blocks. The sum of these weights cannot exceed $\sum_{k \geq 4} 2kt_k$, and so we have

$$\frac{4}{3} \sum_{k \geq 4} kt_k \geq 2F. \quad (2)$$

Let us now look at a typical block I_i which is not an empty block, a deficient block or an F -block. $I_i = \{Q_1, \dots, Q_k\}$. See Figure 2. The triangles are marked with the symbol Δ . The points P and P' are each incident with three or more lines,

FIGURE 1

FIGURE 2

and we have given $\overline{P_i P}$ and $P_{i+1} P'$ weight $\frac{1}{2}$ and we have given $\overline{Q_1 P}$, $\overline{Q_2 P}$, $\overline{Q_{k-1} P'}$ and $\overline{Q_k P'}$ weight one, and block I_i has total weight 5 in the figure drawn. In fact, P and P' could lie on the same side of l and might even coincide. We claim that in all cases the total weight will be at least 4. In Figure 3 we see how 4 can occur. If P_i and P_{i+1} coincide it is to be understood that each occurrence of $\overline{P_i P}$ in Figure 2 or 3 gets weight $\frac{1}{2}$, so that the single edge gets, in fact, a weight of one.

FIGURE 3

The total weight from all of these blocks cannot exceed $\sum_{k \geq 3} 2kt_k$, and the number of these blocks on the line l is

$$\sum_{k \geq 3} t_k(l) - F^l - E^l - D^l.$$

Hence

$$\begin{aligned} \sum_{k \geq 3} 2kt_k &\geq 4 \sum_l \left\{ \sum_{k \geq 3} t_k(l) - F^l - E^l - D^l \right\} \\ &= 4 \left\{ \sum_{k \geq 3} kt_k - F - E - D \right\}, \end{aligned}$$

and we have

$$2E + 2D \geq \sum_{k \geq 3} kt_k - 2F,$$

and adding inequality (2), we obtain

$$2E + 2D \geq 3t_3 - \frac{1}{3} \sum_{k \geq 4} kt_k.$$

Combining with (1), we get

$$3p_3 \leq 2t_2 + \frac{15}{2}t_3 + \frac{19}{6} \sum_{k \geq 4} kt_k. \quad (3)$$

Let p_k be the number of k -gons determined by the region. Counting all of the edges, we see that $\sum kp_k = 2 \sum kt_k$, which we rewrite as

$$3p_3 \leq 4t_2 + 6t_3 + 2 \sum_{k \geq 4} kt_k - \sum_{k \geq 4} kp_k. \quad (4)$$

Adding $\frac{4}{5}$ of (3) to $\frac{1}{5}$ of (4) we obtain

$$\begin{aligned} 3p_3 &\leq \frac{12}{5}t_2 + \frac{36}{5}t_3 + \frac{44}{15} \sum_{k \geq 4} kt_k \\ &\leq \frac{12}{5} \left\{ t_2 + 3t_3 + \sum_{k \geq 4} \binom{k}{2} t_k \right\} = \frac{6}{5}n(n-1), \end{aligned}$$

and $p_3 \leq \frac{2}{5}n(n-1)$. To see that equality can occur for $n = 6$, see Figure 4, and Theorem 1 is proved.

PROOF OF THEOREM 2. Thomas O. Strommer [3] proved

$$p_3 \leq \frac{1}{3}n(n-1) + 4 - \frac{2}{3}t_2 + \sum_{k \geq 4} (k-4)p_k. \quad (5)$$

Adding this to (4) and multiplying the result by $\frac{3}{4}$ we obtain

$$3p_3 \leq \frac{1}{4}n(n-1) + 3 + \frac{5}{2}t_2 + \frac{9}{2}t_3 + \frac{3}{2} \sum_{k \geq 4} kt_k - 3 \sum_{k \geq 4} p_k. \quad (6)$$

FIGURE 4. $n = 6, p_3 = 12 = \frac{2}{5}n(n-1)$

If we now add $\frac{1}{3}$ of (6) to $\frac{2}{3}$ of (3) we obtain

$$\begin{aligned} 3p_3 &< \frac{1}{12}n(n-1) + 1 + \frac{13}{6}t_2 + \frac{39}{6}t_3 + \frac{47}{18}\sum_{k \geq 4} kt_k \\ &< \frac{1}{12}n(n-1) + 1 + \frac{13}{6}\left\{t_2 + 3t_3 + \sum_{k \geq 4} \binom{k}{2}t_k\right\} \\ &= \frac{14}{12}n(n-1) + 1, \end{aligned}$$

and Theorem 2 follows.

REMARK. Inequality (6) gives a rather curious upper bound on $\sum_{k \geq 3} p_k$, the total number of regions.

ACKNOWLEDGEMENT. We thank T. O. Strommer for helpful suggestions regarding this exposition.

REFERENCES

1. B. Grünbaum, *Arrangements and spreads*, CBMS Regional Conf. Ser. in Math., no. 10, Amer. Math. Soc., Providence, R. I., 1972.
2. G. Purdy, *Triangles in arrangements of lines*, Discrete Math. **25** (1979), 157–163.
3. T. O. Strommer, *Triangles in arrangements of lines*, J. Combinatorial Theory Ser. A **23** (1977), 314–320.

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843