AUTOMORPHISMS OF $GL_n(R)$

WILLIAM C. WATERHOUSE¹

ABSTRACT. Let R be a commutative ring in which 2 is invertible. Then all automorphisms of $GL_n(R)$ for $n > 3$ are of standard type.

Let R be a commutative ring in which 2 is invertible. This paper completes the proof that all automorphisms of the group $GL_n(R)$ for $n > 3$ are of standard type. This was previously known only under various additional hypotheses, the best result being that it held when R has no nontrivial idempotents [4]. Basically all that I do here is to establish in general the initial results needed to carry out McDonald's proof in [4], and I am grateful to him for raising this problem at the Pennsylvania State University conference on abstract automorphisms of algebraic groups. I have included a few remarks on the nature of standard automorphisms, since the group scheme properties involved have not been made explicit in the papers on this subject. These also simplify the proof that for our rings the group $E_n(R)$ is characteristic in $GL_n(R)$.

1. The nature of standard automorphisms. The automorphisms of $GL_n(R)$ called standard are the composites of three different types. First, there are the radial automorphisms: for certain homomorphisms $\chi: GL_n(R) \rightarrow R^*$ the map $A \mapsto \chi(A)A$ may be an automorphism of $GL_n(R)$. Secondly, every ring automorphism of R induces an automorphism of $GL_n(R)$. Finally, there are automorphisms of $GL_n(R)$ which are induced by automorphisms of the whole algebraic group scheme GL_n over R . These are known for all reductive group schemes [2, p. 328], and in this sense $\text{Aut}(GL_n)$ is simply the semidirect product of PGL_n and the constant group scheme $\mathbb{Z}/2\mathbb{Z}$ of order 2. But it is worth pausing to see what this means concretely.

It is true quite formally that the constant group scheme of order 2 acts on GL_n by transpose inverse. When in addition $\text{Spec } R$ is connected, $(\mathbb{Z}/2\mathbb{Z})(R)$ has just two elements, and the action on $GL_n(R)$ is the obvious one. But in general $(\mathbb{Z}/2\mathbb{Z})(R)$ has many more elements, one for each clopen subset of $\text{Spec } R$. It is easy to compute that their action on GL_n , and in particular on $GL_n(R)$, is trivial over one piece of $\text{Spec } R$ and transpose inverse over the other.

The standard automorphisms coming from PGL_n , finally, reflect the action of GL_n on itself by inner automorphisms. The center G_m acts trivially, so since the action is algebraic it factors through the group scheme quotient PGL_n of GL_n by G_m . That is, though the map $GL_n(R) \rightarrow PGL_n(R)$ need not be surjective, the formal

Received by the editors August 6, 1979.

1980 *Mathematics Subject Classification*. Primary 20G35.

¹This work was supported in part by the National Science Foundation.

© 1980 American Mathematical Society
0002-9939/80/0000-0300/\$02.25

properties make the latter group operate as automorphisms of $GL_n(R)$. This is the *only* reason why in previous papers one finds "standard automorphisms" performed by conjugations from some larger $GL_n(S)$. We will see below that only the action of $PGL_n(R)$ is needed, and the somewhat arbitrary choice of the extension S is irrelevant.

This analysis also shows *a priori* how big such an S must be if one wants it. The cohomology exact sequence (see e.g. [6, Chapter 18]) gives

$$1 \rightarrow G_n(R) \rightarrow GL_n(R) \rightarrow PGL_n(R) \rightarrow \text{Pic}(R) \rightarrow H^1(/R, GL_n).$$

Interpreting GL_n as automorphisms of the free module, one can easily see that the last map sends an invertible module P (twisted form of R) to the twisted form P^n of R^n . The kernel then is those P with P^n free. Mapping to the sequence for an extension S , we see that all of $PGL_n(R)$ comes from $GL_n(S)$ whenever $P \otimes_R S$ is free for all such P . Needless to say, this is the condition which did indeed come up in [4]. But specifically what one needs is the following automorphisms not coming from $GL_n(R)$.

PROPOSITION. *Suppose $V = R^n = \bigoplus Rb_i$ is expressed as $P_1 \oplus \cdots \oplus P_n$ with the P_i all isomorphic. Let $\sigma'_i: V \rightarrow V$ be the reflection that is -1 on P_i and 1 on the other summands. Let σ_i be defined similarly from the free decomposition $V = \bigoplus Rb_i$. Then some element of $PGL_n(R)$ defines an automorphism of $GL_n(R)$ that takes the σ'_i to σ_i .*

PROOF. Fix an identification of each P_i with P_1 , and define a group of automorphisms $\{\pi'\}$ of V (isomorphic to S_n) by permuting the n -tuples in $P_1 \oplus \cdots \oplus P_1$. Choose some faithfully flat $R \rightarrow S$ with $P_1 \otimes S$ free. Choose a basis element c_1 of $P_1 \otimes S$. The $\pi' \otimes \text{id}$ send this to a well-determined set of basis elements c_i of $P_i \otimes S$, and are simply the permutations of this basis of $V \otimes S$.

Let φ in $GL_n(S)$ be the map sending c_i to b_i . Then $\varphi^{-1}(\sigma_i \otimes \text{id})\varphi = \sigma'_i \otimes \text{id}$ and $\varphi^{-1}(\pi \otimes \text{id})\varphi = \pi' \otimes \text{id}$, where $\{\pi\}$ are the permutations of the b_i . Since $\sigma'_i \otimes \text{id}$ comes from $GL_n(R)$, it has the same image $\sigma'_i \otimes \text{id} \otimes \text{id}$ under both the maps $d^0, d^1: GL_n(S) \rightarrow GL_n(S \otimes S)$. This says that

$$(d^1\varphi)^{-1}(\sigma'_i \otimes \text{id} \otimes \text{id})(d^1\varphi) = (d^0\varphi)^{-1}(\sigma_i \otimes \text{id} \otimes \text{id})(d^0\varphi),$$

which is equivalent to

$$(d^0\varphi)(d^1\varphi)^{-1}(\sigma_i \otimes \text{id} \otimes \text{id}) = (\sigma_i \otimes \text{id} \otimes \text{id})(d^0\varphi)(d^1\varphi)^{-1}.$$

Thus $(d^0\varphi)(d^1\varphi)^{-1}$ commutes with the extensions of the σ_i , and hence it is diagonal in $GL_n(S \otimes S)$. It similarly commutes with the extensions of the π , and hence it is scalar. Thus $d^0(\bar{\varphi})$ and $d^1(\bar{\varphi})$ in $PGL_n(S \otimes S)$ are equal, which by faithfully flat descent means that $\bar{\varphi}$ in $PGL_n(S)$ actually comes from an element in $PGL_n(R)$. By construction this element has the desired effect. $\square$

2. Automorphisms of $GL_n(R)$. We will need the following simple lemma. It undoubtedly must be known, but I cannot find a reference for it, so I sketch a proof.

LEMMA. *Let R be a commutative ring. Let M be a finitely generated projective R -module whose local rank at some prime is greater than one. Then $\text{Aut}_R(M)$ is noncommutative.*

PROOF. Embed M as a summand of some R^m . Only finitely many elements of R are needed to write down the projections onto M and onto a complement of M . Let R' be the subring they generate. Then $M = M' \otimes_{R'} R$ for a projective M' over R' , and $\text{Aut}_{R'}(M')$ injects into $\text{Aut}_R(M)$. Replacing R and M by R' and M' , we may assume R is noetherian. The rank is locally constant [1, p. 138], so we can decompose R as $\prod R_i$ with $M = \prod M_i$ having M_i of rank i . Keeping the other factors fixed, we can embed $\text{Aut}_{R_i}(M_i)$ in $\text{Aut}_R(M)$. Replacing R by R_i , we may assume M has constant rank.

Let S be the localization of R in which all elements not divisors of zero are inverted. Then S is semilocal [1, p. 151], and $M \otimes S$ is free of rank greater than one. We have $\text{End}_R(M) \otimes S \simeq \text{End}_S(M \otimes S)$, and $\text{End}_R(M)$ injects into $\text{End}_S(M \otimes S)$. Choose a basis of $M \otimes S$, and let e_{12} and e_{21} be matrix units for that basis. There is some f in R , not a divisor of zero, with fe_{12} in $\text{End}_R(M)$, and similarly some ge_{21} is in $\text{End}_R(M)$. Now $1 + fe_{12}$ in $\text{End}_R(M)$ actually lies in $\text{Aut}_R(M)$, since its inverse $1 - fe_{12}$ in $\text{End}_S(M \otimes S)$ also lies in $\text{End}_R(M)$. Similarly $1 + ge_{21}$ is in $\text{Aut}_R(M)$. As f and g are not zero-divisors, these elements do not commute. $\square$

From now on we assume 2 is invertible in R and n is ≥ 3 . Denote by $\{b_i\}$ the standard basis of R^n , and let $\sigma_i: R^n \rightarrow R^n$ send b_i to $-b_i$ and the other b_j to themselves. Let Λ be an arbitrary automorphism of $GL_n(R)$.

LEMMA. *For each subset α of $\{1, \dots, n\}$, let*

$$P_\alpha = \{v | \Lambda(\sigma_i)v = -v \text{ for } i \in \alpha \text{ and } \Lambda(\sigma_i)v = v \text{ for } i \notin \alpha\}.$$

Then $R^n = \bigoplus P_\alpha$, and $P_\alpha = 0$ unless $|\alpha|$ equals 1 or $n - 1$. All P_α for α of the same size are isomorphic.

PROOF. Since 2 is invertible, trivially $R^n = \bigoplus P_\alpha$. In particular, all P_α are projective. Let $\{\pi\}$ now be the permutations of the b_i . We have $\pi\sigma_i\pi^{-1} = \sigma_{\pi(i)}$, and hence $\Lambda(\pi)\Lambda(\sigma_i)\Lambda(\pi)^{-1} = \Lambda(\sigma_{\pi(i)})$. Simple computation shows then $\Lambda(\pi)P_\alpha \subseteq P_{\pi(\alpha)}$, which implies $\Lambda(\pi)P_\alpha = P_{\pi(\alpha)}$ since we have a group acting. The π act transitively on the α of a fixed size, so all such P_α are isomorphic.

Consider an α with $1 < |\alpha| < n - 1$. If P_α is nonzero, we can find a prime where its localization is nonzero. The π -action gives us then $\binom{n}{|\alpha|}$ different free summands of the localized module, and of course this is too many. Thus $P_\alpha = 0$. For $|\alpha| = 1$ or $|\alpha| = n - 1$ the same argument shows that at any prime where P_α is nonzero, all summands for different size subsets must vanish.

If now P_ϕ or $P_{\{1, \dots, n\}}$ is nonzero, then at primes in their support we can have no terms but these two occurring. Hence the local rank of one of them must be at least two. By the previous lemma it has noncommutative automorphism group. That group embeds as a subgroup of $GL_n(R)$ commuting with all $\Lambda(\sigma_i)$. But this is

impossible, because the centralizer of the σ_i is the commutative group of diagonal matrices. $\square$

It is worth noting that P_α may be nonzero both for $|\alpha| = 1$ and $|\alpha| = n - 1$ at once. For example, let k be a perfect field, $\text{char}(k) = 5$, and let $R = k \times k$. On $GL_4(R) = GL_4(k) \times GL_4(k)$, the map $\Lambda(A \times B) = A \times (\det B)B$ is an automorphism. Here $\Lambda(\sigma_i)$ has first component that of σ_i and second component that of $-\sigma_i$, and the $\Lambda(\sigma_i)$ split R^4 into eight eigenspaces P_α . This disproves McDonald's conjecture in [4] that all automorphisms are "stable" in the sense defined there. But the failure is not serious, because it arises only from a complexity in standard (radial) automorphisms.

PROPOSITION. *Changing Λ by an automorphism from $PGL_n(R)$, one can make all $\Lambda(\sigma_i)$ equal $e\sigma_i$ for some e in R with $e^2 = 1$.*

PROOF. The previous lemma shows that R^n splits into n isomorphic summands P_α with $|\alpha| = 1$ and n more isomorphic summands (with complementary support) P_α with $|\alpha| = n - 1$. Let e be the element of R which is 1 on the support of $P_{\{1\}}$ and -1 on its complement. Grouping together complementary summands, we express R^n as $P_1 \oplus \cdots \oplus P_n$ with all P_i isomorphic, and we have $\Lambda(\sigma_i)v$ equal to $(1 - 2e)v$ for v in P_i and equal to ev for v in the other P_j . Changing Λ by the automorphism constructed in §1, we see the composite sends σ_i precisely to $e\sigma_i$. $\square$

COROLLARY. *Let $\psi_{ij} = \sigma_i\sigma_j$. Then after the change of Λ one has $\Lambda(\psi_{ij}) = \psi_{ij}$.*

PROOF. $\Lambda(\psi_{ij}) = \Lambda(\sigma_i)\Lambda(\sigma_j) = e\sigma_i e\sigma_j = e^2\psi_{ij} = \psi_{ij}$. $\square$

THEOREM. *Let R be a commutative ring in which 2 is invertible. Then every automorphism of $GL_n(R)$ for $n > 3$ is standard.*

PROOF. The arguments in [4] after p. 163, including those cited from [5], require nothing but the corollary above. Furthermore, the extension from R to S is not needed once the corollary is established, and thus the automorphisms are indeed standard in the sense explained in §1. $\square$

COROLLARY. *For such R and n , let I be an ideal of R mapped to itself by all automorphisms of R . Then the group of elementary matrices $E_n(I)$ is characteristic in $GL_n(R)$. In particular, $E_n(R)$ is characteristic.*

PROOF. Clearly automorphisms of R will send $E_n(I)$ to itself, and radial automorphisms are trivial on it because it is in the commutator subgroup. Let Λ now come from a group scheme automorphism. Suslin proved that $E_n(I)$ is normal (see [4, Appendix]), and so $\Lambda(E_n(I))$ is normal. It follows [3], [7] that for some J ,

$$E_n(J) \subseteq \Lambda(E_n(I)) \subseteq C_n(J),$$

where $C_n(J)$ is the subgroup that maps to the center of $GL_n(R/J)$. But group scheme automorphisms induce compatible automorphisms of $GL_n(R/J)$, and hence they send $C_n(J)$ to itself. Applying this to Λ^{-1} , we find $E_n(I) \subseteq C_n(J)$, which implies $I \subseteq J$. Thus $E_n(I) \subseteq E_n(J)$ is contained in $\Lambda(E_n(I))$ for all such Λ . The reverse containment follows from the statement for Λ^{-1} . $\square$

Technically, I believe the proof of the theorem itself requires the fact that standard automorphisms preserve $E_n(R)$. Specifically this seems to be needed in the first paragraph on p. 168 of [4]; the matrix P there is not in $GL_n(R)$, and thus Suslin's result is not directly applicable. But there is no logical circle involved, since in the corollary we used the theorem only to restrict our considerations to standard automorphisms.

REFERENCES

1. N. Bourbaki, *Algèbre commutative*, Hermann, Paris, 1961.
2. M. Demazure, A. Grothendieck, et al., *SGA 3: Schémas en groupes*. III, Lecture Notes in Math., vol. 153, Springer-Verlag, New York, 1970.
3. I. Z. Golubčik, *On the full linear group over an associative ring*, *Uspehi Mat. Nauk* **28** (171) (1973), 179–180. (Russian).
4. B. R. McDonald, *Automorphisms of $GL_n(R)$* , *Trans. Amer. Math. Soc.* **246** (1978), 155–171.
5. J. Pomfret and B. R. McDonald, *Automorphisms of $GL_n(R)$, R a local ring*, *Trans. Amer. Math. Soc.* **173** (1972), 379–388.
6. W. C. Waterhouse, *Introduction to affine group schemes*, Graduate Texts in Math., no. 66, Springer-Verlag, New York, 1979.
7. J. S. Wilson, *The normal and subnormal structure of general linear groups*, *Proc. Cambridge Philos. Soc.* **71** (1972), 163–177.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PENNSYLVANIA 16802