IDEMPOTENT MULTIPLIERS ON SPACES OF CONTINUOUS FUNCTIONS WITH p -SUMMABLE FOURIER TRANSFORMS

LYNETTE M. BLOOM AND WALTER R. BLOOM

ABSTRACT. Let G denote a compact abelian group, and A^p the space of functions continuous on G and having p -summable Fourier transforms. The idempotent multipliers from A^p to A^q are characterised for $p, q \in [1, 2]$.

Throughout G will denote a compact Hausdorff abelian group with character group Γ . Given $p \in [1, 2]$ we shall write A^p for the Banach space of functions continuous on G with p -summable Fourier transforms, and normed by $\|f\| = \|f\|_\infty + \|\hat{f}\|_p$. Note that we can identify A^2 with C and A^1 with A , the spaces of functions on G that are continuous and have absolutely convergent Fourier series respectively. An application of the Hahn-Banach theorem shows that the dual $(A^p)'$ of A^p can be identified with $M + Fl^{p'}$, where M is the space of Radon measures on G and $Fl^{p'}$ is the space of pseudomeasures on G with Fourier transforms in $l^{p'}$. Here p' is the exponent conjugate to p , that is, $p' = p/(p - 1)$ with the usual convention if $p = 1$. The duality is expressed by

$$h(f) = \mu * f(0) + \sigma * f(0), \quad f \in A^p,$$

where $h \in (A^p)'$, $\mu \in M$ and $\sigma \in Fl^{p'}$.

We shall write (A^p, A^q) for the set of pseudomeasures that are multipliers from A^p to A^q ; that is, $\sigma \in (A^p, A^q)$ if and only if for every $f \in A^p$ there exists $g \in A^q$ with $\sigma * f = g$. Using the above characterisation of the dual of A^p the following result can be proved.

LEMMA 1. *Given $p \in [1, 2]$ and $q \in [p, 2]$,*

$$(A^p, A^q) = (A^p, A^p) = (A^p, C) = M + Fl^{p'}.$$

The A^p spaces and their multiplier spaces have been considered previously; see [1], [2], [5] and [6]. Here we are interested in characterising the set of multipliers $\phi \in (A^p, A^q)$ that are idempotent; that is, satisfy $\phi * \phi = \phi$. We shall write $(A^p, A^q)_I$ for the set of all idempotent elements of (A^p, A^q) . In addition, we shall denote by J the set of all idempotent measures on G , by ξ_Ξ the characteristic function of the set Ξ , and by $|\Xi|$ the cardinality of Ξ if Ξ is finite.

By considering Fourier transforms we see that to each $\phi \in (A^p, A^q)_I$ there corresponds unique $\Xi \subseteq \Gamma$ for which $\hat{\phi} = \xi_\Xi$. In the case $p = 1$, $q \in [1, 2]$, it is easy to see (using Lemma 1) that

Received by the editors October 20, 1978.

AMS (MOS) subject classifications (1970). Primary 43A22, 43A15.

© 1980 American Mathematical Society
 0002-9939/80/0000-0319/\$02.00

$$(A, A^q)_I = (A, A)_I = \{\phi: \hat{\phi} = \xi_{\Xi} \text{ for some } \Xi \subseteq \Gamma\}.$$

For other values of p, q , $p < q$, the idempotent multipliers are precisely the idempotent measures.

THEOREM 1. For $p \in (1, 2]$, $q \in [p, 2]$, $(A^p, A^q)_I = (A^p, A^p)_I = J$.

PROOF. The first equality follows directly from Lemma 1. As for the second, consider $\phi \in (A^p, A^p)_I$. By Lemma 1 we can write $\phi = \mu - \sigma$, where $\mu \in M$ and $\hat{\sigma} \in l^{p'}$. Choose a positive integer n satisfying $2n < p' < 2(n+1)$. Then $\hat{\sigma}^{n+1} \in l^2$ and hence $\sigma^{*r} \in L^2$ for all $r > n+1$, where σ^{*r} denotes the r -fold convolution of σ with itself. Now

$$\mu^{*s} = (\phi + \sigma)^{*s} = \sum_{r=0}^s \binom{s}{r} \phi^{*s-r} * \sigma^{*r} = \sum_{r=0}^{s-1} \binom{s}{r} \phi * \sigma^{*r} + \sigma^{*s}$$

(recall that $\phi^{*k} = \phi$ for all $k > 1$) and consideration of this expression for $s = n+1, n+2, \dots, 2n+1$ gives

$$\begin{bmatrix} 1 & \binom{n+1}{1} & \dots & \binom{n+1}{n} \\ 1 & \binom{n+2}{1} & \dots & \binom{n+2}{n} \\ \vdots & \vdots & & \vdots \\ 1 & \binom{2n+1}{1} & \dots & \binom{2n+1}{n} \end{bmatrix} \begin{bmatrix} \phi \\ \phi * \sigma \\ \vdots \\ \phi * \sigma^{*n} \end{bmatrix} = \begin{bmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_{n+1} \end{bmatrix}$$

where $\mu_i \in M$, $i = 1, 2, \dots, n+1$. The coefficient matrix is nonsingular; indeed using the identity $\binom{t+1}{s} - \binom{t}{s} = \binom{t}{s-1}$, $t > s$, it can be reduced to the upper triangular matrix (α_{ij}) , where $\alpha_{ij} = \binom{n+1}{j-i}$ for $i < j$. It follows that $\phi \in M$ and hence $\phi \in J$, thus showing that $(A^p, A^p)_I \subseteq J$. The reverse inclusion is clear, completing the proof of the theorem. $\square$

For $p \in (1, 2]$, $q \in [1, p)$ the only idempotent multipliers ϕ from A^p to A^q are those given by $\hat{\phi} = \xi_{\Xi}$, where Ξ is finite. To prove this we require some preliminary results.

Let $\mathcal{K}$ denote the family of all cosets in Γ and $\mathcal{R}$ the Boolean ring generated by $\mathcal{K}$. Every $\Xi \in \mathcal{R}$ can be written as a finite union of sets of the form $\Pi = \bigcap_{i=1}^n \Lambda_i \cap \bigcap_{i=n+1}^r \Lambda_i^c$, where $\Lambda_i \in \mathcal{K}$ and Λ_i^c denotes the complement of Λ_i . If Ξ is infinite then so must be at least one such set Π .

LEMMA 2. Given an infinite set Π of the above type there exists a finite set $\mathcal{T}$ with the property that $\Pi \cup \mathcal{T}$ contains an infinite coset.

PROOF. First we show that a finite intersection of cosets is either empty or itself a coset. Indeed consider $\Lambda_1, \Lambda_2 \in \mathcal{K}$ such that $\Lambda_1 \cap \Lambda_2$ is nonempty. Then $\Lambda_1 = \eta_1 + \Omega_1$, $\Lambda_2 = \eta_2 + \Omega_2$, where Ω_1, Ω_2 are subgroups of Γ , and there exist $\chi_1 \in \Omega_1$, $\chi_2 \in \Omega_2$ such that $\eta_1 + \chi_1 = \eta_2 + \chi_2$. Hence

$$\Lambda_1 \cap \Lambda_2 = (\eta_2 + \chi_2 + \Omega_1) \cap (\eta_2 + \chi_2 + \Omega_2) = \eta_2 + \chi_2 + \Omega_1 \cap \Omega_2 \in \mathcal{K}.$$

Thus a nonempty intersection of two cosets is itself a coset and the result extends to arbitrary finite intersections by induction.

Applying this to (the nonempty set) Π we see that

$$\Pi = \bigcap_{i=1}^n \Lambda_i^c \cap \Lambda, \quad \Lambda_i, \Lambda \in \mathcal{K}.$$

By relabelling if necessary we can assume that $\Pi \cup \mathcal{T}$ contains the infinite set $\bigcap_{i=1}^n \Lambda_i^c \cap \Lambda$, where $\mathcal{T}$ is finite and each Λ_i is infinite. Now $\Lambda_i = \eta_i + \Omega_i$, $\Lambda = \eta + \Omega$ for some subgroups Ω_i, Ω of Γ . We consider the following cases:

(1) $\bigcap_{i=1}^n \Omega_i \cap \Omega$ is infinite. Then $\Pi \cup \mathcal{T}$ contains $\bigcap_{i=1}^n \Lambda_i^c \cap \Lambda$ which, for suitable cosets Λ'_i of Ω_i , is nonempty; just use the fact that each Λ_i^c is a union of cosets of Ω_i . By the argument used in the first part of the proof of the lemma, $\bigcap_{i=1}^n \Lambda'_i \cap \Lambda$ is an infinite coset in Γ .

(2) $\bigcap_{i=1}^n \Omega_i \cap \Omega$ is finite, $\bigcap_{i \in I} \Omega_i \cap \Omega$ is infinite for some $I \subseteq \{1, 2, \dots, n\}$ with $|I| = n - 1$. Then there are cosets Λ'_i of Ω_i such that $\Pi \cup \mathcal{T}$ contains the nonempty set $\bigcap_{i \in I} \Lambda'_i \cap \Lambda \cap \Lambda_j^c$, where $\{j\} = \{1, 2, \dots, n\} \setminus I$, and $\mathcal{T}' = \bigcap_{i \in I} \Lambda'_i \cap \Lambda \cap \Lambda_j$ is (empty or) finite. Thus $\Pi \cup \mathcal{T} \cup \mathcal{T}' \supseteq \bigcap_{i \in I} \Lambda'_i \cap \Lambda$, an infinite coset.

(3) $\bigcap_{i \in I} \Omega_i \cap \Omega$ is finite for all $I \subseteq \{1, 2, \dots, n\}$ with $|I| = n - 1$. In this case we return to (2) above and repeat the process.

After $n - 1$ steps we have that $\Omega_i \cap \Omega$ is finite for each $i \in \{1, 2, \dots, n\}$ and

$$\Pi \cup \mathcal{T} \cup \bigcup_{i=1}^n (\Lambda_i \cap \Lambda) \supseteq \left(\bigcap_{i=1}^n \Lambda_i^c \cap \Lambda \right) \cup \left(\bigcup_{i=1}^n \Lambda_i \cap \Lambda \right) = \Lambda$$

is the required infinite coset. $\square$

THEOREM 2. Let $\Xi \in \mathcal{R}$ be infinite and $p \in (1, 2]$. Then there exists a continuous function f with $\text{supp } \hat{f} \subseteq \Xi$ and $\hat{f} \in l^p \setminus l^q$ for all $q \in [1, p)$.

PROOF. By Lemma 2 we have the existence of a finite set $\mathcal{T}$ such that $\Xi \cup \mathcal{T}$ contains an infinite coset Λ . We shall construct f as in the statement of the theorem with $\text{supp } \hat{f} \subseteq \Lambda$, and then the result will follow. There are two cases:

(a) The subgroup $\Omega = -\eta + \Lambda$ has an element χ of infinite order. Now construct Rudin-Shapiro polynomials f_n in the usual way ([7, (37.19)]) so that $\hat{f}_n$ takes values in $\{-1, 0, 1\}$,

$$\text{supp } \hat{f}_n = \{0, \chi, \dots, (2^n - 1)\chi\} \quad \text{and} \quad \|f_n\|_\infty \leq 2^{(n+1)/2}.$$

Choose $\chi_n \in \Omega$ so that the spectra of the functions $\chi_n f_n$ are pairwise disjoint and put $f = \sum_{n=0}^\infty (n+1)^{-2} 2^{-n/p} \eta \chi_n f_n$. Then

$$\sum_{n=0}^\infty (n+1)^{-2} 2^{-n/p} \|\eta \chi_n f_n\|_\infty < \infty,$$

so that $f \in C$. Clearly $\text{supp } \hat{f} \subseteq \Lambda$. That $f \in A^p$ follows from the estimate

$$\|\hat{f}\|_p \leq \sum_{n=0}^\infty (n+1)^{-2} 2^{-n/p} 2^{n/p} < \infty,$$

whereas $\|\hat{f}\|_q^q = \sum_{n=0}^\infty (n+1)^{-2q} 2^{-nq/p} 2^n$ shows that $f \notin A^q$ for any $q < p$.

(b) $-\eta + \Lambda = \Omega$ is a torsion subgroup of Γ , in which case there is a strictly increasing sequence (Ω_n) of finite subgroups of Ω . For each n construct f_n (as in [4]) with the properties $|\hat{f}_n| = 1$ on Ω_n , $\text{supp } \hat{f}_n = \Omega_n$ and $\|f_n\|_\infty = |\Omega_n|^{1/2}$. As in (a) choose $\chi_n \in \Omega$ so that the spectra of the functions $\chi_n f_n$ are pairwise disjoint and put $f = \sum_{n=0}^\infty (n+1)^{-2} 2^{-n/p} \eta \chi_n f_n$. $\square$

THEOREM 3. *Let $p \in (1, 2]$, $q \in [1, p)$ be given. Then $\phi \in (A^p, A^q)_I$ if and only if $\hat{\phi} = \xi_\Xi$ for some finite set $\Xi \subseteq \Gamma$.*

PROOF. Sufficiency is obvious. As for the other direction, consider $\phi \in (A^p, A^q)_I$. Using $(A^p, A^q)_I \subseteq (A^p, A^p)_I$ we see from Theorem 1 that $\phi \in J$ and hence, by [3, Theorem 3], $\hat{\phi} = \xi_\Xi$ for some set Ξ belonging to the coset ring $\mathcal{R}$ of Γ . Suppose now that Ξ is infinite. By Theorem 2 there exists $f \in C$ with $\text{supp } \hat{f} \subseteq \Xi$ and $\hat{f} \in l^p \setminus l^q$. This contradicts the assumption that $\phi \in (A^p, A^q)_I$. $\square$

COROLLARY. *Suppose G is infinite and let $p \in (1, 2]$, $q \in [1, p)$ be given. Then $(A^p, A^q)_I \subsetneq (A^p, A^p)_I$.*

PROOF. That this inclusion is strict follows from Theorem 3 together with the fact that δ_0 (the Dirac measure at 0) belongs to $(A^p, A^p)_I$. $\square$

ACKNOWLEDGEMENT. We should like to thank Dr. John F. Price for suggesting some of the problems considered in this paper.

REFERENCES

1. Lynette M. Bloom, *The Fourier multiplier problem for spaces of continuous functions with p -summable transforms*, J. Austral. Math. Soc. **17** (1974), 319–331.
2. Lynette M. Butler, *Certain nonalgebras in harmonic analysis*, Bull. Austral. Math. Soc. **4** (1971), 247–254.
3. Paul J. Cohen, *On a conjecture of Littlewood and idempotent measures*, Amer. J. Math. **82** (1960), 191–212.
4. Colin C. Graham, *The Sidon constant of a finite abelian group*, Proc. Amer. Math. Soc. **68** (1978), 83–84.
5. Michael B. Gregory, *p -Helson sets*, $1 < p < 2$, Israel J. Math. **12** (1972), 356–368.
6. Edwin Hewitt and Gunter Ritter, *Über die Integrierbarkeit von Fourier-Transformierten auf Gruppen. Teil 1. Stetige Funktionen mit kompaktem Träger und eine Bemerkung über hyperbolische Differentialoperatoren*, Math. Ann. **224** (1976), 77–96.
7. Edwin Hewitt and Kenneth A. Ross, *Abstract harmonic analysis*. Vol. II, Die Grundlehren der Math. Wissenschaften, Band 152, Springer-Verlag, Berlin and New York, 1970.

SCHOOL OF MATHEMATICAL AND PHYSICAL SCIENCES, MURDOCH UNIVERSITY, MURDOCH, WESTERN AUSTRALIA, 6150, AUSTRALIA