DESINGULARIZING MAPS OF CORANK ONE

S. J. BLANK AND C. CURLEY

ABSTRACT. In 1960, A. Haefliger presented necessary and sufficient conditions for factoring a smooth map from a surface into the plane through an immersion into R^3 . Here, necessary and sufficient conditions are given for factoring a map between manifolds of dimension n , $n > 2$, through an immersion into a line bundle over the range. In addition, conditions are given for factoring such a map through a submersion from a line bundle over the domain.

By a desingularization of a smooth map $f: M \rightarrow V$ between manifolds we mean one of the following.

- (1) An immersion $F: M \rightarrow E$ for some line bundle $\pi: E \rightarrow V$ such that $f = \pi \circ F$ or
- (2) a submersion $F': E' \rightarrow V$ for some line bundle $\pi': E' \rightarrow M$ with zero section s such that $f = F' \circ s$.

We say f desingularizes in the first (resp. second) sense with respect to E (resp. E'). If f has a desingularization in the first sense, then the pull-back of the normal bundle to the image of F provides a line bundle with respect to which f has a desingularization in the second sense. The converse is not true. The map for Figure 1 desingularizes in the second sense but not the first. In this paper we present necessary and sufficient conditions for desingularizing a generic map with respect to a given bundle. In [1] A. Haefliger provides necessary and sufficient conditions for factoring a generic map from a closed surface into the plane through an immersion into R^3 . Let $f: M \rightarrow V$ be a smooth map where M is a closed smooth n -manifold and V is a smooth n -manifold and let $\pi: E \rightarrow V$ be a line bundle. If there is an immersion $F: M \rightarrow E$ with $f = \pi \circ F$ we say f factors through an immersion into E . If f does factor then $\text{rank } df > n - 1$ everywhere, or equivalently, the corank of f does not exceed one. The singularities of a generic f , by which we mean 2-generic and finite-to-one [2], with $\text{rank } df > n - 1$ everywhere, are stratified by the manifolds

$$S_1 = \{x \in M \mid \text{rank } df_x = n - 1\}, \quad S_{1,1} = \{x \in S_1 \mid \text{rank } d(f|_{S_1})_x = n - 2\}.$$

The map f being 2-generic ensures that S_1 and $S_{1,1}$ are submanifolds of M .

If we regard the Klein bottle as $\{(\phi, \theta) \mid \phi \in S^1, \theta \in [0, 2\pi]\} \text{ mod the relation } (\phi, 0) = (-\phi, 2\pi)$ then $f(\phi, \theta) = (\cos \phi + 2, \theta)$ is a map from the Klein bottle to the plane (in polar coordinates). We will show that this map desingularizes in the second sense but not in the first sense. Figure 1 represents the image of this map.

Received by the editors June 21, 1979 and, in revised form, September 29, 1979.

AMS (MOS) subject classifications (1970). Primary 57D35, 57D40; Secondary 57D20, 57D45.

Key words and phrases. Immersion, submersion, line bundle, 2-generic, corank, Stiefel-Whitney class.

© 1980 American Mathematical Society
0002-9939/80/0000-0572/\$02.00

FIGURE 1: Image of $f: (\text{Klein bottle}) \rightarrow \mathbb{R}^2$

For $n = 2$ a point in $S_{1,1}$ is called a cusp. In this setting Haefliger's theorem is: *A generic map $f: M^2 \rightarrow \mathbb{R}^2$ factors through an immersion into $\mathbb{R}^3$ if and only if each connected component of S_1 with an even (resp. odd) number of cusps has an orientable (resp. nonorientable) neighborhood.*

The following example of a map $f: S^2 \rightarrow \mathbb{R}^2$ which cannot be desingularized is due to A. Haefliger:

FIGURE 2

The set S_1 consists of two parallel curves C' and C'' each containing exactly one cusp; the image of the annulus bounded by two parallel curves between C' and C'' is a ribbon forming a figure eight bounded by the dashed curves; the two thin curves represent the images of the boundaries of polar caps.

For $n > 2$ there is the following theorem due to Y. Saito [3]: *A generic map $f: M^n \rightarrow \mathbb{R}^n$ factors through an immersion into $\mathbb{R}^{n+1}$ if (i) M is orientable, (ii) $\text{rank } df > n - 1$ everywhere and (iii) $S_{1,1} = \emptyset$.*

Let $N \rightarrow S_1$ be a tubular neighborhood of S_1 and let $K \rightarrow S_1$ be the line bundle

given by the kernel of df if $\text{rank } df \geq n - 1$ everywhere. We denote the first Stiefel-Whitney classes of these bundles by $w(N)$ and $w(K)$ respectively. Similarly we denote the class of the bundle $E \rightarrow V$ by $w(E)$. Let $[S_{1,1}]$ be the class in $H^1(S_1; \mathbb{Z}_2)$ dual to the homology class represented by $S_{1,1}$ and let $i: S_1 \rightarrow M$ be the inclusion map.

THEOREM 1. *A generic map $f: M \rightarrow V$ factors through an immersion into E if and only if the following conditions hold.*

- (i) $\text{rank } df \geq n - 1$ everywhere and
- (ii) $[S_{1,1}] + w(N) + i^*f^*w(E) = 0$.

REMARK. If M is orientable and E is trivial, then condition (ii) is equivalent to the statement that $S_{1,1}$ bounds in S_1 .

PROOF OF THEOREM 1. Even if f does not factor, we have $[S_{1,1}] + w(N) + w(K) = 0$.

We present an elementary proof of this fact. Let $\alpha: K \rightarrow N$ be the restriction to K of the projection map from the tangent space onto the normal bundle and let $\sigma: S_1 \rightarrow K$ be a section transverse to the zero section of K . Since f is 2-generic the section $\alpha \circ \sigma$ is transverse to the zero section of N . The homology class dual to $w(K)$ is represented by the intersection of α and the zero section. Similarly we can represent the dual of $w(N)$ by the intersection of $\alpha \circ \sigma$ and the zero section in N . Since $\alpha \circ \sigma(x)$ is in the zero section if and only if either $x \in S_{1,1}$ or $\sigma(x)$ is in the zero section, these homology classes are equal. Hence $[S_{1,1}] + w(K) = w(N)$.

If f factors through an immersion F , then dF defines a bundle map from K to E over $f \circ i$, which is an isomorphism on the fibres. Hence $i^*f^*w(E) = w(K)$.

Conversely, if $i^*f^*w(E) = w(K)$ there is a bundle map $L: K \rightarrow E$ over $f \circ i$ which is an isomorphism on the fibres. We will use this map in factoring f locally. For each $p \in M$ choose a closed disk B_p containing p such that the bundles K and E are trivial over $B_p \cap S_1$ and $f(B_p)$ respectively. Furthermore, since f is finite-to-one we can choose these disks so that $f(p) \notin f(\partial B_p)$. From this first condition we obtain, by integrating a vector field extending K , an immersion $F_p: B_p \rightarrow E$ for each p . Furthermore we require these immersions to satisfy $f = \pi \circ F_p$ and $d(F_p)_x(k) = \lambda_{p,x}L(k)$ where $x \in S_1 \cap B_p$, $k \in K$ and $\lambda_{p,x} > 0$. Here we identify a vector v from the tangent space of E with a vector in E if $d\pi(v) = 0$. From the second condition and the compactness of M we can cover M with finitely many open disks $D_1, D_2, \dots, D_r$ such that $\bar{D}_j \subset B_{p_j}$ and $f(\bar{D}_j) \cap f(\partial B_{p_j}) = \emptyset$ for each j and corresponding p_j .

Next choose smooth maps $\phi_j: V \rightarrow \mathbb{R}^1$ such that (i) $\phi_j > 0$, (ii) $\phi_j \circ f \equiv 0$ near ∂B_{p_j} and (iii) $\phi_j \circ f > 0$ on D_j . For each j let $\sigma_j: M \rightarrow \mathbb{R}^1$ be the map defined by $\sigma_j(x) = \phi_j \circ f(x)$ for $x \in B_{p_j}$ and $\sigma_j(x) = 0$ otherwise. Let $\sigma_j F_{p_j}: M \rightarrow E$ be the fibre product of σ_j and F_{p_j} and let $F = \sum_{j=1}^r \sigma_j F_{p_j}$ be the fibre sum. Note that $\ker dF \subset \ker d(\pi \circ F) = \ker df$. Since df is 1-1 away from S_1 , so is dF . Since $\dim \ker df_x = 1$ for $x \in S_1$ it suffices to show $dF_x(k) \neq 0$ for k a nonzero vector in the kernel of df_x . We will calculate dF in a neighborhood of x in which each F_{p_j} is of the form (f, h_j) where h_j maps into the fibre of E . Then

$$\begin{aligned} dF_x(k) &= \sum_{j=1}^r d(\sigma_j F_{p_j})_x(k) = \sum_{j=1}^r d(\sigma_j h_j)_x(k) = \sum_{j=1}^r \sigma_j d(h_j)_x(k) \\ &= \sum_{j=1}^r \sigma_j \lambda_{p,x} L(k) = \mu L(k) \end{aligned}$$

where $\mu > 0$. Therefore F is an immersion.

We now consider desingularization in the second sense. For a given line bundle $\pi': E' \rightarrow M$ with zero section $s: M \rightarrow E'$ we have

THEOREM 2. *For a generic map $f: M \rightarrow V$ there is a submersion $F': E' \rightarrow V$ with $f = F' \circ s$ if and only if the following conditions hold.*

- (i) $\text{rank } df \geq n - 1$ everywhere and
- (ii) $[S_{1,1}] = i^*w(E')$.

PROOF OF THEOREM 2. At a point $p \in S_1 - S_{1,1}$ the map f is a *fold*, i.e. f can be written as $f(x, y) = (x^2, y)$ for $x \in \mathbb{R}^1$ and $y \in \mathbb{R}^{n-1}$. If we have a submersion F' there is a unique *positive* direction on the fibre E'_p , $p \in S_1 - S_{1,1}$ determined by: v is *positive* if and only if for any neighborhood of p , N_p , we have $F(\alpha v) \in f(N_p)$ for all sufficiently small $\alpha > 0$. Since f is 2-generic we can choose a section of $E'_{|S_1}$ which is positive on $S_1 - S_{1,1}$ and which meets the zero section transversally with intersection $S_{1,1}$. Hence $[S_{1,1}]$ is the Stiefel-Whitney class of $E'_{|S_1}$ as is $i^*w(E')$.

Conversely if $[S_{1,1}] = i^*w(E')$ consider the line bundle over S_1 whose fibres are the lines normal to the image $f(S_1)$. For a 2-generic f this latter bundle is well defined and has Stiefel-Whitney class $[S_{1,1}]$. Therefore this bundle is equivalent to $E'_{|S_1}$. Using a bundle isomorphism we can define F' on $E'_{|S_1}$. We then extend F' so that it is constant on the fibres outside of a neighborhood of S_1 .

Since any class in $H^1(M; \mathbb{Z}_2)$ is the Stiefel-Whitney class of some line bundle over M , the following are immediate for a map f with $\text{rank } df \geq n - 1$ everywhere.

COROLLARY 1. *A generic $f: M \rightarrow V$ desingularizes in the first sense if and only if there is a class $w \in H^1(V; \mathbb{Z}_2)$ which satisfies the equation $i^*f^*(w) + [S_{1,1}] + w(N) = 0$.*

COROLLARY 2. *A generic $f: M \rightarrow V$ desingularizes in the second sense if and only if there is a class $w \in H^1(M; \mathbb{Z}_2)$ which satisfies the equation $i^*(w) = [S_{1,1}]$.*

With these corollaries we can establish the desingularization statements for the map of Figure 1.

REFERENCES

1. A. Haefliger, *Quelques remarques sur les applications différentiables d'une surface dans le plan*, Ann. Inst. Fourier (Grenoble) **10** (1960), 47–60.
2. M. Golubitsky and V. Guillemin, *Stable mappings and their singularities*, Springer-Verlag, New York, 1973.
3. Y. Saito, *On decomposable mappings of manifolds*, J. Math. Kyoto Univ. **1** (1961/62), 425–455.

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MASSACHUSETTS 02115

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, CHESTNUT HILL, MASSACHUSETTS 02167