FINITE GROUPS AND INVARIANT SOLUTIONS TO ONE-DIMENSIONAL PLATEAU PROBLEMS

DAVID BINDSCHADLER

ABSTRACT. Let G be a finite group of isometries acting on a complete Riemannian manifold. Suppose that B is a 0-dimensional boundary which is G -invariant. If the order of G divides the product of the cardinality of the orbit and the density of B at each point, then a G -invariant absolutely length minimizing integral current with boundary B can be constructed.

1. Introduction. Let G be a finite group of isometries acting on a complete Riemannian manifold. Let B denote a boundary consisting of a finite collection of points with integral densities. It is shown that if the order of the group divides the product of the cardinality of the orbit and the density of B at each point, then there exists an invariant solution to the oriented Plateau problem. (That is, there exists an invariant collection of oriented length minimizing geodesics with boundary B which minimizes total length among all collections of oriented curves with boundary B .) Known examples show that if the divisibility condition fails, there may exist boundaries with no invariant solution. (See [F1, 5.4.17].)

For other results concerning invariant solutions to the Plateau problem see [L], [B] and [BJ].

The author would like to thank the referee for suggesting the use of polyhedral chains instead of more abstract objects.

2. Polyhedral 1-chains in a metric space.

2.1. *Definitions and notations.* Let (X, d) be a metric space.

A 0-chain in X is an integer valued function β on X such that $\{x: \beta(x) \neq 0\}$ is a finite set.

A polyhedral 1-chain (or 1-pchain) in X is a nonnegative function $\alpha: X \times X \rightarrow \mathbb{Z}$ such that $\{(x, y) \in X \times X: \alpha(x, y) \neq 0\}$ is finite and $\alpha(x, y)\alpha(y, x) = 0$ for all $(x, y) \in X \times X$.

Let δ_x and $[x, y]$ denote the characteristic functions of $\{x\} \subset X$ and $\{(x, y)\} \subset X \times X$, respectively. Then for any 0-chain β and 1-pchain α , we have

$$\beta = \sum_{x \in X} \beta(x) \delta_x \quad \text{and} \quad \alpha = \sum_{(x, y) \in X \times X} \alpha(x, y) [x, y].$$

If β is a 0-chain and α is a 1-pchain, we define the *mass* of β , the *mass* of α and

Received by the editors August 27, 1979 and, in revised form, January 25, 1980.

AMS (MOS) subject classifications (1970). Primary 49F22; Secondary 49F20.

Key words and phrases. Plateau problem, area minimizing, integral current, invariant solution.

© 1980 American Mathematical Society
 0002-9939/80/0000-0617/\$02.50

the *boundary* of α by the equations

$$\mathbf{M}(\beta) = \sum_{x \in X} |\beta(x)|,$$

$$\mathbf{M}(\alpha) = \sum_{(x,y) \in X \times X} \alpha(x,y) d(x,y),$$

$$\partial\alpha = \sum_{(x,y) \in X \times X} \alpha(x,y)(\delta_y - \delta_x) = \sum_{y \in X} \sum_{x \in X} (\alpha(x,y) - \alpha(y,x))\delta_y.$$

A *cycle* is a 1-pchain with zero boundary.

If α and γ are two 1-pchains their *chain sum* is given by

$$\alpha \oplus \gamma(x,y) = \max\{0, \alpha(x,y) - \alpha(y,x) + \gamma(x,y) - \gamma(y,x)\}$$

for all $(x,y) \in X \times X$. The set of 1-pchains forms a commutative group under chain sums. Indeed, closure, existence of identity and commutativity are obvious. The inverse of α is the 1-pchain α given by $\ominus\alpha(x,y) = \alpha(y,x)$. And associativity follows from the equality $a = \max\{0, a\} - \max\{0, -a\}$. Also observe that

$$\alpha \oplus \gamma(x,y) + \alpha \oplus \gamma(y,x) = |\alpha(x,y) - \alpha(y,x) + \gamma(x,y) - \gamma(y,x)|,$$

$$\partial \ominus \alpha = -\partial\alpha \quad \text{and} \quad \partial(\alpha \oplus \gamma) = \partial\alpha \oplus \partial\gamma.$$

A 1-pchain is called *simple* if it has the form $\sum_{i=1}^l [x_{i-1}, x_i]$, where $x_i = x_j$ implies $i = j$ or $\{i, j\} = \{0, l\}$.

In case X were a complete Riemannian manifold and $T \in \mathbf{I}_1(X)$ (see [F1, pp. 670, 671]) consisted of a sum of oriented length minimizing geodesic arcs with integral density, then one can associate with T a 1-pchain α such that $\mathbf{M}(T) = \mathbf{M}(\alpha)$ and $\partial\alpha$ is essentially ∂T . Also, given a 1-pchain α , we can associate a sum of oriented length minimizing geodesic arcs with integral densities T by associating with $\alpha(x,y) > 0$ any length minimizing geodesic arc with boundary $\delta_y - \delta_x$ with density $\alpha(x,y)$. Then T and α have the same boundary and $\mathbf{M}(T) \leq \mathbf{M}(\alpha)$. It is this connection that lets us exploit mass minimizing 1-pchain in the study of mass minimizing integral currents.

2.2. THEOREM. *Given a polyhedral 1-chain α , there exists simple 1-pchains σ_i , $i = 1, \dots, l$, such that $\alpha = \ominus_{i=1}^l \sigma_i = \sum_{i=1}^l \sigma_i$, $\mathbf{M}(\alpha) = \sum_{i=1}^l \mathbf{M}(\sigma_i)$, and $\mathbf{M}(\partial\alpha) = \sum_{i=1}^l \mathbf{M}(\partial\sigma_i)$.*

PROOF. By induction on the integer $\sum_{(x,y) \in X \times X} \alpha(x,y)$, it suffices to find a nonzero simple 1-pchain σ so that

- (i) $\alpha \ominus \sigma = \alpha - \sigma$,
- (ii) $\mathbf{M}(\alpha) = \mathbf{M}(\alpha \ominus \sigma) + \mathbf{M}(\sigma)$,
- (iii) $\mathbf{M}(\partial\alpha) = \mathbf{M}(\partial(\alpha \ominus \sigma)) + \mathbf{M}(\partial\sigma)$.

This will be accomplished by finding points $x_M, \dots, x_N$, $M, N \in \mathbb{Z}$ with $M < N$ such that

- (a) $x_i \neq x_j$ unless $i = j$ or $\{i, j\} = \{M, N\}$,
- (b) $\alpha(x_{i-1}, x_i) > 0$ for $i = M+1, \dots, N$,
- (c) either (1) $x_M = x_N$ or (2) $\partial\alpha(x_M) < 0 < \partial\alpha(x_N)$.

For setting $\sigma = \sum_{i=M+1}^N [x_{i-1}, x_i]$, one obtains (i), hence (ii), from (a) and (b) and (iii) from (c).

The theorem is trivial if $\alpha = 0$, so choose x_0 and x_1 such that $\alpha(x_0, x_1) > 0$. If $\partial\alpha(x_1) > 0$, then set $N = 1$. If $\partial\alpha(x_1) < 0$, then we may assume that distinct points $x_0, \dots, x_k$ have been chosen such that $\alpha(x_{i-1}, x_i) > 0 > \partial\alpha(x_i)$ for all $i \in \{1, \dots, k\}$. Since $0 > \partial\alpha(x_k) = \sum_{y \in X} \alpha(y, x_k) - \alpha(x_k, y)$, there exists x_{k+1} with the property that $\alpha(x_k, x_{k+1}) > 0$.

If $x_{k+1} = x_i$ for some $i \in \{0, \dots, k\}$, then $x_i, \dots, x_{k+1}$ satisfies (a), (b), (c)(1).

If $x_{k+1} \notin \{x_0, \dots, x_k\}$ and $\partial\alpha(x_{k+1}) > 0$, then set $N = k + 1$.

If $x_{k+1} \notin \{x_0, \dots, x_k\}$ and $\partial\alpha(x_{k+1}) < 0$, then continue.

Since $\{(x, y): \alpha(x, y) > 0\}$ is a finite set, we either obtain the desired set or distinct points $x_0, \dots, x_N$ such that $\alpha(x_N) > 0$ and $\alpha(x_{i-1}, x_i) > 0$ for all $i \in \{1, \dots, N\}$. In the latter case a similar argument using decreasing indices can be used to complete the proof.

2.3. Given a 0-chain β such that $\sum_{x \in X} \beta(x) = 0$, we denote by M_β the number

$$\inf\{M(\alpha): \alpha \text{ is a 1-pchain with } \partial\alpha = \beta\}.$$

Observe that M_β is finite since one can construct a 1-pchain with boundary β by adding an appropriate number of characteristic functions $[x, y]$ where $\beta(x) < 0 < \beta(y)$. Note also that $M_\beta = \inf\{M(\alpha_0 \oplus \zeta): \zeta \text{ is a cycle}\}$, where α_0 is any 1-pchain with $\partial\alpha_0 = \beta$.

2.4. THEOREM. *If β is a 0-chain in X such that $\sum \beta(x) = 0$, then there exists a 1-pchain α_0 with $\partial\alpha_0 = \beta$ and $M(\alpha_0) = M_\beta$.*

PROOF. For any 1-pchain α with $\partial\alpha = \beta$ one can find σ_i , $i = 1, \dots, l$, as in 2.2. Now define

$$\gamma_i = \begin{cases} [x, y] & \text{if } \partial\sigma_i = \delta_y - \delta_x, \\ 0 & \text{if } \partial\sigma_i = 0, \end{cases}$$

and $\gamma = \bigoplus_{i=1}^l \gamma_i = \sum_{i=1}^l \gamma_i$. Then $\partial\gamma = \sum_{i=1}^l \partial\sigma_i = \alpha$ and by the triangle inequality

$$M(\gamma) = \sum_{i=1}^l M(\gamma_i) < \sum_{i=1}^l M(\sigma_i) = M(\alpha).$$

Observing that $\gamma(x, y) > 0$ implies $\beta(x) < 0 < \beta(y)$ we conclude

$$M_\beta = \inf\{M(\gamma): \partial\gamma = \beta \text{ and } \gamma(x, y) > 0 \text{ implies } \beta(x) < 0 < \beta(y)\}.$$

The infimum is attained, since the latter set is finite.

2.5. THEOREM. *For any positive integer k and 0-chain β in X with $\sum_{x \in X} \beta(x) = 0$, the equality $M_{k\beta} = kM_\beta$ holds.*

PROOF. If α is a 1-pchain, then $\partial k\alpha = k\partial\alpha$ and $M(k\alpha) = kM(\alpha)$ from which $M_{k\beta} \leq kM_\beta$ follows.

For the reverse inequality, we suppose α is a 1-pchain such that $\partial\alpha = \beta$ and $M(\alpha) = M_\beta$. Given any cycle ζ , we use 2.2 to decompose ζ into a sum of simple cycles σ_i , $i = 1, \dots, l$. For each $i \in \{1, \dots, l\}$ and $(x, y) \in X \times X$, let

$$s_i(x, y) = \begin{cases} -1 & \text{if } (\alpha(x, y) - \alpha(y, x))(\sigma_i(x, y) - \sigma_i(y, x)) < 0, \\ 1 & \text{otherwise.} \end{cases}$$

Since

$$\begin{aligned} \alpha \oplus \sigma_i(x, y) + \alpha \oplus \sigma_i(y, x) &= |\alpha(x, y) - \alpha(y, x) + \sigma_i(x, y) - \sigma_i(y, x)| \\ &= |\alpha(x, y) - \alpha(y, x)| + s_i(x, y)|\sigma_i(x, y) - \sigma_i(y, x)|, \end{aligned}$$

we have

$$\mathbf{M}(\alpha \oplus \sigma_i) = \mathbf{M}(\alpha) + \frac{1}{2} \sum_{(x, y) \in X \times X} s_i(x, y) |\sigma_i(x, y) - \sigma_i(y, x)| d(x, y).$$

By the mass minimality of α , we conclude

$$\sum_{(x, y) \in X \times X} s_i(x, y) |\sigma_i(x, y) - \sigma_i(y, x)| d(x, y) \geq 0.$$

For each $(x, y) \in X \times X$ let $I(x, y) = \{i: s_i(x, y) < 0\}$. Then

$$\begin{aligned} k\alpha \oplus \sum_{i=1}^l \sigma_i(x, y) + k\alpha \oplus \sum_{i=1}^l \sigma_i(y, x) &= \left| k\alpha(x, y) - k\alpha(y, x) + \sum_{i=1}^l \sigma_i(x, y) - \sigma_i(y, x) \right| \\ &= \left| k\alpha(x, y) - k\alpha(y, x) + \sum_{i \notin I(x, y)} \sigma_i(x, y) - \sigma_i(y, x) \right. \\ &\quad \left. + \sum_{i \in I(x, y)} \sigma_i(x, y) - \sigma_i(y, x) \right| \\ &= \left| |k\alpha(x, y) - k\alpha(y, x)| + \sum_{i \notin I(x, y)} |\sigma_i(x, y) - \sigma_i(y, x)| \right. \\ &\quad \left. - \sum_{i \in I(x, y)} |\sigma_i(x, y) - \sigma_i(y, x)| \right| \\ &= \left| |k\alpha(x, y) - k\alpha(y, x)| + \sum_{i=1}^l s_i(x, y) |\sigma_i(x, y) - \sigma_i(y, x)| \right| \\ &> |k\alpha(x, y) - k\alpha(y, x)| + \sum_{i=1}^l s_i(x, y) |\sigma_i(x, y) - \sigma_i(y, x)|. \end{aligned}$$

Hence

$$\begin{aligned} \mathbf{M}(k\alpha \oplus \zeta) &> \mathbf{M}(k\alpha) + \frac{1}{2} \sum_{(x, y) \in X \times X} \sum_{i=1}^l s_i(x, y) |\sigma_i(x, y) - \sigma_i(y, x)| d(x, y) \\ &> \mathbf{M}(k\alpha) = k\mathbf{M}_\beta. \end{aligned}$$

Thus the inequality $\mathbf{M}_{k\beta} > k\mathbf{M}_\beta$ is established.

3. Invariant solutions to Plateau problems.

3.1. Let G be a finite group acting on X so that $d(g(x), g(y)) = d(x, y)$ for all $x, y \in X$ and $g \in G$. Let $G(x)$ denote the orbit of x and G_x denote the isotropy

subgroup of G at x . Identifying orbits of G to points gives the orbit space Z . Let $\pi: X \rightarrow Z$ be the canonical projection. For each $u, v \in Z$ define

$$d'(u, v) = \min\{d(x, y): x \in \pi^{-1}(u), y \in \pi^{-1}(v)\}.$$

The 0-chain β and the 1-chain α are called invariant if $\beta(gx) = \beta(x)$ and $\alpha(g(x), g(y)) = \alpha(x, y)$ for all $x, y \in X$.

3.2. THEOREM. *If β is a G -invariant 0-chain with $\sum_{x \in X} \beta(x) = 0$ and the order of G divides $\beta(x) \text{ card } G(x)$ for each $x \in X$, then there exists a G -invariant 1-pchain α such that $\partial\alpha = \beta$ and $\mathbf{M}(\alpha) = \mathbf{M}_\beta$.*

PROOF. Let r be the order of G . Define $\beta': Z \rightarrow \mathbf{Z}$ by

$$\beta'(u) = \frac{1}{r} \sum_{x \in \pi^{-1}(u)} \beta(x).$$

Since β is G -invariant and r divides $\beta(x) \text{ card } G(x)$, β' is a 0-chain in Z . If γ is a 1-pchain in X with $\partial\gamma = \beta$, then one can define the 1-pchain in Z by

$$\gamma'(u, v) = \sum_{(x, y) \in \pi^{-1}(u) \times \pi^{-1}(v)} \gamma(x, y).$$

Then $\partial\gamma' = r\beta'$ and $\mathbf{M}(\gamma') \leq \mathbf{M}(\gamma)$. Hence by 2.5, $\mathbf{M}_\beta \geq \mathbf{M}_{r\beta'} = r\mathbf{M}_{\beta'}$.

Now choose a 1-pchain α' such that $\partial\alpha' = \beta'$ and $\mathbf{M}(\alpha') = \mathbf{M}_{\beta'}$. For each $(u, v) \in Z \times Z$ such that $\alpha'(u, v) > 0$, we find $x_u \in \pi^{-1}(u)$ and $y_{u,v} \in \pi^{-1}(v)$ such that $d(x_u, y_{u,v}) = d'(u, v)$. Let α be the 1-pchain in X given by

$$\alpha = \sum_{(u,v) \in Z \times Z} \sum_{g \in G} \alpha'(u, v) [g(x_u), g(y_{u,v})].$$

Then

$$\begin{aligned} \mathbf{M}(\alpha) &= \sum_{(u,v) \in Z \times Z} \sum_{g \in G} \alpha(g(x_u), g(y_{u,v})) d(x_u, y_{u,v}) \\ &= \sum_{(u,v) \in Z \times Z} \sum_{g \in G} \alpha'(u, v) d'(u, v) = r\mathbf{M}(\alpha'), \end{aligned}$$

and

$$\begin{aligned} \partial\alpha &= \sum_{(u,v) \in Z \times Z} \sum_{g \in G} \alpha'(u, v) \delta_{g y_{u,v}} - \sum_{(u,v) \in Z \times Z} \sum_{g \in G} \alpha'(u, v) \delta_{g x_u} \\ &= \sum_{u \in Z} \left(\sum_{v \in Z} \alpha'(v, u) - \alpha'(u, v) \right) \sum_{g \in G} \delta_{g w_u} \\ &= \sum_{u \in Z} \beta'(u) \sum_{g \in G} \delta_{g w_u}, \end{aligned}$$

where w_u is any point in $\pi^{-1}(u)$. Since $(\text{card } G(x))(\text{card } G_x) = r$, we have that

$$\partial\alpha(x) = \beta'(\pi(x)) \text{ card } G_x = \beta(x).$$

Hence $\mathbf{M}_\beta \leq \mathbf{M}(\alpha) = r\mathbf{M}(\alpha') = r\mathbf{M}_{\beta'} \leq \mathbf{M}_\beta$.

3.3. For an explanation of the terms and notation used in the following corollary, we refer the reader to [F1, pp. 670, 671].

COROLLARY. Let X be a complete Riemannian manifold and G be a finite group of isometries of X . If B is a G -invariant 0-dimensional rectifiable current which is a boundary and r divides $\Theta(\|B\|, x) \operatorname{card} G(x)$ for all $x \in X$, then there exists a G -invariant $T \in I_1(X)$ such that $\partial T = B$ and

$$M(T) = \inf \{M(R) : R \in I_1(X) \text{ and } \partial R = B\}.$$

PROOF. Repeat the argument of 3.2 except in the definition of α use $g_{\#} L(x_u, y_{u,v})$ where $L(x_u, y_{u,v})$ is any oriented length minimizing arc from x_u to $y_{u,v}$.

REFERENCES

- [B] D. E. Bindschadler, *Invariant solutions to the oriented Plateau problem of maximal codimension*, Trans. Amer. Math. Soc. **261** (1980), 439–462.
- [BJ] J. E. Brothers, *Invariance of solutions to invariant parametric variational problems*, Trans. Amer. Math. Soc. **262** (1980), 159–179.
- [F1] H. Federer, *Geometric measure theory*, Springer-Verlag, New York, 1969.
- [F2] ———, *Real flat chains, cochains and variational problems*, Indiana Univ. Math. J. **24** (1974), 351–407.
- [L] H. B. Lawson, *The equivariant Plateau problem and interior regularity*, Trans. Amer. Math. Soc. **173** (1973), 231–249.

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MICHIGAN 48202