ON OPERATOR RANGES

BHUSHAN L. WADHWA

ABSTRACT. If f is any vector-valued bounded function defined on open set D of the complex plane, and T is any bounded linear operator on a Hilbert space such that $\sigma_R(T^*)$ is empty and if $(T - zI)^2f(z) = x$ for all z in D then f is analytic.

Let T be a bounded linear operator on a Hilbert space H . Let f be an H -valued function defined on an open set D of the complex plane. Suppose $(T - zI)f(z) = x$ for all z in D and for a fixed x in H . The question arises: What type of conditions on the operator T and on the function f will be sufficient to insure that f is analytic on D ? This question has been implicitly discussed in Clancey [1], Johnson [4], Putnam [5], [6], Radjabalipour [7], and Stampfli-Wadhwa [9], [10], under various conditions related to normality on T . One of the main results is: If T is a hyponormal operator ($TT^* \leq T^*T$) and f is any function satisfying the above equation then f is analytic in D . See [1]. On the other hand, there is a cohyponormal operator T , (T^* is hyponormal) and a bounded function f satisfying the above equation which fails to be analytic on D . See [6].

In this note, by a simple argument, we shall show that if $(T - zI)^2f(z) = x$ for all z in D and if f is bounded on D and $\sigma_R(T^*) = \emptyset$ (range of $T^* - \bar{z}I$ is dense in H for all z) then f is analytic. We shall use this result to give an alternative proof of a classical result of Stampfli [8] about quadratically hyponormal operators.

The following lemma is implicitly contained in most of the references mentioned previously. We include it for the sake of completeness.

LEMMA. Let $(T - zI)f(z) = x$ for all z in D be such that f is bounded and $\sigma_R(T^*) = \emptyset$ then f is weakly continuous.

PROOF. For any z and z_0 in D ,

$$\begin{aligned}(f(z) - f(z_0), (T^* - \bar{z}_0I)y) &= ((T - z_0I)(f(z) - f(z_0)), y) \\ &= (z - z_0)(f(z), y) \quad \text{for all } y \text{ in } H.\end{aligned}$$

Since f is bounded and range of $(T^* - \bar{z}_0I)$ is dense in H , it follows that f is weakly continuous.

THEOREM. Let g be a bounded vector-valued function such that $(T - zI)^2g(z) = x$ for all z in D and let $\sigma_R(T^*) = \emptyset$. Then g is analytic on D .

PROOF. Let $f(z) = (T - zI)g(z)$. Then $(T - zI)f(z) = x$ for all z in D , f is bounded and hence weakly continuous on each bounded subset D_0 of D . Now for

any z and z_0 in D_0 ,

$$((f(z) - f(z_0))/(z - z_0), (T^* - \bar{z}_0 I)y) = (f(z), y),$$

and

$$\begin{aligned} \lim_{z \rightarrow z_0} ((f(z) - f(z_0))/(z - z_0), (T^* - \bar{z}_0 I)y) &= (f(z_0), y) \\ &= ((T - z_0 I)g(z_0), y) = (g(z_0), (T^* - \bar{z}_0 I)y) \end{aligned}$$

for all y in H .

Since range of $(T^* - \bar{z}_0 I)$ is dense in H , f is analytic and $f'(z) = g(z)$ for all z in D .

Let T be a quadratically hyponormal operator ($aT^2 + bT + cI$ is hyponormal for all complex numbers a, b and c). Let $\rho(T, x)$ be the local resolvent of the vector x with respect to the operator T (see Dunford and Schwartz [3, p. 1935]).

COROLLARY (STAMPFLI [8]). *If T is a quadratically hyponormal operator with $\sigma_R(T) = \emptyset$ then $\rho(T; x) \subset \overline{\rho(T^*, x)}$. (The bar denotes the complex conjugate of the set.)*

PROOF. Let $z_0 \in \rho(T, x)$; thus there exists an analytic function defined on a bounded set D containing z_0 such that $(T - zI)f(z) = x$. Since f is analytic, a simple computation shows that $(T - zI)^2 f'(z) = x$ for all z in D . Since T is quadratically hyponormal, $(T - zI)^2(T^* - \bar{z}I)^2 \leq (T^* - \bar{z}I)^2(T - zI)^2$. By Douglas [2], there is a contraction $K(z)$ such that $(T - zI)^2 = (T^* - \bar{z}I)^2 K(z)$. Consequently, $(T^* - \bar{z}I)^2 g(\bar{z}) = x$ where $g(\bar{z}) = K(z)f'(z)$. Thus $g(\bar{z})$ is bounded for $z \in D$. Using the Theorem we conclude that $g(z)$ is analytic for all z in $\bar{D}$ and $(T^* - zI)(T^* - \bar{z}I)g(z) = x$ for all z in $\bar{D}$; thus $z_0 \in \overline{\rho(T^*, x)}$.

REFERENCES

1. K. Clancey, *On local spectra of seminormal operators*, Proc. Amer. Math. Soc. **72** (1978), 473–479.
2. R. G. Douglas, *On majorization, factorization and range inclusion of operators on Hilbert space*, Proc. Amer. Math. Soc. **17** (1966), 413–415.
3. N. Dunford and J. Schwartz, *Linear operators. III: Spectral operators*, Pure and Appl. Math., Vol. VII, Interscience, New York-London-Sydney, 1971.
4. B. E. Johnson, *Continuity of linear operators commuting with continuous linear operators*, Trans. Amer. Math. Soc. **128** (1967), 88–102.
5. C. R. Putnam, *Ranges of normal and subnormal operators*, Michigan Math. J. **18** (1971), 33–36.
6. ———, *Hyponormal contractions and strong power convergence*, Pacific J. Math. **57** (1975), 531–538.
7. M. Radjabalipour, *Ranges of hyponormal operators*, Illinois J. Math. **21** (1977), 70–75.
8. J. G. Stampfli, *Analytic extensions and spectral localization*, J. Math. Mech. **19** (1966), 287–296.
9. J. G. Stampfli and B. L. Wadhwa, *An asymmetric Putnam-Fuglede theorem for dominant operators*, Indiana Univ. Math. J. **25** (1976), 359–365.
10. ———, *On dominant operators*, Monatsh. Math. **84** (1977), 143–153.

DEPARTMENT OF MATHEMATICS, CLEVELAND STATE UNIVERSITY, CLEVELAND, OHIO 44115