PREDICTION n UNITS OF TIME AHEAD¹

TAKAHIKO NAKAZI AND KATUTOSHI TAKAHASHI

ABSTRACT. The purpose of this note is to give a simple expression in terms of w of the quantities

$$\rho_n(w) = \inf_f \int_0^{2\pi} |1 + e^{in\theta} f|^2 w \, d\theta / 2\pi \quad (n = 0, 1, 2, \dots),$$

where f ranges over the analytic trigonometric polynomials with mean value zero and w is nonnegative and summable on the circle.

Helson [2, pp. 21–22] said that it is unreasonable to expect to have a simple expression in terms of w for the quantities ρ_n except $n = 0$. $\rho_0(w) = \exp \int_0^{2\pi} \log w \, d\theta / 2\pi$ is the famous Szegő theorem. We may assume $\log w$ is summable, because otherwise $\rho_n(w) = 0$ for all n .

THEOREM. *Let w be nonnegative and summable on the circle. Suppose $\log w$ is summable and*

$$\log w(\theta) \sim \sum_{j=-\infty}^{\infty} a_j e^{ij\theta}.$$

Then

$$\begin{aligned} \rho_n(w) &= \inf_f \int_0^{2\pi} |1 + e^{in\theta} f|^2 w \, d\theta / 2\pi \\ &= e^{a_0} \times \sum_{j=0}^n \left| \sum' \frac{a_1^{m_1} a_2^{m_2} \cdots a_n^{m_n}}{m_1! m_2! \cdots m_n!} \right|^2 \end{aligned}$$

where $n \geq 1$ and f ranges over the analytic trigonometric polynomials with mean value zero and Σ' is the summation of all permutations of nonnegative integers $m_1, m_2, \dots, m_n$ with $m_1 + 2m_2 + \cdots + nm_n = j$ for each j .

PROOF. Set $g_1 = \sum_{l=1}^n a_l z^l$ and $g_2 = a_0/2 + \sum_{l \geq n+1} a_l z^l$, then their radial limits satisfy $w(\theta) = |\exp g_1(e^{i\theta})|^2 |\exp g_2(e^{i\theta})|^2$ a.e. θ . $\exp(g_1 + g_2)$ is an outer function [3, p. 61] and so $\exp g_2$ is outer. Hence, if we note that there exist positive numbers ε and M with $0 < \varepsilon < |\exp g_1(e^{i\theta})|^2 < M < \infty$, as in the proof of Szegő's theorem

Received by the editors July 9, 1979 and, in revised form, September 11, 1979 and November 9, 1979.
AMS (MOS) subject classifications (1970). Primary 60G25, 30A98, 46J15.

Key words and phrases. Prediction theory, Szegő's theorem, n units of time ahead, outer function, Fourier coefficients.

¹Research supported in part from Grant-in-Aid Encouragement of Young Scientists.

© 1980 American Mathematical Society
0002-9939/80/0000-0627/\$01.50

$$\begin{aligned}
\rho_n(w) &= \inf_f \int |1 + e^{in\theta} f|^2 |\exp g_2|^2 |\exp g_1|^2 d\theta / 2\pi \\
&= \inf_f \int |\exp g_2 + e^{in\theta} f \exp g_2|^2 |\exp g_1|^2 d\theta / 2\pi \\
&= \inf_f \int |e^{a_0/2} + e^{in\theta} f|^2 |\exp g_1|^2 d\theta / 2\pi \\
&= e^{a_0} \inf_f \int |1 + e^{in\theta} f|^2 |\exp g_1|^2 d\theta / 2\pi.
\end{aligned}$$

Since $\exp g_1$ is an outer function, if the Fourier coefficients of $\exp g_1$ are $\{b_j\}$, then (cf. [1, pp. 184–187], [2, p. 22])

$$\begin{aligned}
&\inf_f \int |1 + e^{in\theta} f|^2 |\exp g_1|^2 d\theta / 2\pi \\
&= \sum_{j=0}^n |b_j|^2 \exp g_1(z) = \prod_{i=1}^n \exp(a_i z^i) = \sum \frac{(a_1 z)^{m_1} \cdots (a_n z^n)^{m_n}}{m_1! \cdots m_n!}
\end{aligned}$$

where the m_j range independently over nonnegative integers. This implies the theorem.

The theorem shows the following:

$$\begin{aligned}
\rho_1(w) &= \inf_f \int_0^{2\pi} |1 + e^{i\theta} f|^2 w d\theta / 2\pi \\
&= \exp \int_0^{2\pi} \log w d\theta / 2\pi \left\{ 1 + \left| \int_0^{2\pi} (\log w) e^{i\theta} d\theta / 2\pi \right|^2 \right\}.
\end{aligned}$$

If w is nonnegative and summable, and $\log w$ is summable, it is known (cf. [2, p. 20]) that $w = |g|^2$ for some outer g . The theorem gives a simple expression in terms of w of the Fourier coefficients of outer function g (cf. [4]).

We wish to express our appreciation to the referee for several valuable suggestions.

REFERENCES

1. U. Grenander and G. Szegő, *Toeplitz forms and their applications*, Univ. of California Press, Berkeley and Los Angeles, 1958.
2. H. Helson, *Lectures on invariant subspaces*, Academic Press, New York, 1964.
3. K. Hoffman, *Banach spaces of analytic functions*, Prentice-Hall, Englewood Cliffs, N.J., 1962.
4. T. L. Kriete III, *On the Fourier coefficients of outer functions*, Indiana Univ. Math. J. **20** (1970), 147–155.

DIVISION OF APPLIED MATHEMATICS, RESEARCH INSTITUTE OF APPLIED ELECTRICITY, HOKKAIDO UNIVERSITY, SAPPORO, JAPAN