REPRESENTATIONS OF BAER *-SEMIGROUPS

STANLEY P. GUDDER AND JOHN R. MICHEL

ABSTRACT. Various types of Hilbert space representations for a Baer *-semigroup S are defined. The representations are characterized in terms of *-positive functions on S which possess additivity and consistency properties.

1. Introduction. In this paper we characterize those Baer *-semigroups that admit various types of representations in Hilbert space. These characterizations are formulated in terms of the existence of *-positive functions possessing certain properties. Such formulations have been used for general semigroups and *-semigroups [1], [10], [12], but the additional properties needed for Baer *-semigroups have not been previously considered.

Representations of Baer *-semigroups are not only of interest in their own right, they can be important in the study of quantum logics [7], [11], [13], [18] and operational quantum mechanics [2], [3], [7], [9]. Because of the intimate connection between Baer *-semigroups and orthomodular lattices [5], [6], representations of the former can provide embeddings of the latter [8]. This connection is not only of a purely mathematical character; it also results from a deeper understanding of the physical meaning of "operation" in quantum mechanics [15], [16]. For this reason, representations of Baer *-semigroups may be physically interpreted as representations of the set of operations for a quantum system. One then obtains a representation theory which is more fundamental than the usual representation theory for the C^* -algebra of bounded observables for a quantum system [4], [9].

2. Definitions and notation. Let S be a Baer *-semigroup [5], [15] whose set of closed projections we denote by $P'(S)$ and let $L(H)$ denote the set of bounded linear operators on a complex Hilbert space H . For $A \in L(H)$, A' denotes the orthogonal projection onto the null space of A . A *-representation ρ of S in H is a map $\rho: S \rightarrow L(H)$ satisfying

$$\rho(0) = 0, \quad \rho(1) = I, \quad (1)$$

$$\rho(st) = \rho(s)\rho(t) \quad \text{for all } s, t \in S, \quad (2)$$

$$\rho(s^*) = \rho(s)^* \quad \text{for all } s \in S, \quad (3)$$

$$\|\rho(s)\| \leq 1 \quad \text{for all } s \in S. \quad (4)$$

Received by the editors September 26, 1979 and, in revised form, January 10, 1980 and April 14, 1980.
 1980 *Mathematics Subject Classification*. Primary 20M30; Secondary 81B10.

Key words and phrases. Baer *-semigroups, Hilbert space representations, positive definite functions, quantum logic.

© 1981 American Mathematical Society
 0002-9939/81/0000-0050/\$02.75

We now define various types of $*$ -representations. A *weak Baer $*$ -representation* of S is a $*$ -representation ρ for which

$$\rho(e') = \rho(e)' \quad \text{for all } e \in P'(S). \quad (5)$$

A *Baer $*$ -representation* of S is a $*$ -representation ρ for which

$$\rho(s') = \rho(s)' \quad \text{for all } s \in S. \quad (6)$$

A *weak σ^* -representation* of S is a $*$ -representation ρ satisfying

$$\begin{aligned} &\text{if } e_i \in P'(S) \text{ are mutually orthogonal, } i = 1, 2, \dots, \text{ and} \\ &\bigvee e_i \text{ exists, then } \rho(\bigvee e_i) = \sum \rho(e_i). \end{aligned} \quad (7)$$

A *σ^* -representation* of S is a Baer $*$ -representation ρ satisfying (7).

A $*$ -representation $\rho: S \rightarrow L(H)$ is *cyclic* if there exists an $x \in H$ such that the linear hull of $\{\rho(s)x: s \in S\}$ is dense in H . It is not difficult to show that any $*$ -representation ρ is a direct sum of cyclic ones and that these cyclic representations preserve (5), (6), or (7) if they hold for ρ . For this reason, we shall mainly restrict our attention to cyclic $*$ -representations.

A bounded complex-valued function ϕ on S is *$*$ -positive* if $\phi(0) = 0$ and for any finite collections $\{\lambda_1, \dots, \lambda_n\} \subseteq C$, $\{s_1, \dots, s_n\} \subseteq S$, we have

$$\sum_{i,j=1}^n \bar{\lambda}_i \lambda_j \phi(s_i^* s_j) > 0.$$

Notice that if ϕ is $*$ -positive and $t \in S$, then the function $\phi_t(s) = \phi(t^* s t)$ is also $*$ -positive. A $*$ -positive function ϕ is *additive* if $\phi_t(e) + \phi_t(e') = \phi_t(1)$ for every $t \in S$, $e \in P'(S)$, and ϕ is *σ -additive* (or *normal*) if ϕ is additive and for any sequences $e_i \in P'(S)$ which are mutually orthogonal and $\bigvee_{i=1}^{\infty} e_i$ exists for n sufficiently large, we have

$$\lim_{n \rightarrow \infty} \phi_t\left(\bigvee_{i=1}^n e_i\right) = 0 \quad \text{for every } t \in S.$$

LEMMA 1. Let ϕ be a $*$ -positive function on S .

(a) If $e, f \in P'(S)$ and $ef = fe$, then $\phi(e) + \phi(f) - \phi(e f) \leq \phi(e \vee f)$.

(b) ϕ is additive if and only if for any finite set of mutually orthogonal $e_i \in P'(S)$, $i = 1, \dots, n$, $\phi_t(\bigvee e_i) = \sum \phi_t(e_i)$ for every $t \in S$.

(c) ϕ is σ -additive if and only if for every sequence of mutually orthogonal $e_i \in P'(S)$ such that $\bigvee_{i=1}^{\infty} e_i$ exists for every n , $\phi_t(\bigvee e_i) = \sum \phi_t(e_i)$ for every $t \in S$.

PROOF. (a) Take $s_1 = e \vee f$, $s_2 = e$, $s_3 = f$, $s_4 = ef$, $\lambda_1 = \lambda_4 = -1$, $\lambda_2 = \lambda_3 = 1$, in the definition of a $*$ -positive function.

(b) If $e_1 \perp e_2$, then from (a) we have

$$\phi_t(e'_1) + \phi_t(e'_2) - \phi_t(e'_1 e'_2) \leq \phi_t(e'_1 \vee e'_2).$$

Assuming ϕ is additive gives

$$\begin{aligned} &[\phi_t(1) - \phi_t(e_1)] + [\phi_t(1) - \phi_t(e_2)] - [\phi_t(1) - \phi_t((e'_1 e'_2)')] \\ &\leq \phi_t(1) - \phi_t((e'_1 \vee e'_2)'). \end{aligned}$$

Using De Morgan's laws and rearranging gives $\phi_t(e_1 \vee e_2) \leq \phi_t(e_1) + \phi_t(e_2)$. The opposite inequality follows from (a).

(c) If ϕ is σ -additive, then

$$\phi_t(\bigvee e_i) = \phi_t\left(\bigvee_{i=1}^n e_i\right) + \phi_t\left(\bigvee_{i=n+1}^{\infty} e_i\right) = \sum_{i=1}^n \phi_t(e_i) + \phi_t\left(\bigvee_{i=n+1}^{\infty} e_i\right).$$

Hence,

$$\lim_{n \rightarrow \infty} \left| \phi_t(\bigvee e_i) - \sum_{i=1}^n \phi_t(e_i) \right| = \lim_{n \rightarrow \infty} \phi_t\left(\bigvee_{i=n+1}^{\infty} e_i\right) = 0. \quad \square$$

We say that a *-positive function ϕ is *consistent* if $\phi(s*s) = 0$ implies that $\phi(s') = \phi(1)$. In a Baer *-semigroup, $s*s = 0$ implies $s' = 1$. Hence, a consistent *-positive function is one that respects the prior implication. For example, if ϕ is faithful ($\phi(s*s) = 0$ implies $s = 0$), then ϕ_t is consistent for every $t \in S$. Indeed, $\phi_t(s*s) = 0$ implies that $\phi(st*st) = 0$. Hence, $st = 0$ so that $s't = t$ and

$$\phi_t(s') = \phi(t*s't) = \phi(t*t) = \phi_t(1).$$

In the next paragraph we shall define ϕ to be strongly consistent if limits of certain *-positive functions derived from ϕ are consistent.

Let ϕ be a *-positive function on S and let $\{c_1, \dots, c_m\} \subseteq C$, $\{t_1, \dots, t_m\} \subseteq S$. Define the function $\phi_1 \equiv \phi\{c_i, t_i\}$ by $\phi_1(s) = \sum_{k,i} \bar{c}_i c_k \phi(t_k^* s t_i)$. It is not hard to show that ϕ_1 is again *-positive. Now let $\{c_1^k, \dots, c_{n(k)}^k\} \subseteq C$, $\{t_1^k, \dots, t_{n(k)}^k\} \subseteq S$, $k = 1, 2, \dots$, be a sequence of finite sets and let $\phi_k \equiv \phi\{c_i^k, t_i^k\}$. We say that the sequence ϕ_k converges if

$$\lim_{k \rightarrow \infty} \phi_k(1) = \operatorname{Re} \lim_{k, l \rightarrow \infty} \sum_{i,j} \bar{c}_i^l c_j^k \phi(t_i^{l*} t_j^k).$$

It can be shown using the techniques of Theorem 2 to follow, that if ϕ_k converges, then $\lim_{k \rightarrow \infty} \phi_k(s)$ exists for every $s \in S$. It is clear that the function $\psi(s) \equiv \lim_{k \rightarrow \infty} \phi_k(s)$ is again *-positive on S . We call ψ a *limit function* for ϕ . We say that ϕ is *strongly consistent* if every limit function for ϕ is consistent.

3. Representation theorem.

THEOREM 2. (a) Let $\rho: S \rightarrow L(H)$ be a *-representation, $x \in H$, and $\phi(s) = \langle \rho(s)x, x \rangle$, $s \in S$. If ρ is weak Baer, weak σ , Baer, or σ , respectively, then ϕ is an additive, σ -additive, strongly consistent, σ -additive strongly consistent, respectively, *-positive function.

(b) Conversely, if ϕ is an additive, σ -additive, strongly consistent, or σ -additive strongly consistent, respectively, *-positive function on S , then there exists a weak Baer, weak σ , Baer, σ , respectively, *-representation $\rho: S \rightarrow L(H)$ which is cyclic with cyclic vector $x \in H$ such that $\phi(s) = \langle \rho(s)x, x \rangle$ for all $s \in S$.

(c) There exists an injective *-representation of each of the previous types if and only if there exists a separating set of *-positive functions of the prior corresponding type.

PROOF. (a) We first show that $\phi(s) = \langle \rho(s)x, x \rangle$ is $*$ -positive. Clearly, $\phi(0) = 0$ and ϕ is bounded by Schwarz's inequality. Moreover,

$$\begin{aligned} \sum \bar{\lambda}_i \lambda_j \phi(s_i^* s_j) &= \sum \bar{\lambda}_i \lambda_j \langle \rho(s_j)x, \rho(s_i)x \rangle \\ &= \left\langle \sum \lambda_j \rho(s_j)x, \sum \lambda_i \rho(s_i)x \right\rangle \\ &= \left\| \sum \lambda_i \rho(s_i)x \right\|^2 > 0. \end{aligned}$$

If ρ is a weak Baer $*$ -representation, then ϕ is additive since for $e \in P'(S)$, $t \in S$, we have

$$\begin{aligned} \phi_t(e) + \phi_t(e') &= \langle \rho(t^*et)x, x \rangle + \langle \rho(t^*e't)x, x \rangle \\ &= \langle \rho(e)\rho(t)x, \rho(t)x \rangle + \langle \rho(e')\rho(t)x, \rho(t)x \rangle. \end{aligned}$$

Now $\rho(e)$ is an orthogonal projection since $\rho(e)^2 = \rho(e^2) = \rho(e) = \rho(e^*) = \rho(e)^*$. Hence, $\rho(e) + \rho(e') = I$ and

$$\phi_t(e) + \phi_t(e') = \langle \rho(t)x, \rho(t)x \rangle = \langle \rho(t^*t)x, x \rangle = \phi_t(1).$$

If ρ is a weak σ^* -representation, then ρ is a weak Baer $*$ -representation, so ϕ is additive. Moreover, ϕ is σ -additive since for $e_i \in P'(S)$ satisfying the requirements of the definition we have

$$\begin{aligned} \phi_t\left(\bigvee_{i=1}^{\infty} e_i\right) &= \left\langle \rho\left(\bigvee_{i=1}^{\infty} e_i\right)\rho(t)x, \rho(t)x \right\rangle \\ &= \left\langle \sum_{i=1}^{\infty} \rho(e_i)\rho(t)x, \rho(t)x \right\rangle = \sum_{i=1}^{\infty} \langle \rho(e_i)\rho(t)x, \rho(t)x \rangle \\ &= \sum_{i=1}^{\infty} \|\rho(e_i)\rho(t)x\|^2 \leq \|\rho(t)x\|^2. \end{aligned}$$

Since the infinite series converges,

$$\lim_{n \rightarrow \infty} \phi_t\left(\bigvee_{i=m}^{\infty} e_i\right) = \lim_{n \rightarrow \infty} \sum_{i=n}^{\infty} \|\rho(e_i)\rho(t)x\|^2 = 0.$$

Suppose ρ is a Baer $*$ -representation and ψ is a limit function for ϕ . Then there exist sets $\{c_1^k, \dots, c_{n(k)}^k\} \subseteq C$, $\{t_1^k, \dots, t_{n(k)}^k\} \subseteq S$ such that $\phi_k \equiv \phi\{c_i^k, t_i^k\}$ converges and $\psi(s) = \lim \phi_k(s)$ for every $s \in S$. Let $x_k = \sum_i c_i^k \rho(t_i^k)x$. Then x_k is Cauchy since

$$\begin{aligned} \lim_{k, l \rightarrow \infty} \|x_k - x_l\|^2 &= \lim_{k, l \rightarrow \infty} [\|x_k\|^2 + \|x_l\|^2 - 2 \operatorname{Re} \langle x_k, x_l \rangle] \\ &= 2 \left[\lim_{k \rightarrow \infty} \|x_k\|^2 - \lim_{k, l \rightarrow \infty} \operatorname{Re} \langle x_k, x_l \rangle \right] \\ &= 2 \left[\lim_{k \rightarrow \infty} \sum_{i,j} \bar{c}_i^k c_j^k \langle \rho(t_i^{k*} t_j^k)x, x \rangle \right. \\ &\quad \left. - \lim_{k, l \rightarrow \infty} \operatorname{Re} \sum_{i,j} \bar{c}_i^l c_j^k \langle \rho(t_i^{l*} t_j^k)x, x \rangle \right] \\ &= 2 \left[\lim_{k \rightarrow \infty} \phi_k(1) - \operatorname{Re} \lim_{k, l \rightarrow \infty} \sum_{i,j} \bar{c}_i^l c_j^k \phi(t_i^{l*} t_j^k) \right] = 0. \end{aligned}$$

Suppose that $\lim x_k = x_0 \in H$. Then

$$\psi(s) \equiv \lim \phi_k(s) = \lim \langle \rho(s)x_k, x_k \rangle = \langle \rho(s)x_0, x_0 \rangle.$$

If $\psi(s^*s) = 0$, then $\|\rho(s)x_0\|^2 = 0$, so that $\rho(s)x_0 = 0$. Hence, $\rho(s')x_0 = \rho(s)'x_0 = x_0$ and

$$\psi(s') = \langle \rho(s')x_0, x_0 \rangle = \|x_0\|^2 = \psi(1).$$

It follows that ϕ is strongly consistent. If ρ is a σ^* -representation, the results are similar.

(b) Let ϕ be a $*$ -positive function on S . Let

$$l_1(S) = \left\{ f: S \rightarrow C: \|f\| \equiv \sum_{s \neq 0} |f(s)| < \infty \right\}$$

where we identify $f, g \in l_1(S)$ if $f(s) = g(s)$ for all $s \neq 0$. Define addition and scalar multiplication pointwise in $l_1(S)$, define $f^*(s) = \overline{f(s^*)}$ for all $s \in S$, and define

$$(fg)(s) = \sum \{f(r)g(t): r, t \in S, rt = s\}.$$

Then $l_1(S)$ is a Banach $*$ -algebra [8], [10], [17] and every $f \in l_1(S)$ has a unique representation $f = \sum c_i \delta(s_i)$, $c_i \in C$, $[\delta(s)](t) \equiv \delta_{s,t}$ (the Kronecker delta). If we define $\omega: l_1(S) \rightarrow C$ by $\omega(f) = \sum c_i \phi(s_i)$, then ω is linear. Also, ω is positive since

$$\omega(f^*f) = \sum_{i,j} \bar{c}_i c_j \phi(s_i^* s_j) = \lim_{n \rightarrow \infty} \sum_{i,j=1}^n \bar{c}_i c_j \phi(s_i^* s_j) > 0.$$

By the GNS theorem [4] there exist a Hilbert space H and a cyclic $*$ -representation $\Pi: l_1(S) \rightarrow L(H)$ with cyclic vector x such that $\omega(f) = \langle \Pi(f)x, x \rangle$ for all $f \in l_1(S)$. Define $\rho: S \rightarrow L(H)$ by $\rho(s) = \Pi[\delta(s)]$. Then it is not difficult to show that ρ is a cyclic $*$ -representation of S in H with cyclic vector x , and $\phi(s) = \langle \rho(s)x, x \rangle$.

Now suppose that ϕ is additive. Then for $e \in P'(S)$ we have for every $t \in S$

$$\begin{aligned} \langle \rho(e)\rho(t)x, \rho(t)x \rangle &= \langle \rho(t^*et)x, x \rangle = \phi_t(e) \\ &= \phi_t(1) - \phi_t(e') = \langle [I - \rho(e')] \rho(t)x, \rho(t)x \rangle. \end{aligned}$$

Since $I - \rho(e) - \rho(e')$ is an orthogonal projection, it follows that

$$\|[I - \rho(e) - \rho(e')] \rho(t)x\|^2 = 0.$$

Hence, $[I - \rho(e) - \rho(e')] \rho(t)x = 0$. Since linear combinations of vectors $\rho(t)x$, $t \in S$, are dense in H , we have $\rho(e') = I - \rho(e) = \rho(e)'$. The proof that the σ -additivity of ϕ implies that ρ is a weak $*$ -representation is similar [14].

Now suppose that ϕ is strongly consistent. To prove (6), since $ss' = 0$, we have $\rho(s)\rho(s') = 0$. This shows that if $\rho(s')x_0 = x_0$ for $x_0 \in H$, then $\rho(s)x_0 = 0$. Conversely, suppose that $\rho(s)x_0 = 0$. Since x is cyclic, there exists a sequence $x_k = \sum_{i=1}^{n(k)} c_i^k \rho(t_i^k)x$ such that $\lim x_k = x_0$. For the sequence $\{c_i^k\} \subseteq C$, $\{t_i^k\} \subseteq S$, $k = 1, 2, \dots$, let $\phi_k = \phi\{c_i^k, t_i^k\}$. Since x_k is Cauchy, we have

$$\begin{aligned}
\lim_{k \rightarrow \infty} \phi_k(1) &= \lim_{k \rightarrow \infty} \sum_{i,j} \bar{c}_i^k c_j^k \phi(t_i^{k*} t_j^k) \\
&= \lim_{k \rightarrow \infty} \sum_{i,j} \bar{c}_i^k c_j^k \langle \rho(t_i^{k*} t_j^k) x, x \rangle \\
&= \lim_{k \rightarrow \infty} \|x_k\|^2 = \frac{1}{2} \left[\lim_{k, l \rightarrow \infty} \|x_k - x_l\|^2 + 2 \operatorname{Re} \lim_{k, l \rightarrow \infty} \langle x_k, x_l \rangle \right] \\
&= \operatorname{Re} \lim_{k, l \rightarrow \infty} \sum_{i,j} \bar{c}_i^l c_j^k \phi(t_i^{l*} t_j^k).
\end{aligned}$$

Hence, ϕ_k converges and

$$\psi(s) \equiv \lim \phi_k(s) = \lim \langle \rho(s) x_k, x_k \rangle = \langle \rho(s) x_0, x_0 \rangle.$$

Therefore, $\psi(s^*s) = \|\rho(s)x_0\|^2 = 0$. Since ϕ is strongly consistent,

$$\|\rho(s')x_0\|^2 = \langle \rho(s')x_0, x_0 \rangle = \psi(s') = \psi(1) = \|x_0\|^2.$$

Hence, $\rho(s')x_0 = x_0$. It follows that $\rho(s') = \rho(s)'$.

(c) Suppose that $\rho: S \rightarrow L(H)$ is an injective $*$ -representation. If $s \neq t$, then $\rho(s) \neq \rho(t)$. Hence, there exists a vector $x \in H$ such that $\langle \rho(s)x, x \rangle \neq \langle \rho(t)x, x \rangle$. It follows that the set of $*$ -positive functions $\{\langle \rho(\cdot)x, x \rangle: x \in H\}$ separates elements of S . If ρ is a $*$ -representation of a certain type, it follows from part (a) that the previous $*$ -positive function is of the corresponding type. Conversely, suppose there exists a separating set $\{\phi_\alpha\}$ of strongly consistent $*$ -positive functions on S . By part (b), there exists a cyclic Baer $*$ -representation $\rho_\alpha: S \rightarrow L(H_\alpha)$ corresponding to each ϕ_α . Let $H = \bigoplus H_\alpha$ and define $\rho: S \rightarrow L(H)$ by $[\rho(s)f](\alpha) = \rho_\alpha(s)f(\alpha)$ for every $f \in H$. It is not difficult to show that ρ is a Baer $*$ -representation of S in H . To show that ρ is injective, suppose that $s, t \in S$ and $s \neq t$. By hypothesis, there exists a ϕ_α such that $\phi_\alpha(s) \neq \phi_\alpha(t)$. Let $f \in H$ be defined by $f_\alpha(\beta) = x_\alpha \delta_{\alpha,\beta}$ where x_α is the cyclic vector for ρ_α . Then

$$\langle \rho(s)f, f \rangle = \langle \rho_\alpha(s)x_\alpha, x_\alpha \rangle = \phi_\alpha(s) \neq \phi_\alpha(t) = \langle \rho(t)f, f \rangle.$$

Hence, $\rho(s) \neq \rho(t)$ and ρ is injective. The proofs for the other types of $*$ -positive functions are similar. $\square$

The existence of Baer $*$ -representations for finite Baer $*$ -semigroups becomes much simpler. In this case one only needs a $*$ -positive function ϕ such that $\phi\{c_i, t_i\}$ is consistent for all finite sets $\{c_1, \dots, c_n\} \subseteq C, \{t_1, \dots, t_n\} \subseteq S$.

One can also consider the unitary equivalence of representations, subrepresentations, the irreducibility of representations, the unitary implementation of automorphisms, etc. [14]. In particular, it can be shown that the cyclic representation of S corresponding to a $*$ -positive function ϕ is unique to within unitary equivalence [14]. These results, along with their proofs, follow closely to those for C^* -algebras.

REFERENCES

1. A. Clifford and G. Preston, *The algebraic theory of semigroups*, Vol. 1, Math. Surveys, No. 7, Amer. Math. Soc., Providence, R. I., 1961.
2. E. B. Davies, *Quantum theory of open systems*, Academic Press, New York, 1976.
3. E. B. Davies and J. Lewis, *An operational approach to quantum probability*, Comm. Math. Phys. 17 (1970), 239–260.

4. G. Emch, *Algebraic methods in statistical mechanics and quantum field theory*, Interscience, New York, 1972.
5. D. Foulis, *Baer *-semigroups*, Proc. Amer. Math. Soc. **11** (1960), 648–654.
6. D. Foulis, *A note on orthomodular lattices*, Portugal. Math. **21** (1962), 65–72.
7. S. Gudder, *Stochastic methods in quantum mechanics*, Elsevier-North-Holland, New York, 1979.
8. S. Gudder and J. Michel, *Embedding quantum logics in Hilbert space*, Lett. Math. Phys. **3** (1979), 379–386.
9. R. Haag and D. Kastler, *An algebraic approach to quantum field theory*, J. Math. Phys. **5** (1964), 848–861.
10. E. Hewitt and H. Zuckerman, *The l_1 -algebra of a commutative semigroup*, Trans. Amer. Math. Soc. **83** (1956), 70–97.
11. J. Jauch, *Foundations of quantum mechanics*, Addison-Wesley, Reading, Mass., 1968.
12. R. Lindahl and P. Maserick, *Positive-definite functions on involution semigroups*, Duke Math. J. **39** (1971), 771–782.
13. G. Mackey, *The mathematical foundations of quantum mechanics*, Benjamin, New York, 1963.
14. J. Michel, *Hilbert space representations of Baer *-semigroups and quantum logics*, Dissertation, Univ. Wisconsin, 1974.
15. J. Pool, *Baer *-semigroups and the logic of quantum mechanics*, Comm. Math. Phys. **9** (1968), 118–141.
16. ———, *Semi-modularity and the logic of quantum mechanics*, Comm. Math. Phys. **9** (1968), 212–228.
17. G. Rütiman, *On the logical structure of quantum mechanics*, Found. Phys. **1** (1972), 173–182.
18. V. Varadarajan, *Geometry of quantum theory*, Vols. 1 and 2, Van Nostrand, Princeton, N. J., 1968 and 1970.

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, UNIVERSITY OF DENVER, DENVER, COLORADO 80208

DEPARTMENT OF MATHEMATICS, MARIETTA COLLEGE, MARIETTA, OHIO 45750