EXTREME VALUES FOR THE SIDON CONSTANT

DONALD I. CARTWRIGHT, ROBERT B. HOWLETT AND JOHN R. McMULLEN

ABSTRACT. Let G be a compact group and let $\phi \neq P \subseteq \hat{G}$. We consider the inequalities $1 < \kappa(P) < (\sum_{\sigma \in P} d_{\sigma}^2)^{1/2}$, where $\kappa(P)$ denotes the Sidon constant of P . The condition $\kappa(P) = 1$ essentially characterizes an example of Figà-Talamanca and Rider. The condition $\kappa(P) = (\sum_{\sigma \in P} d_{\sigma}^2)^{1/2}$ for finite P is equivalent to the existence of certain interesting functions on G . We show that $\kappa(\hat{G}) = |G|^{1/2}$ for a very large class of finite groups G , and this implies the existence of " G -circulant" unitary matrices whose entries all have modulus $|G|^{-1/2}$.

Let G be a compact group and let P be any nonempty subset of $\hat{G}$, the dual hypergroup of G . We denote by $\mathfrak{T}_P(G)$ the space of trigonometric polynomials f on G whose Fourier transforms $\hat{f}$ vanish off P . The point of departure of this article is the easily proved pair of inequalities

$$1 < \kappa(P) < \left(\sum_{\sigma \in P} d_{\sigma}^2 \right)^{1/2} \quad (1)$$

(see Proposition 1), in which $\kappa(P)$ denotes the Sidon constant of P , defined by

$$\kappa(P) = \sup \left\{ \|\hat{f}\|_1 = \sum_{\sigma \in P} d_{\sigma} \|\hat{f}(\sigma)\|_{\phi_1} : f \in \mathfrak{T}_P(G), \|f\|_{\infty} < 1 \right\},$$

and where, as usual, d_{σ} denotes the degree of the representation σ in $\hat{G}$, and $\|\cdot\|_{\phi_1}$ denotes the trace-class norm.

We investigate conditions under which $\kappa(P)$ takes one of the two bounding values in (1). For the case $\kappa(P) = 1$ we give a complete answer. Indeed, we show that the example of Figà-Talamanca and Rider [1] of a Sidon set P constructed using the projections on the various factors of a product of unitary groups gives essentially the only situation in which $\kappa(P) = 1$ can occur.

We give in Proposition 3 various necessary and sufficient conditions for $\kappa(P) = (\sum_{\sigma \in P} d_{\sigma}^2)^{1/2}$ to hold. In Theorem 2 and its corollary, we show that this is the case whenever $P = \hat{G}$ and G belongs to a large class of finite groups. This special case permits construction of certain generalized circulant Hadamard matrices, or equivalently certain elements of the group algebra CG . It also strongly generalizes the result of Graham [2], who established this when G is a finite abelian group.

Finally, we show by example that there exist two finite groups A and B , a hypergroup isomorphism $\phi: \hat{A} \rightarrow \hat{B}$, and a set $P \subseteq \hat{A}$ such that $\kappa(P) \neq \kappa(\phi(P))$. Thus the result of Vrem [7], that whether a subset P of $\hat{G}$ is a *central* Sidon set

Received by the editors March 17, 1980.

1980 *Mathematics Subject Classification*. Primary 43A46, 20C05, 05B20.

Key words and phrases. Sidon constant, Sidon set, circulant Hadamard matrix.

© 1981 American Mathematical Society
0002-9939/81/0000-0156/\$02.75

depends only on the hypergroup $\hat{G}$ and not on the compact group G giving rise to it, is unlikely to hold sway for general Sidon sets.

Unexplained notation will be that of Hewitt and Ross [3].

1. The basic inequalities. Henceforth when $P \subseteq \hat{G}$ we use the notation $\|P\| = \sum_{\sigma \in P} d_{\sigma}^2$. Then using [3, (D.51) and (28.43)] and the Cauchy-Schwarz inequality it is easy to obtain

PROPOSITION 1. *Let G be a compact group and let $P \subseteq \hat{G}$ be a finite nonempty set. Then for each $f \in \mathfrak{T}_P(G)$ we have*

$$\|f\|_{\infty} < \|f\|_1 < \|P\|^{1/2} \|f\|_{\infty}, \quad (2)$$

and hence

$$1 < \kappa(P) < \|P\|^{1/2}. \quad \square \quad (1')$$

2. Sets P with $\kappa(P) = 1$. Let $G = \prod_{\alpha \in A} G_{\alpha}$ be a Cartesian product in which each factor G_{α} is a unitary group. Let pr_{α} denote the projection on the factor G_{α} , and for each $\alpha \in A$ let $\sigma_{\alpha} \in \hat{G}_{\alpha}$ be the self-representation. In [1], Figà-Talamanca and Rider showed that $P = \{\sigma_{\alpha} \circ \text{pr}_{\alpha} | \alpha \in A\}$ is a Sidon set. In fact it is plain that $\kappa(P) = 1$. The following theorem characterizes situations of this kind.

THEOREM 1. *Let G be a compact group, and let P be any subset of $\hat{G}$. Let ϕ_P denote the homomorphism $\prod_{\sigma \in P} \sigma$ of G into $\mathfrak{U}_P = \prod_{\sigma \in P} \mathfrak{U}(H_{\sigma})$. Then $\kappa(P) = 1$ holds if and only if every element of $\mathfrak{U}_P$ can be written $\lambda \phi_P(x)$, with $\lambda \in \mathbb{T}$ and $x \in G$. In particular, if $P = \{1\} \cup Q$ with $1 \notin Q$, then the condition becomes: $\phi_Q: G \rightarrow \mathfrak{U}_Q$ is a surjection.*

The proof requires the following technical lemma.

LEMMA. *Let G be a compact group, let $P \subseteq \hat{G}$ be finite, and let $U_{\tau} \in \mathfrak{U}(H_{\tau})$ be given for each $\tau \in P$. Then there exist $x_1, \dots, x_n \in G$ and $\mu_1, \dots, \mu_n \in \mathbb{C}$ such that $\sum |\mu_k| = 1$ and such that $U_{\sigma} = \kappa(P) \sum_{k=1}^n \mu_k \sigma(x_k)$, for each $\sigma \in P$. Indeed $\kappa(P)$ is the smallest constant with this property.*

The lemma follows from the bipolar theorem applied to the inclusion $A \subseteq \kappa(P)B^{00}$, in which $B = \{\bigoplus_{\sigma \in P} \sigma(x) | x \in G\}$ and $A = \{\bigoplus_{\sigma \in P} U_{\sigma} | U_{\sigma} \in \mathfrak{U}(H_{\sigma}) \text{ for all } \sigma \in P\}$ are regarded as subsets of $\mathcal{L}(\bigoplus_{\sigma \in P} H_{\sigma})$, and the absolute bipolars are taken with respect to the pairing $\langle T, S \rangle = \text{trace}(TS)$. (Notice that the circled convex hull $\Gamma(B)$ of B is closed, since B is compact and $\mathcal{L}(\bigoplus_{\sigma \in P} H_{\sigma})$ is finite-dimensional.) $\square$

Now the assertion of the theorem for finite P follows immediately from the lemma, since $\mathfrak{U}(H)$ is the set of extreme points of the unit ball of $\mathcal{L}(H)$ (see [5, Theorem 1.6.4]). The case of infinite P then follows by a routine compactness argument. $\square$

REMARK. If $G = \mathbb{T}$, then Theorem 1 shows that a subset P of $\mathbb{Z}$ whose Sidon constant is 1 has at most two elements, since $\phi_P(G)$ has dimension at most one as a submanifold of $\mathfrak{U}_P$, while the latter has dimension $\text{card } P$. Similarly, if G is a

compact Lie group of dimension n and if $P \subseteq \hat{G}$ with $\kappa(P) = 1$, then P may contain at most $n + 1$ elements.

Before leaving this topic we record without proof a curious geometric interpretation of the Sidon constant afforded by the above lemma. We let $d(\cdot, \cdot)$ denote the metric obtained from the operator norm on $\mathcal{L}(\oplus_{\sigma \in P} H_\sigma)$, and define $\rho(P) = \sup\{d(U, \Gamma(B)) \mid U \in A\}$ where B , $\Gamma(B)$ and A are defined as in the proof of the lemma. Thus for a single $\sigma \in P$ we have

$$\rho(\sigma) = \sup\{d(U, \Gamma(\sigma(G))) \mid U \in \mathcal{U}(H_\sigma)\}.$$

PROPOSITION 2. $\kappa(P) = (1 - \rho(P))^{-1}$ for any finite $P \subseteq \hat{G}$. $\square$

3. Sets P with $\kappa(P) = \|P\|^{1/2}$.

PROPOSITION 3. Let G be a compact group and let $P \subseteq \hat{G}$ be nonempty and finite. Then $\kappa(P) = \|P\|^{1/2}$ holds if and only if there exists a complex-valued function f on G satisfying one of the following equivalent conditions:

- (a) $f \in \mathcal{T}_P(G)$, $\|\hat{f}\|_1 = \|P\|^{1/2}$ and $\|f\|_\infty \leq 1$.
- (b) $|f| \equiv 1$, and $\|P\|^{1/2}\hat{f}(\sigma)$ vanishes for $\sigma \notin P$ and is unitary for $\sigma \in P$.
- (c) $|f| \equiv 1$, and $f * f^* = \|P\|^{-1} \sum_{\sigma \in P} d_\sigma \chi_\sigma$.

PROOF. Since $\mathcal{T}_P(G)$ is finite-dimensional, the mapping $f \mapsto \|\hat{f}\|_1$ attains its maximum on $\{f \in \mathcal{T}_P(G) : \|f\|_\infty \leq 1\}$. Thus $\kappa(P) = \|P\|^{1/2}$ if and only if a function f exists satisfying (a). When proving Proposition 1 one sees that for equality to hold in the inequality $\|\hat{f}\|_1 \leq \|P\|^{1/2} \|f\|_\infty$ it is necessary and sufficient that $|f|$ be constant and that for some $c > 0$ independent of σ , $c\hat{f}(\sigma)$ is unitary for all $\sigma \in P$. This shows the equivalence of (a) and (b). The equivalence of (b) and (c) is easily seen by taking Fourier transforms. $\square$

It will be convenient to write δ_P for the function $\|P\|^{-1} \sum_{\sigma \in P} d_\sigma \chi_\sigma$ appearing in (c). In a special situation, δ_P assumes a simple form:

PROPOSITION 4. Let N be an open normal subgroup of the compact group G . Let $\rho \in \hat{N}$ be G -invariant (i.e. the representation $\rho^x: n \mapsto \rho(x^{-1}nx)$ is equivalent to ρ for each $x \in G$). Let $P = \rho^G$, where we define $\rho^G = \{\sigma \in \hat{G} : \rho \text{ is a component of } \sigma|_N\}$. Then P is finite, $\|P\| = [G : N]d_\rho^2$, and

$$\delta_P(x) = \begin{cases} 0 & \text{if } x \notin N, \\ \frac{\chi_\rho(x)}{d_\rho} & \text{if } x \in N. \end{cases}$$

PROOF. Put

$$g(x) = \begin{cases} 0 & \text{if } x \notin N, \\ \chi_\rho(x) & \text{if } x \in N. \end{cases}$$

If $\sigma \in P$ we see that $\hat{g}(\sigma) = [G : N]^{-1} \int_N \chi_\rho(n) \sigma(n^{-1}) dn$ so that by Clifford's theorem $\hat{g}(\sigma) = [G : N]^{-1} d_\rho^{-1} I_{H_\sigma}$. If $\sigma \notin P$ then ρ is not a component of $\sigma|_N$ and therefore $\hat{g}(\sigma) = 0$. From this we see that the equation $d_\rho [G : N] g = \sum_{\sigma \in P} d_\sigma \chi_\sigma$ holds, as both members are continuous functions and their Fourier transforms agree. Evaluation at the identity yields $[G : N] d_\rho^2 = \|P\|$ and hence $d_\rho \delta_P = g$ follows. $\square$

Our next theorem implies that $\kappa(P) = \|P\|^{1/2}$ holds for sets P of the form ρ^G , at least when G is a finite solvable group and $d_\rho = 1$.

THEOREM 2. *Let N be a normal subgroup of a finite solvable group G , and let ρ be a linear character of N satisfying $\rho(x^{-1}nx) = \rho(n)$ for all $x \in G$ and $n \in N$. Then there exists a function $f: G \rightarrow \mathbb{C}$ satisfying*

- (i) $|f(x)| = 1$ for all $x \in G$.
- (ii) $f(xn) = f(x)\rho(n)$ for all $x \in G$ and $n \in N$.
- (iii) $f * f^* = 0$ off N .

Note. (i) and (ii) entail $f * f^*|_N = \rho$.

PROOF. The key idea is lifting f from a subgroup M_1 to a subgroup M_3 , where $M_1 \triangleleft M_2 \triangleleft M_3$ and $[M_2 : M_1] = [M_3 : M_2] = p$ a prime. Straightforward lifting of f from M_1 to M_3 fails, in general.

Assuming the theorem is false, let G be a counterexample of minimal order. Choose $N \triangleleft G$ and a G -invariant linear character ρ of N for which no function f exists with the desired properties. We obtain a contradiction by means of several steps.

1. ρ is faithful. Since ρ is G -invariant, $H = \ker \rho$ is a normal subgroup of G . Let $\tilde{\rho}$ be the character of N/H induced by ρ . If $H \neq \{1\}$, then by minimality of G there exists $\tilde{f}: G/H \rightarrow \mathbb{C}$ satisfying (i), (ii) and (iii) with G replaced by G/H and N by N/H and ρ by $\tilde{\rho}$. Define $f: G \rightarrow \mathbb{C}$ by $f(x) = \tilde{f}(xH)$ and it is easy to see that f has the desired properties. This contradiction shows that $H = \{1\}$.

2. N is contained in the centre of G . This follows from step 1 since $\rho(x^{-1}nx) = \rho(n)$ for all $x \in G$ and $n \in N$.

3. There do not exist proper subgroups $G_1/N, \dots, G_r/N$ of G/N such that each $\xi \in G/N$ is uniquely expressible as a product $\xi_1 \xi_2 \cdots \xi_r$ with $\xi_j \in G_j/N$ for each j . Suppose G/N has such subgroups. Since $|G_j| < |G|$ for each j , there exist functions $f_j: G_j \rightarrow \mathbb{C}$ satisfying (i), (ii) and (iii) (with G replaced by G_j). Our hypotheses and step 2 show that each $x \in G$ may be expressed in the form $x_1 x_2 \cdots x_r$ (with $x_j \in G_j$ for all j) in exactly $|N|^{r-1}$ ways. Property (ii) of the functions f_j shows that a function $f: G \rightarrow \mathbb{C}$ is well defined if we set $f(x) = f_1(x_1)f_2(x_2) \cdots f_r(x_r)$ for any such decomposition of x . It is clear that f satisfies properties (i) and (ii). To see that (iii) holds, notice that $f = (|G|/|N|)^{r-1} \tilde{f}_1 * \cdots * \tilde{f}_r$, where

$$\tilde{f}_j(x) = \begin{cases} 0 & \text{if } x \in G \setminus G_j, \\ f_j(x) & \text{if } x \in G_j. \end{cases}$$

Notice also that $\tilde{f}_j * \tilde{f}_j^* = [G : G_j]^{-1} \delta$, that $\delta * \delta = [G : N]^{-1} \delta$, where

$$\delta(x) = \begin{cases} 0 & \text{if } x \in G \setminus N, \\ \rho(x) & \text{if } x \in N, \end{cases}$$

and that δ is central in $\mathfrak{T}(G)$ ($\delta = \delta_\rho$ in the notation of Proposition 4). This completes step 3.

4. G/N is a p -group for some prime p . If G/N is not a p -group, then being solvable it has nontrivial proper Hall subgroups G_1/N and G_2/N corresponding to

complementary sets of prime divisors of $|G/N|$. Since $G/N = G_1/N \cdot G_2/N$ and $G_1/N \cap G_2/N = \{1\}$, step 3 yields a contradiction.

5. If $|N| > 1$ then ρ does not extend to a character of G . Suppose that $|N| > 1$ and that ρ extends to a character $\tilde{\rho}$ of G . By minimality of G there exists a function $\tilde{f}: G/N \rightarrow \mathbb{C}$ satisfying (i), (ii) and (iii) (with G replaced by G/N , N by the trivial subgroup of G/N and of course ρ by the trivial representation). Define $f: G \rightarrow \mathbb{C}$ by $f(x) = \tilde{f}(xN)\tilde{\rho}(x)$ and clearly a contradiction is obtained.

6. G is not abelian. If G is abelian, then $N = \{1\}$ by step 5. But then the set P of Proposition 4 is the whole dual group $\hat{G}$ and so $\kappa(P) = \|P\|^{1/2} (= |G|^{1/2})$ (see Graham [2]). Hence a function f exists satisfying (i), (ii) and (iii).

7. For the p -group G/N we have $p = 2$. By steps 2 and 6 we have $N \subsetneq G$, and so there is a subgroup M of G with $N \subset M$, $M/N \subseteq Z(G/N)$ and $[M : N] = p$. Then M is abelian and normal. Let $\Lambda = \{\lambda \in \hat{M} : \lambda|_N = \rho\}$, and fix any $\lambda_0 \in \Lambda$. Then $\Lambda = \{\lambda_0 \nu : \nu \in N^0\}$, where N^0 denotes the subgroup of $\hat{M}$ whose elements are trivial on N . Now G acts on Λ and fixes N^0 elementwise. Let H_0 be the stabilizer of λ_0 . Since $|\Lambda| = p$ we must have $H_0 = G$ or $[G : H_0] = p$. Since M is abelian, we have $M \subseteq H_0$ and $M \neq G$. Hence there is a subgroup H of G with $[G : H] = p$ and $M \subseteq H \subseteq H_0$. Since $|H| < |G|$ there is for each $\lambda \in \Lambda$ a function $f_\lambda: H \rightarrow \mathbb{C}$ satisfying (i), (ii) and (iii) with G replaced by H , N replaced by M , and ρ replaced by λ . Choose coset representatives $\{x_\lambda : \lambda \in \Lambda\}$ for H in G (note that $|\Lambda| = [G : H] = p$) and define $f: G \rightarrow \mathbb{C}$ by $f(x_\lambda h) = f_\lambda(h)$ for $\lambda \in \Lambda$ and $h \in H$. Clearly f satisfies (i) and (ii), and one calculates that

$$(f * f^*)(x) = \begin{cases} 0 & \text{if } x \notin M, \\ \frac{1}{p} \sum_{\lambda \in \Lambda} \lambda(x_\lambda^{-1} x x_\lambda) & \text{if } x \in M. \end{cases} \quad (3)$$

(Note that $\lambda^H \cap \lambda'^H = \emptyset$ when $\lambda, \lambda' \in \Lambda$ are distinct, since λ is the only component of $\sigma|_M$ if $\sigma \in \lambda^H$. Therefore $f_\lambda * f_{\lambda'}^* = 0$ if $\lambda \neq \lambda'$, taking the convolution over H here, since $f_\lambda \in \mathfrak{T}_Q(H)$ with $Q = \lambda^H$.)

If the coset representatives x_λ are chosen so that the characters $\{\lambda^{x_\lambda} : \lambda \in \Lambda\}$ are distinct, then the sum in (3) is simply $\sum_{\lambda \in \Lambda} \lambda(x)$ and so f satisfies (iii). Such a choice of x_λ 's is achieved as follows: Suppose first that $H_0 \neq G$. Pick $x \in G \setminus H_0$. Thus $\lambda_0^x = \lambda_0 \nu$ for some nontrivial $\nu \in N^0$. The powers ν^k of ν for $k = 0, \dots, p-1$ are distinct, and so $\lambda_0 \nu^k$ runs through all of Λ . For each k we have $(\lambda_0 \nu^k)^{x^k} = \lambda_0 \nu^{2k}$, and if $p \neq 2$ the characters ν^{2k} , $k = 0, \dots, p-1$, are all distinct. Hence if we set $x_\lambda = x^k$ whenever $\lambda = \lambda_0 \nu^k$, we have achieved our aim. In the case $H_0 = G$, we have $\lambda^x = \lambda$ for all $x \in G$ and $\lambda \in \Lambda$, and any choice of x_λ 's will do. Hence $p \neq 2$ leads to a contradiction.

8. G/N is not a 2-group. The argument used in step 7 breaks down if the two extensions λ_1 and λ_2 of ρ to M are interchanged by some $x \in G$. Assume then that this is the case, so that $H = H_0$ in the above notation. If H is the only maximal subgroup of G containing M , then G/M is cyclic, and since $M/N \subseteq Z(G/N)$, G/N is abelian. But then G/N is cyclic by step 3 and the structure theory of abelian groups. But this and $N \subseteq Z(G)$ shows that G is abelian, contradicting step 6. So let K be another maximal subgroup of G containing M , and let $L = H \cap K$.

Pick any $x \in K \setminus H$. Then $K = L \cup Lx$, and x interchanges λ_1 and λ_2 . Let $g_1, g_2: L \rightarrow \mathbb{C}$ satisfy (i), (ii) and (iii) with L replacing G , M replacing N , and λ_1 (resp. λ_2) replacing ρ . Define $f_1, f_2: K \rightarrow \mathbb{C}$ by $f_1(l) = g_1(l)$ for $l \in L$, $f_1(lx) = g_1(l)$, $f_2(l) = g_2(l)$, and $f_2(lx^{-1}) = -g_2(l)$. Then f_1 and f_2 satisfy (i) and (iii) (but not (ii)) with K replacing G , M replacing N and λ_1 (resp. λ_2) replacing ρ . Also $f_1 * f_2^* = 0$, taking the convolution over K . The calculations needed here are not difficult. For example, if $\xi \in Lx$, write $\xi = l_0x$ with $l_0 \in L$, and we have

$$\begin{aligned} & \frac{1}{|K|} \sum_{k \in K} f_1(k) \overline{f_1(\xi^{-1}k)} \\ &= \frac{1}{|K|} \sum_{l \in L} \{ f_1(l) \overline{f_1(x^{-1}l_0^{-1}l)} + f_1(lx) \overline{f_1(x^{-1}l_0^{-1}lx)} \} \\ &= \frac{1}{|K|} \sum_{l \in L} \{ g_1(l) \overline{g_1(x^{-1}l_0^{-1}lx^{-1})} + g_1(l) \overline{g_1(x^{-1}l_0^{-1}lx)} \} \\ &= \frac{1}{|M||K|} \sum_{l \in L, m \in M} \{ g_1(lm) \overline{g_1(x^{-1}l_0^{-1}lmx^{-1})} + g_1(lm) \overline{g_1(x^{-1}l_0^{-1}lmx)} \} = 0, \end{aligned}$$

using the property (ii) for g_1 and the fact that x interchanges λ_1 and λ_2 .

Now pick any $y \in H \setminus K$. Then $G = K \cup yK$, and we may define $f: G \rightarrow \mathbb{C}$ by $f(k) = f_1(k)$ and $f(yk) = f_2(k)$ if $k \in K$. It is routine to check that f satisfies properties (i), (ii) and (iii). $\square$

COROLLARY. Let G be a finite group having solvable subgroups $G_1, \dots, G_r$ ($r > 1$) so that each $x \in G$ is uniquely expressible as a product $x_1 \cdots x_r$ with $x_j \in G_j$ for each j . Then $\kappa(\hat{G}) = \|\hat{G}\|^{1/2} = |G|^{1/2}$.

PROOF. When G is solvable this follows from Theorem 2 above, taking N to be the trivial subgroup. The general case follows from the solvable case as in step 3 of the proof. $\square$

REMARK. We conjecture that every finite group can be decomposed in the above manner. If this conjecture is true, then $\kappa(\hat{G}) = |G|^{1/2}$ and Theorem 2 are valid for every finite group G .

4. Circulant matrices. Let G be a finite group. We will say that a matrix $(\xi_{ij})_{i,j=1}^n$ is G -circulant if there is an enumeration $x_1 = e, x_2, \dots, x_n$ of G and a function $h: G \rightarrow \mathbb{C}$ such that $\xi_{ij} = h(x_i^{-1}x_j)$ ($1 \leq i, j \leq n$). If $G = (a^k)_{k=0}^{n-1}$ is cyclic and if $x_j = a^{j-1}$ for $j = 1, 2, \dots, n$ and if we put $m_j = \xi_{1j}$, then such a matrix takes the form

$$M = \begin{pmatrix} m_1 & m_2 & \cdot & \cdot & \cdot & \cdot & m_n \\ m_n & m_1 & m_2 & \cdot & \cdot & \cdot & m_{n-1} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ m_2 & m_3 & \cdot & \cdot & \cdot & m_n & m_1 \end{pmatrix},$$

and is circulant in the usual sense.

THEOREM 3. *Let G be any finite group of order n . Then there exists a unitary matrix which is G -circulant, and whose entries all have modulus $n^{-1/2}$, if and only if $\kappa(\hat{G}) = n^{1/2}$.*

PROOF. Apply Propositions 3 and 4 with N the trivial subgroup of G . If f is a function satisfying (c) of Proposition 3, then the matrix (ξ_{ij}) with $\xi_{ij} = n^{-1/2}f(x_i^{-1}x_j)$ has the desired properties. $\square$

The proof of Theorem 2 shows that the matrix in Theorem 3 can be chosen to have the form $A/\sqrt{n}$, where the entries of A are roots of unity. For cyclic G , this matrix is real-valued if and only if A is an $n \times n$ circulant Hadamard matrix. Such matrices A are known not to exist for $5 < n < 12100$. For A to have ± 1 , $\pm i$ as entries, n is still very restricted (see, e.g. [6]).

5. A small example. Let D and Q denote respectively the dihedral and quaternion groups of order 8, presented by $D = \langle a, b | a^4 = 1 = b^2, b^{-1}ab = a^{-1} \rangle$ and $Q = \langle a, b | a^4 = 1, b^2 = a^2, b^{-1}ab = a^{-1} \rangle$. Then $\hat{D}$ and $\hat{Q}$ are isomorphic as hypergroups (see McMullen and Price [4]), each consisting of four linear representations, and a two-dimensional representation. Denote the two-dimensional representation of D by σ_1 and that of Q by σ_2 . Then explicit calculations reveal that $\kappa(\{1, \sigma_1\}) = 2$ but $\kappa(\{1, \sigma_2\}) = \sqrt{5}$. The set $P_2 = \{1, \sigma_2\} \subseteq \hat{Q}$ therefore satisfies the equality $\kappa(P) = \|P\|^{1/2}$, while $P_1 = \{1, \sigma_1\} \subseteq \hat{D}$ does not. Thus the Sidon constant is not preserved by hypergroup isomorphisms. We also note that P_2 is not of the special form ρ^G introduced in Proposition 4.

REFERENCES

1. A. Figà-Talamanca and D. Rider, *A theorem of Littlewood and lacunary series for compact groups*, Pacific J. Math **16** (1966), 505–514. MR **34** #6444.
2. C. C. Graham, *The Sidon constant of a finite abelian group*, Proc. Amer. Math. Soc. **68** (1978), 83–84. MR **56** #16262.
3. E. Hewitt and K. A. Ross, *Abstract harmonic analysis*, II, Springer-Verlag, Berlin and New York, 1970.
4. J. R. McMullen and J. F. Price, *Reversible hypergroups*, Rend. Sem. Mat. Fis. Milano **47** (1977), 67–85.
5. S. Sakai, *C*-algebras and W*-algebras*, Ergebnisse der Math. und ihrer Grenzgebiete, Band 60, Springer-Verlag, Berlin and New York, 1971.
6. R. J. Turyn, *Complex Hadamard matrices*, Combinatorial Structures and their Applications (Proc. Calgary Internat. Conf., Calgary, Alberta, 1969), Gordon and Breach, New York, 1970, pp. 435–437. MR **42** #5821.
7. R. C. Vrem, *Lacunarity on compact hypergroups*, Math. Z. **164** (1978), 93–104.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SYDNEY, SYDNEY, N.S.W. 2006, AUSTRALIA