

MULTIPLES OF WEIERSTRASS POINTS AS SPECIAL DIVISORS

R. F. LAX

ABSTRACT. Complex spaces \mathcal{W}_n^r of Weierstrass points are isomorphic to the intersection, on the n th symmetric product of the universal curve over the Teichmüller space, of complex spaces \mathcal{G}_n^r of special divisors with the diagonal Δ_n consisting of divisors which are multiples of a point. The tangent space at a point of this intersection is described and it is shown that $\mathcal{G}_n^1 - \mathcal{G}_n^2$ and Δ_n intersect transversally.

Let $T = T_g$ denote the Teichmüller space for Teichmüller surfaces of genus $g > 1$ and let $\pi: V \rightarrow T$ denote the universal curve of genus g . Denote by $V_T^{(n)}$ the n th symmetric product of V over T . Let \mathcal{G}_n^r denote the closed complex subspace of $V_T^{(n)}$ whose points are divisors of degree n and projective dimension at least r (see [3], [2]). We have proved

THEOREM 1 ([3]). *Suppose $n \leq g$. Then $\mathcal{G}_n^1 - \mathcal{G}_n^2$ is smooth of pure dimension $2n + 2g - 4$.*

For $2 \leq n \leq g$, let \mathcal{W}_n^r denote the closed complex subspace of V consisting of those $(t, P) \in V$ such that there are at least r gaps less than or equal to n in the Weierstrass gap sequence at P on V_t . These spaces were introduced in [4] and, by employing methods similar to those used in the proof of Theorem 1, we proved

THEOREM 2 ([4]). *For $2 \leq n \leq g$, $\mathcal{W}_n^1 - \mathcal{W}_n^2$ is smooth of pure dimension $n + 2g - 3$.*

In this note, we describe the relationship between the \mathcal{G}_n^r and the \mathcal{W}_n^r and show how Theorem 2 may be derived in a direct fashion from Theorem 1.

Let Δ_n denote the image of $\delta_n: V \rightarrow V_T^{(n)}$, the closed immersion which takes a point (t, P) to the point (t, nP) . The following proposition follows easily from the definitions.

PROPOSITION 1. *For $2 \leq n \leq g$,*

$$\delta_n|_{\mathcal{W}_n^r}: \mathcal{W}_n^r \xrightarrow{\cong} \mathcal{G}_n^r \cap \Delta_n.$$

We now explicitly consider the intersection of \mathcal{G}_n^r and Δ_n . Suppose $(t, nP) \in \mathcal{G}_n^r \cap \Delta_n$. Put $X = V_t$. Let z be a local coordinate on X centered at P and let z_1, \dots, z_n denote n copies of z . Let $\sigma_1, \dots, \sigma_n$ denote the n elementary symmetric functions in z_1, \dots, z_n . Let c_1, \dots, c_{3g-3} denote Patt's local coordinates on T

Received by the editors May 8, 1980.

1980 *Mathematics Subject Classification.* Primary 32G15, 14H15.

Key words and phrases. Complex space, Weierstrass point, special divisor, Teichmüller space, universal curve.

centered at t (see [8]). Then $c_1, \dots, c_{3g-3}, \sigma_1, \dots, \sigma_n$ are local coordinates on $V_T^{(n)}$ centered at (t, nP) . Put

Z = tangent space to $V_T^{(n)}$ at (t, nP) ;

Z_1 = tangent space to \mathcal{G}_n^r at (t, nP) ;

Z_2 = tangent space to Δ_n at (t, nP) .

We describe coordinates for Z . Suppose $\xi \in Z$. We may view ξ as a \mathbb{C} -homomorphism of local rings

$$\xi: \mathcal{O}_{V_T^{(n)}, (t, nP)} \rightarrow \mathbb{C}[\varepsilon]/(\varepsilon^2)$$

(cf. [6, p. 332]). Then ξ is determined by its values on a set of local parameters of $\mathcal{O}_{V_T^{(n)}, (t, nP)}$. So, if $\xi(c_m) = b_m \varepsilon$, $m = 1, \dots, 3g-3$, and $\xi(\sigma_i) = u_i \varepsilon$, $i = 1, \dots, n$, then $(u_1, \dots, u_n, b_1, \dots, b_{3g-3})$ serve as coordinates for Z .

PROPOSITION 2. $\xi = (u_1, \dots, u_n, b_1, \dots, b_{3g-3})$ is in Z_2 if and only if $u_2 = u_3 = \dots = u_n = 0$.

PROOF. Suppose $(t, Q_1 + \dots + Q_n) \in V_T^{(n)}$ is a point near (t, nP) . Then $(t, Q_1 + \dots + Q_n) \in \Delta_n \Leftrightarrow z(Q_1) = \dots = z(Q_n) = z_0 \Leftrightarrow z_0$ is an n -fold root of

$$F(Y) = \prod_{i=1}^n (Y - z(Q_i)) = Y^n - \sigma_1(z(Q_1), \dots, z(Q_n)) Y^{n-1}$$

$$+ \dots + (-1)^n \sigma_n(z(Q_1), \dots, z(Q_n))$$

$$\Leftrightarrow F(z_0) = F'(z_0) = \dots = F^{(n-1)}(z_0) = 0$$

$$\Leftrightarrow \sigma_k(z(Q_1), \dots, z(Q_n)) = \binom{n}{k} [\sigma_1(z(Q_1), \dots, z(Q_n))]^k / n^k$$

$$\text{for } k = 2, 3, \dots, n,$$

and $\sigma_1(z(Q_1), \dots, z(Q_n)) = nz_0$.

So, near (t, nP) , Δ_n is defined by the equations $\{\sigma_k = \binom{n}{k} \sigma_1^k / n^k\}$, $k = 2, \dots, n$. Thus ξ is tangent to Δ_n at (t, nP) if and only if $\xi(\sigma_k) = 0$ for $k = 2, 3, \dots, n$.

We next recall the description of Z_1 , which was given in [3]. Let $1, \gamma_2, \dots, \gamma_g$ denote the Weierstrass gaps at $P \in X$. Choose a basis of holomorphic 1-forms $d\xi_1, \dots, d\xi_g$ on X such that $\text{ord}_P d\xi_j = \gamma_j - 1$. Write

$$d\xi_j = \sum_{i=0}^{\infty} a_{i,j} z^i dz.$$

For details concerning the following result, we refer the reader to [3].

PROPOSITION 3. Suppose $n < g$ and $\xi \in Z$. Then $\xi \in Z_1$ if and only if all minors of order $n - r + 1$ of the matrix

$$\mathfrak{M} = \left[\begin{array}{c|c} (-1)^i a_{i,j} & \varepsilon \left[\sum_{l=1}^n (-1)^{i+l-1} a_{i+l} u_l + \sum_{m=1}^{3g-3} \tau'_{P,i}(Q_m) \xi'_j(Q_m) b_m \right] \right] \\ \hline i = 0, \dots, n-1 & i = 0, \dots, n-1 \\ j = 1, \dots, n-r & j = n-r+1, \dots, g \end{array} \right]$$

vanish, where $\tau_{P,k}$ is an elementary integral of the second kind on X with pole of order $k+1$ at P and where (Q_1, \dots, Q_{3g-3}) is any point chosen from an open subset of X^{3g-3} .

Now, suppose $(t, nP) \in \mathcal{G}_n^r - \mathcal{G}_n^{r+1}$, $n < g$. Then \mathcal{N} will have a nonzero minor of order $n - r$, call it μ , and in order that all minors of order $n - r + 1$ of \mathcal{N} vanish, it is sufficient that those minors of order $n - r + 1$ which contain μ should vanish. This gives rise to $r(g - n + r)$ linear equations $\{E_k\}$ in $u_1, \dots, u_n, b_1, \dots, b_{3g-3}$. These equations are of the form

$$E_k: \sum_{l=1}^n e_{k,l} u_l + \sum_{m=1}^{3g-3} \alpha_k(Q_m) b_m = 0,$$

where the α_k are (not necessarily finite) quadratic differentials on X . (The α_k arise from the products $d\tau_{P,i} d\zeta_j$ which appear in \mathcal{N} —see [3].)

THEOREM 3. *Suppose $n < g$, $r(g - n + r) < 3g - 3$, and $(t, nP) \in \mathcal{G}_n^r - \mathcal{G}_n^{r+1}$. If the above α_k , $k = 1, \dots, r(g - n + r)$, are linearly independent quadratic differentials, then:*

- 1) $\dim Z_1 = 3g - 3 + (r + 1)(n - r) - rg + r$ and \mathcal{G}_n^r is smooth at (t, nP) .
- 2) $\dim Z_1 \cap Z_2 = 3g - 2 - r(g - n + r)$ and \mathcal{G}_n^r and Δ_n intersect transversally at (t, nP) .

PROOF. One may show, as in [3], that if the α_k are linearly independent, then since (Q_1, \dots, Q_{3g-3}) is any point from an open subset of X^{3g-3} , the matrix $[\alpha_k(Q_m)]$, $k = 1, \dots, r(g - n + r)$ and $m = 1, \dots, 3g - 3$, will have maximum rank. It then follows that the systems of equations which define Z_1 and $Z_1 \cap Z_2$ will have maximum rank, establishing the theorem.

We showed in [3] that at least $g - n + r$ of the α_k are linearly independent. In particular, if $r = 1$, then all the α_k are linearly independent. As a consequence we have

THEOREM 4. *Suppose $(t, nP) \in \mathcal{G}_n^r - \mathcal{G}_n^{r+1}$, $n < g$. Then*

- (1) $\dim Z_1 \leq 2g + 2n - r - 3$; in particular, if $r = 1$, then $\dim Z_1 = 2g + 2n - 4$ and \mathcal{G}_n^1 is smooth at (t, nP) .
- (2) $\dim Z_1 \cap Z_2 \leq 2g + n - r - 2$; in particular, if $r = 1$, then $\dim Z_1 \cap Z_2 = 2g + n - 3$ and \mathcal{G}_n^1 and Δ_n intersect transversally at (t, nP) .

COROLLARY. *For $n < g$,*

- (1) $\dim \mathcal{W}_n^r \leq 2g + n - r - 2$;
- (2) $\mathcal{W}_n^1 - \mathcal{W}_n^2$ is smooth of pure dimension $2g + n - 3$.

REMARKS. (1) The smoothness of $\mathcal{G}_n^1 - \mathcal{G}_n^2$ has also recently been demonstrated by Arbarello-Cornalba [1] and Namba [7].

(2) Arbarello-Cornalba [1] have shown that $\mathcal{G}_n^2 - \mathcal{G}_n^3$ is smooth, but it does not necessarily follow that the α_k are then linearly independent or that this space intersects Δ_n transversally.

(3) In [5], we defined \mathcal{W}_n^r for $n > g$. The points of this space are those $(t, P) \in V$ such that there are at least r gaps greater than n in the gap sequence at $P \in V_t$. We showed that for $n > g$, $\mathcal{W}_n^1 - \mathcal{W}_n^2$ is smooth of pure dimension $4g - n - 3$. This result can also be obtained as above by considering the intersection of \mathcal{G}_n^r and Δ_n for $n > g$, but we note that, by our definition of \mathcal{W}_n^r , for $n > g$, $\mathcal{W}_n^r = \delta_n^{-1}(\mathcal{G}_n^{n-g+r})$.

REFERENCES

1. E. Arbarello and M. Cornalba, *Su una congettura di Petri*, Comment Math. Helv. (to appear).
2. R. C. Gunning, *Lectures on Riemann surfaces: Jacobi varieties*, Princeton Univ. Press, Princeton, N. J., 1972.
3. R. F. Lax, *On the dimension of varieties of special divisors*, Trans. Amer. Math. Soc. **203** (1975), 141–159.
4. ———, *Weierstrass points of the universal curve*, Math. Ann. **216** (1975), 35–42.
5. ———, *Gap sequences and moduli in genus 4*, Math. Z. **175** (1980), 67–75.
6. D. Mumford, *Introduction to algebraic geometry* (preliminary version of first 3 chapters), Harvard notes.
7. M. Namba, *Families of meromorphic functions on compact Riemann surfaces*, Lecture Notes in Math., vol. 767, Springer-Verlag, Berlin, 1979.
8. C. Patt, *Variations of Teichmüller and Torelli surfaces*, J. Analyse Math. **11** (1963), 221–247.

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MASSACHUSETTS 02115

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA (Current address)