COMMENTS ON NOTES OF STANOJEVIĆ ET AL.

MAHLON M. DAY

ABSTRACT. Some further inequalities which characterize inner-product spaces among normed linear spaces are derived by averaging characters of locally compact groups.

1. Comparing results in Penico and Stanojević (Proc. Amer. Math. Soc. **79** (1980), 427–430) with the Schoenberg characterization:

$(S, \sim)$ For all x, y in S , $2^{-1}(\|x + y\|^2 + \|x - y\|^2) \sim 2$, in words:

the average of $\|x + zy\|^2$ over the group $\{1, -1\} \sim 2$,

of inner-product spaces (see Day, *Normed linear spaces*, Chapter VII, §3, for a general view of such characterizations) suggested that a modification of the proof of the Dvoretzky-Rogers Theorem (*op. cit.*, p. 80, Lemma 1) would prove a family of “averaging” characterizations of inner-product spaces. For notation, Γ is the circle group in the complex plane, S is the unit sphere $\{x \mid \|x\| = 1\}$ in a normed space N , and $\sim$ means one of the relations $<$, $=$, or $>$ but the same one for all x and y in S .

THEOREM A. Let $G \neq 1$ be a closed subgroup of the circle group Γ and let H be Haar measure on G normalized so that $H(G) = 1$. Let

$$\Psi_G(x, y) = \int_G \|x + zy\|^2 dH(z).$$

Then a complex-linear normed space N is an inner-product space if and only if

$$(\Psi_G, \sim) \quad \text{for all } x, y \text{ in } S, \Psi_G(x, y) \sim 2.$$

The note of Stanojević and Suchanek (Proc. Amer. Math. Soc. **81** (1981), 101–103) gives a more general result. Their paper does not, however, suggest the finite examples which are prominent in Theorem A because the only closed proper subgroups of Γ are the groups G_k of k th roots of unity. If θ is a principal k th root of unity, then N is an inner-product space if and only if for each (or for one) $k > 2$

$$(\Psi_k, \sim) \quad \text{for each } x \text{ and } y \text{ in } S, k^{-1} \sum_{p < k} \|x + \theta^p y\|^2 \sim 2.$$

The case $k = 2$ is $(S, \sim)$. The case $k = 3$ already gives an unfamiliar condition

$$(\|x + y\|^2 + \|x + \omega y\|^2 + \|x + \omega^2 y\|^2)/3 \sim 2,$$

where, of course, $\omega^3 = 1$.

Received by the editors April 23, 1980.

AMS (MOS) subject classifications (1970). Primary 46C10.

© 1981 American Mathematical Society
0002-9939/81/0000-0220/\$01.50

A result similar to Theorem A gives an averaging analog of the condition of Penico and Stanojević; replace Ψ_G above by

$$\Phi_G(x, y) = \int_G \|(\operatorname{Re}(z))x + (\operatorname{Im}(z))y\|^2 dH(z).$$

One can make a corresponding analog of the theorem of Stanojević and Suchanek by replacing their integrand by $\|(\operatorname{Re}(\gamma(g)))x + (\operatorname{Im}(\gamma(g)))y\|^2$.

2. Von Neumann (Trans. Amer. Math. Soc. **36** (1934), p. 445) pointed out that the space $AP(G)$ of almost periodic functions on an arbitrary group G is a natural generalization of the space of continuous functions on a compact group. One aspect of this is the existence on $AP(G)$ of a unique invariant mean m , which specializes to a normalized Haar measure when G is compact. Since each character of G is almost periodic, this point of view allows a generalization of the theorem of Stanojević and Suchanek sufficiently broad to allow as a corollary the generalization of Theorem A to all subgroups of Γ .

THEOREM B. *A complex-linear space N is an inner-product space if there exists a topological group G and a continuous character $\gamma \not\equiv 1$ of G such that*

$$(\Psi_\gamma, \sim) \quad \text{for all } x, y \text{ in } S, \quad \Psi_\gamma(x, y) = m_g(\|x + \gamma(g)y\|^2) \sim 2.$$

Conversely, if N is an inner-product space, $(\Psi_\gamma, =)$ holds for all G and all $\gamma \not\equiv 1$.

The proof is that of Theorem A or of the theorem of Stanojević and Suchanek.

Since the invariant mean for the almost periodic functions on the integers or on the reals can be calculated as a limit of averages, special cases of Theorem B say that N is an inner-product space if and only if for each (or for one) $z \neq 1$ in Γ and for all x, y in S

$$(i) \quad \lim_{K \rightarrow \infty} K^{-1} \sum_{p < K} \|x + z^p y\|^2 \sim 2,$$

or

$$(ii) \quad \lim_{K \rightarrow \infty} (2K)^{-1} \int_{-K}^K \|x + \exp(it)y\|^2 dt \sim 2.$$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, ILLINOIS 61801