ON THE DINI TEST AND DIVERGENCE OF FOURIER SERIES

CALIXTO P. CALDERON¹

ABSTRACT. In this note we prove that no condition weaker than the Dini assures the pointwise convergence of a Fourier series in a set of positive measure.

1. Statement of results. It is well known that if a function f belonging to $L^1[0, 2\pi]$ satisfies at a point x the condition

$$\int_{-\delta}^{\delta} \frac{|f(x) - f(x-t)|}{|t|} dt < \infty \quad (1.1)$$

then the partial sums S_m of the Fourier series of f converge to $f(x)$.

The above condition is optimal for individual points. In fact, "Given any continuous $\mu(t) \geq 0$ such that $\mu(t)/t$ is not integrable in a neighborhood of $t = 0$, we can find a continuous function f , such that $|f(t) - f(0)| < \mu(t)$ for small t and $S_m(f)$ diverges at $t = 0$." (See [5, Vol. I, Theorem 2.4, p. 303].) On the other hand, if a periodic function f satisfies the conditions $|f(x+t) - f(x)| < \mu(t)$, $0 < t < \delta(x)$, x in a set E of positive measure, then $S_m(f)$ converges at almost every x in E . This fact is an easy consequence of Carleson's theorem (see [1]).

The purpose of this note is to show that the condition (1.1) is the optimal one for x belonging to a set E of positive measure, if we give a suitable definition of what we understand by a weaker condition. More precisely, for each $w(t)$ that satisfies

$$\begin{aligned} \text{(i)} \quad & w(t) \text{ increasing and continuous, } w(0) = 0, \\ \text{(ii)} \quad & \int_0^{\delta} w(t) dt/t = \infty, \end{aligned} \quad (1.2)$$

we have

2. THEOREM. *There is a function g belonging to $L^1[0, 2\pi]$ and a set F of positive measure such that*

$$(\alpha) \int_{-\delta}^{\delta} |g(x+t) - g(x)| w(|t|)/|t| dt < \infty, x \in F.$$

($\alpha\alpha$) The partial sums $S_n(g)$ of the Fourier series of g diverge a.e. in F .

PROOF. First, we invoke the following theorem due to Marcinkiewicz (see [3]):

THEOREM 2.1. *Let $\varphi(t)$ be a continuous function, increasing and such that $\varphi(0) = 0$. Assume in addition that*

$$(\text{i}) \quad [\varphi(t)]^{-1} = o(\log 1/t), t \rightarrow 0.$$

Then, there is a function $f \in L^1[0, 2\pi]$ such that

$$(\beta) \quad (1/|h|) \int_0^h |f(x+t) - f(x)| dt = O[\varphi(|h|)] \text{ a.e.};$$

($\beta\beta$) The partial sums $S_n(f)$ of the Fourier series of f diverge a.e.

Received by the editors July 23, 1980.

1980 *Mathematics Subject Classification.* Primary 42A20.

¹Research supported by an N.S.F. grant.

© 1981 American Mathematical Society
0002-9939/81/0000-0314/\$01.75

Secondly, we need a lemma of very well-known type.

LEMMA 2.2. Let $\varphi(t)$ be defined by

$$\varphi(t) = \left[\int_t^1 w(s) \frac{ds}{s} \right]^{-1}, \quad 0 < t < 1,$$

and $w(s)$ as in (1.2). Suppose that a function $f \in L^1[0, 2\pi]$ satisfies

(i) $(1/|h|) \int_0^h |f(x+t) - f(x)| dt = O[\varphi(|h|)]$ a.e. in $[0, 2\pi]$.

Then, for each $\varepsilon > 0$ there are a perfect subset F of $[0, 2\pi]$ and a constant C such that

(α) $|F| > 2\pi - \varepsilon$,

($\alpha\alpha$) $|f(x_1) - f(x_2)| < C\varphi(|x_1 - x_2|)$ whenever $x_1, x_2 \in F$ and $0 < |x_1 - x_2| < 1/2$,

($\alpha\alpha\alpha$) $(1/|h|) \int_0^h |f(x) - f(x+t)| dt < C\varphi(|h|)$ whenever $x \in F$.

The proof follows verbatim the lines of the corresponding result in [5, Vol. II, p. 171]. (There, $\varphi(t) = 1/|\log t|$.) One also has to use the fact

$$\varphi(ct) < k\varphi(t), \quad c > 1, \quad 0 < t < 1/4C, \quad (2.2.1)$$

which is an easy consequence of the particular form that $\varphi(t)$ has.

3. Final steps. Consider

$$\varphi(t) = \left[\int_t^1 \bar{w}(s) \frac{ds}{s} \right]^{-1}, \quad 0 < t < 1/2,$$

where $\bar{w}(s) = \max(|\log s|^{-\delta}, w(s))$, $0 < \delta < \frac{1}{4}$, $0 < s < \frac{1}{2}$ and $w(s)$ as in (1.2). On account of the particular form of $\varphi(t)$, we get

$$[\varphi(t)]^{-1} = o(\log(1/t)), \quad t \rightarrow 0. \quad (3.1)$$

Let $f(x)$ be the function whose existence is assured by Theorem 2.1 for the particular choice we made of $\varphi(t)$. Let $\varepsilon > 0$ be a positive constant, F and C be the perfect subset and the constant of Lemma 2.2. Call $\bar{f}$ any continuous extension of f from F to $[0, 2\pi]$ such that

$$|\bar{f}(x_1) - \bar{f}(x_2)| \leq C\varphi(|x_1 - x_2|), \\ x_1, x_2 \in [0, 2\pi] \text{ and } 0 < |x_1 - x_2| < 1/2. \quad (3.2)$$

Write now $f = \bar{f} + g$ and consider

$$\int_F \left(\int_0^{2\pi} |g(x) - g(y)| \frac{\bar{w}(|x-y|)}{|x-y|} dy \right) dx. \quad (3.3)$$

Since $g(x) = 0$, $x \in F$, it is enough to estimate

$$\int_F \left(\int_G |g(y)| \frac{\bar{w}(|x-y|)}{|x-y|} dy \right) dx. \quad (3.4)$$

Here, G denotes the complement of F with respect to $[0, 2\pi]$. Write $G = \bigcup_{i=1}^{\infty} I_k$, $I_i \cap I_j = \emptyset$, $i \neq j$, where $\text{dist}(I_k, F)$ satisfies

$$|I_k| \leq \text{dist}(I_k, F) \leq 2|I_k|, \quad k = 1, 2, \dots \quad (3.5)$$

On account of the definition of $\bar{f}$ and the above inequality we have

$$\int_{I_k} |g(y)| dy < K_0 \varphi(|I_k|) |I_k|, \quad (3.6)$$

which is valid for all k and a suitable choice of K_0 .

Interchanging the order of integration in (3.4) and using (3.6) we get

$$\begin{aligned} \sum_1^\infty \int_{I_k} |g(y)| \left(\int_F \frac{\bar{w}(|x-y|)}{|x-y|} dx \right) dy &< (\text{constant}) \sum_1^\infty \left(\int_{I_k} |g(y)| dy \right) [\varphi(|I_k|)]^{-1} \\ &< \text{constant} \sum_1^\infty |I_k|. \end{aligned} \quad (3.7)$$

The finiteness of (3.3) shows that

$$\int_{-\delta}^{\delta} |g(x) - g(x+t)| \frac{\bar{w}(|t|)}{|t|} dt < \infty \quad (3.8)$$

for almost every x in F . By construction the Fourier series of f diverges a.e. The Fourier series of $\bar{f}$ converges a.e. because

$$\int_0^{2\pi} \int_0^{2\pi} |\bar{f}(x) - \bar{f}(y)|^2 \frac{1}{|x-y|} dx dy < \infty. \quad (3.9)$$

(By construction we have that $\varphi(t) < C(1/|\log t|^{1-\delta})$ with $0 < \delta < 1/4$.) (See Theorem 1.14 in [5, p. 164].)

Finally, (3.8) above also holds for $w(t)$ on account of the fact that $w(t) < \bar{w}(t)$. This concludes the proof.

4. A few additional remarks. The Dini condition (1.1) for $x \in E$, $|E| > 0$, is not comparable to the condition

$$\frac{1}{h} \int_0^h |f(x+t) - f(x)| dt = O\left(\frac{1}{|\log h|}\right), \quad x \in E. \quad (4.1)$$

The reason for that is the fact that (1.1) assures convergence of $S_m(f)$ everywhere in E , while (4.1) assures only convergence a.e. in E (see [4] and [5, Vol. I, p. 302 (2.1)]).

Theorem 2.1 above was proved by Marcinkiewicz to show that condition (4.1) can not be replaced by any similar weaker one.

Finally, I would like to thank Professor Zygmund for many opportune and helpful suggestions concerning the matters of this paper.

REFERENCES

1. L. Carleson, *On the convergence and growth of partial sums of Fourier Series*, Acta Math. **116** (1966), 135–157.
2. Richard Hunt, *On the convergence of Fourier Series*, Orthogonal Expansions and their Continuous Analogues, Southern Illinois University Press, 1968, pp. 235–255.
3. J. Marcinkiewicz, *Sur les séries de Fourier*, Fund. Math. **27** (1936), 38–69.
4. ———, *On the convergence of Fourier Series*, J. London Math. Soc. **10** (1935), 264–268.
5. A. Zygmund, *Trigonometric series*, Vols. I and II, Cambridge University Press, 1968.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS – CHICAGO CIRCLE, CHICAGO, ILLINOIS 60680