DERIVATIVES OF POLYNOMIALS WITH POSITIVE COEFFICIENTS

A. K. VARMA

ABSTRACT. Let $P_n(x)$ be an algebraic polynomial of degree n with positive coefficients. We set

$$I_n = \frac{\|P'_n(x)\omega(x)\|_{L_2[0,\infty)}}{\|P_n(x)\omega(x)\|_{L_2[0,\infty)}}.$$

In this work upper bounds of I_n are investigated. We restrict ourselves here with the case $\omega(x) = x^{\alpha/2}e^{-x/2}$. Results are shown to be best possible.

1. The problems dealt with in the present paper are connected with the classical inequalities of A. Markov [5], P. Erdos [1], G. G. Lorentz [3], [4], G. Szego [7] and P. Turan [8]. First, we describe them briefly.

THEOREM A (A. A. MARKOV). *If $f(x)$ is a polynomial of degree n , such that $|f(x)| \leq 1$ for $a \leq x \leq b$, then*

$$(1.1) \quad \max_{a \leq x \leq b} |f'(x)| \leq \frac{2}{b-a} n^2.$$

Further, this bound is sharp, as can be seen by the example

$$f(x) = T_n(2(x-a)/(b-a) - 1)$$

where T_n is Chebyshev's polynomial.

For special polynomials, one can sometimes improve the estimate (1.1).

Next, two theorems are of this kind.

THEOREM B (P. ERDOS). *Let $f(x)$ be a polynomial of degree n satisfying the inequality $|f(x)| \leq 1$ for $-1 \leq x \leq 1$. Suppose $f(x)$ has only real roots and no root inside $[-1, +1]$; then for $-1 \leq x \leq 1$,*

$$(1.2) \quad |f'(x)| \leq \frac{1}{2} en.$$

This is the best possible result.

Inequalities of the same type (as in Theorem B) hold for the wider class of polynomials with positive coefficients in $1+x$, $1-x$; that is, for the polynomials

$$(1.3) \quad f(x) = \sum_{k+l \leq n} a_{kl}(1+x)^k(1-x)^l, \quad a_{kl} \geq 0, \quad -1 \leq x \leq 1.$$

These polynomials were introduced by G. G. Lorentz [3] and studied extensively by J. T. Scheick [6]. The Lorentz theorem can be stated as follows.

Received by the editors April 21, 1980.

1980 *Mathematics Subject Classification*. Primary 26A60; Secondary 30A40.

© American Mathematical Society
0002-9939/81/0000-0425/\$02.50

THEOREM C (G. G. LORENTZ). *There exists a constant $C > 0$ such that for each polynomial $f(x)$ of the form (1.3),*

$$(1.4) \quad \frac{\|f'\|}{\|f\|} \leq Cn, \quad n = 1, 2, \dots,$$

for the uniform norm on $[-1, +1]$.

In 1964 G. Szego [7] studied the order of magnitude of $\|f'\|/\|f\|$ for unrestricted polynomials f of degree $\leq n$ for the norm

$$(1.5) \quad \|f\| = \sup_{x>0} |f(x)e^{-x}|$$

on $(0, \infty)$. More precisely, he proved the following.

THEOREM D (G. SZEGO). *Let $f(x)$ be a polynomial of fixed degree n and not vanishing identically. Then*

$$(1.6) \quad \frac{\|f'\|}{\|f\|} < Cn, \quad n = 2, 3, \dots,$$

holds (for the weighted uniform norm as defined by (1.5) on $(0, \infty)$).

In 1968 G. G. Lorentz [4] considered the problem of G. Szego for the special polynomials with positive coefficients in x ,

$$(1.7) \quad f(x) = \sum_{k=0}^n a_k x^k, \quad a_k > 0,$$

and the norm of a function f on $(0, \infty)$ is given by

$$\|f\| = \sup_{x>0} |f(x)e^{-\omega(x)}|,$$

where ω increases on $(0, \infty)$. Then the Lorentz theorem can be stated as follows.

THEOREM E (G. G. LORENTZ). *Let $\omega(x)$ satisfy the inequalities $\omega(x) - \omega(0) \leq Ax\omega'(x)$, $x > 0$, and $\omega'(y) \leq A\omega'(x)$, $y \leq x$, for some constant $A > 0$. Then for some constant $C > 0$, the inequality*

$$\frac{\|f'\|}{\|f\|} \leq C \frac{\|p_n'\|}{\|p_n\|}, \quad p_n(x) = x^n,$$

is valid.

Other related interesting results are due to A. Zygmund [10], Hille, Szego and Tamerkin [2] and P. Turan [8], where similar problems are considered either in L_2 norm or even in L_p norm.

We may note that Lorentz raised the problem of determining the best constant in (1.6) and for (1.9).

Now we state the main theorem of this paper. Let $S_n[a, b]$ be the set of all polynomials whose degree is n and whose roots are all real and no root inside (a, b) . It is easy to see that if $p_n \in S_n[0, \infty)$ then it can be expressed in the form

$$(1.8) \quad p_n(x) = \sum_{k=0}^n a_k x^k, \quad a_k \geq 0 \text{ for } k = 0, 1, \dots, n.$$

We now state

THEOREM 1. Let $P_n(x)$ be an algebraic polynomial of degree n with nonnegative coefficients. Then for $\alpha \geq (\sqrt{5} - 1)/2$

$$(1.9) \quad \int_0^\infty (P'_n(x))^2 x^\alpha e^{-x} dx \leq \frac{n^2}{(2n + \alpha)(2n + \alpha - 1)} \int_0^\infty P_n^2(x) x^\alpha e^{-x} dx,$$

equality holding for $P_n(x) = x^n$. For $0 \leq \alpha \leq \frac{1}{2}$ we have

$$(1.10) \quad \int_0^\infty (P'_n(x))^2 x^\alpha e^{-x} dx \leq \frac{1}{(2 + \alpha)(1 + \alpha)} \int_0^\infty P_n^2(x) x^\alpha e^{-x} dx.$$

Moreover (1.10) is also best possible in the sense that for $P_n(x) = x^n + \lambda x$ the expression on the left can be made arbitrarily close to the expression on the right-hand side by choosing λ positive and sufficiently large.

COROLLARY. Let $P_n(x)$ be an algebraic polynomial of degree n with nonnegative coefficients satisfying (1.8) with $a_n > 0$. Then for $\alpha \geq (\sqrt{5} - 1)/2$ we obtain

$$(1.11) \quad \int_0^\infty P_n^2(x) x^\alpha e^{-x} dx \geq (a_n)^2 \Gamma(2n + \alpha + 1),$$

equality holding for $P_n(x) = x^n$ ($a_n = 1$).

Proof of this Corollary is as follows. Repeated application of the inequality (1.9) gives at once

$$(1.12) \quad \frac{\int_0^\infty (P_n^k(x))^2 x^\alpha e^{-x} dx}{\int_0^\infty P_n^2(x) x^\alpha e^{-x} dx} \leq \frac{n^2(n-1)^2 \cdots (n-k+1)^2}{(2n+\alpha)(2n+\alpha-1) \cdots (2n+\alpha-2(k-1))(2n+\alpha-1-2(k-1))}.$$

Next, we note that

$$(1.13) \quad \int_0^\infty (P_n^{(n)}(x))^2 x^\alpha e^{-x} dx = (a_n)^2 (n!)^2 \Gamma(\alpha + 1).$$

Now, we substitute $k = n$ in (1.12) and use (1.13) which leads to (1.11).

2. Proof of Theorem 1. Let $P_n(x)$ be any algebraic polynomial of degree n with positive coefficients. Then we may write

$$(2.1) \quad P_n(x) = a_n x^n + P_{n-1}(x), \quad P_{n-1}(x) = \sum_{k=0}^{n-1} a_k x^k, \quad a_k \geq 0.$$

Using (2.1) and

$$(2.2) \quad \int_0^\infty x^k e^{-x} dx = \Gamma(k + 1)$$

we obtain

$$(2.3) \quad \int_0^\infty (P'_n(x))^2 e^{-x} x^\alpha dx = \int_0^\infty (P'_{n-1}(x))^2 e^{-x} x^\alpha dx \\ + a_n^2 n^2 \Gamma(2n + \alpha - 1) + 2na_n \int_0^\infty P'_{n-1}(x) x^{n+\alpha-1} e^{-x} dx,$$

and

$$(2.4) \quad \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx = a_n^2 \Gamma(2n + \alpha + 1) \\ + \int_0^\infty (P_{n-1}(x))^2 e^{-x} x^\alpha dx + 2a_n \int_0^\infty P_{n-1}(x) e^{-x} x^{n+\alpha} dx.$$

Next we set

$$(2.5) \quad b_n = \frac{n^2}{(2n + \alpha)(2n + \alpha - 1)}$$

and

$$(2.6) \quad \lambda_n = 2n \int_0^\infty P'_{n-1}(x) x^{n+\alpha-1} e^{-x} dx - 2b_n \int_0^\infty P_{n-1}(x) x^{n+\alpha} e^{-x} dx.$$

From (2.3)–(2.6) we may conclude that

$$(2.7) \quad \int_0^\infty (P'_n(x))^2 e^{-x} x^\alpha dx - b_n \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx \\ = \lambda_n a_n + \int_0^\infty (P'_{n-1}(x))^2 e^{-x} x^\alpha dx - b_n \int_0^\infty (P_{n-1}(x))^2 e^{-x} x^\alpha dx.$$

First, we claim that for $\alpha > 0$

$$(2.8) \quad \lambda_n < 0, \quad n = 1, 2, \dots$$

From (2.1), (2.2) and (2.6) we obtain

$$(2.9) \quad \lambda_n = \frac{2n}{(2n + \alpha)(2n + \alpha - 1)} \sum_{k=0}^{n-1} a_k \mu_{kn} \Gamma(k + n + \alpha - 1), \quad a_k > 0,$$

where

$$\mu_{kn} = (2n + \alpha)(2n + \alpha - 1)k - n(k + n + \alpha)(k + n + \alpha - 1) \\ = (k - n)\{n(n - k) + (2\alpha - 1)n + \alpha(\alpha - 1)\}, \quad 0 \leq k \leq n - 1.$$

Clearly for $\alpha > 0$ we have

$$(2.10) \quad \mu_{kn} \leq -\alpha(2n + \alpha - 1) < 0, \quad k = 0, 1, \dots, n - 1.$$

Thus (2.8) follows easily from (2.9) and (2.10). We also note that

$$(2.11) \quad b_n > b_{n-1}, \quad \alpha > \frac{\sqrt{5} - 1}{2}, \quad n = 2, 3, \dots,$$

and

$$(2.12) \quad b_n < b_{n-1}, \quad 0 < \alpha < \frac{1}{2}, \quad n = 2, 3, \dots$$

Using these ideas we can complete the proof of Theorem 1. For $\alpha > (\sqrt{5} - 1)/2$ we use (2.7), (2.8) and (2.11) and we obtain for $k = 2, 3, \dots, n$,

$$(2.13) \quad \begin{aligned} & \int_0^\infty (P'_k(x))^2 e^{-x} x^\alpha dx - b_k \int_0^\infty (P_k(x))^2 e^{-x} x^\alpha dx \\ & < \int_0^\infty (P'_{k-1}(x))^2 e^{-x} x^\alpha dx - b_{k-1} \int_0^\infty (P_{k-1}(x))^2 e^{-x} x^\alpha dx. \end{aligned}$$

Adding all these equations we obtain

$$(2.14) \quad \begin{aligned} & \int_0^\infty (P'_n(x))^2 e^{-x} x^\alpha dx - b_n \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx \\ & < \int_0^\infty (P'_1(x))^2 e^{-x} x^\alpha dx - b_1 \int_0^\infty (P_1(x))^2 e^{-x} x^\alpha dx. \end{aligned}$$

But a simple computation shows that if $P_1(x) = a_1 x + a_0$, $a_1 > 0$, $a_0 > 0$ then

$$(2.15) \quad \int_0^\infty (P'_1(x))^2 e^{-x} x^\alpha dx < b_1 \int_0^\infty (P_1(x))^2 e^{-x} x^\alpha dx,$$

equality holds if $P_1(x) = a_1 x$, $a_1 > 0$. From (2.14) and (2.15) we obtain (1.2) as desired for $\alpha > (\sqrt{5} - 1)/2$. For the proof of (1.3) for $0 < \alpha < \frac{1}{2}$ we use (2.7), (2.8) and (2.12). From these results it follows that

$$(2.16) \quad \begin{aligned} & \int_0^\infty (P'_k(x))^2 e^{-x} x^\alpha dx - b_k \int_0^\infty (P_k(x))^2 e^{-x} x^\alpha dx \\ & \leq \int_0^\infty (P'_{k-1}(x))^2 e^{-x} x^\alpha dx - b_{k-1} \int_0^\infty (P_{k-1}(x))^2 e^{-x} x^\alpha dx \\ & \quad + (b_{k-1} - b_k) \int_0^\infty (P_{k-1}(x))^2 e^{-x} x^\alpha dx. \end{aligned}$$

Also from (2.4) it follows that

$$(2.17) \quad \int_0^\infty P_{k-1}^2(x) x^\alpha e^{-x} dx < \int_0^\infty (P_k(x))^2 x^\alpha e^{-x} dx.$$

From (2.16) and (2.17) we obtain for $k = 2, 3, \dots, n-1, n$,

$$(2.18) \quad \begin{aligned} & \int_0^\infty (P'_k(x))^2 e^{-x} x^\alpha dx - b_k \int_0^\infty (P_k(x))^2 e^{-x} x^\alpha dx \\ & < \int_0^\infty (P'_{k-1}(x))^2 e^{-x} x^\alpha dx - b_{k-1} \int_0^\infty (P_{k-1}(x))^2 e^{-x} x^\alpha dx \\ & \quad + (b_{k-1} - b_k) \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx. \end{aligned}$$

We add all these equations for $k = 2, 3, \dots, n-1, n$ and we obtain

$$(2.19) \quad \begin{aligned} & \int_0^\infty (P'_n(x))^2 e^{-x} x^\alpha dx - b_n \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx \\ & < \int_0^\infty (P'_1(x))^2 e^{-x} x^\alpha dx - b_1 \int_0^\infty (P_1(x))^2 e^{-x} x^\alpha dx \\ & \quad + (b_1 - b_n) \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx \\ & < (b_1 - b_n) \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx. \end{aligned}$$

But clearly (2.18) is equivalent to

$$(2.20) \quad \int_0^\infty (P'_n(x))^2 e^{-x} x^\alpha dx \leq b_1 \int_0^\infty (P_n(x))^2 e^{-x} x^\alpha dx.$$

From (2.19) and (2.5) follows (1.3).

In order to show that (1.3) is best possible we consider $f_0(x) = x^n + \lambda x$, $\lambda > 0$, $0 < \alpha \leq \frac{1}{2}$. A simple computation shows that

$$\frac{\int_0^\infty (f'_0(x))^2 e^{-x} x^\alpha dx}{\int_0^\infty (f_0(x))^2 e^{-x} x^\alpha dx} = \frac{1}{(\alpha + 2)(\alpha + 1)} - C_{n,\lambda},$$

where

$$C_{n,\lambda} = \frac{2\lambda\Gamma(n + \alpha)[\alpha(\alpha + 1) + n^2 - n(\alpha^2 + \alpha - 1)] + d_n}{(\alpha + 2)(\alpha + 1)[\lambda^2\Gamma(\alpha + 3) + 2\lambda\Gamma(n + \alpha + 2) + \Gamma(2n + \alpha + 1)]}$$

and

$$d_n = \Gamma(2n + \alpha - 1)[n^2(2 - 3\alpha - \alpha^2) + 2n(2\alpha - 1) + \alpha(\alpha - 1)].$$

Clearly $C_{n,\lambda} \rightarrow 0$ as $\lambda \rightarrow \infty$ from which our assertion follows.

REFERENCES

1. P. Erdos, *Extremal properties of derivatives of polynomials*, Ann. of Math. (2) **41** (1940), 310–313.
2. E. Hille, G. Szego and J. D. Tamerkin, *On some generalizations of a theorem of A. A. Markhoff*, Duke Math. J. **3** (1937), 729–739.
3. G. G. Lorentz, *The degree of approximation by polynomials with positive coefficients*, Math. Ann. **151** (1963), 239–251.
4. ———, *Derivatives of polynomials with positive coefficients*, J. Approx. Theory **1** (1968), 1–4.
5. A. A. Markov, *On a problem of D. I. Mendeleev*, Izv. Akad. Nauk **62** (1889), 1–24.
6. J. T. Scheick, *Inequalities for derivatives of polynomials of special type*, J. Approx. Theory **6** (1972), 354–358.
7. G. Szego, *On some problems of approximations*, Magyar Tud. Akad. Mat. Kutato Int. Dozl. **2** (1964), 3–9.
8. P. Turan, *Remarks on a theorem of Erhard Schmidt*, Mathematica **2** (25) (1960), 373–378.
9. A. K. Varma, *Some inequalities of algebraic polynomials having real zeros*, Proc. Amer. Math. Soc. **75** (1979), 243–250.
10. A. Zygmund, *A remark on the conjugate series*, Proc. London Math. Soc. **36** (1932), 392–400.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FLORIDA 32611