RIEMANN R_1 -SUMMABILITY OF INDEPENDENT, IDENTICALLY DISTRIBUTED RANDOM VARIABLES

JACK CUZICK

ABSTRACT. Let $X, X_1, X_2, \dots$ be i.i.d. random variables. It is shown that $E|X|\log^+\log^+|X| < \infty$ is a sufficient condition for Riemann R_1 -summability of $\{X_n\}$ to EX . Counterexamples are provided which indicate that the strongest possible necessary condition of moment type is $E|X| < \infty$. However under weak regularity conditions on the tails of the distribution of X the sufficient condition is also shown to be necessary.

1. Introduction. Many regular forms of summability have been studied in great detail for independent, identically distributed random variables $X, X_1, X_2, \dots$; the classic example being the strong law of large numbers which asserts that $E|X| < \infty$ iff $\{X_n\}$ is Cesàro $(C, 1)$ summable to EX . Lai [5] has shown that the existence of the first absolute moment is also necessary and sufficient for Cesàro (C, α) summability for all $\alpha > 1$ as well as for Abel summability, whereas Chow [1] in studying Euler (E, q) , $q > 0$, summability and Borel summability found that the existence of a second moment was equivalent to summability. However, results for Riemann R_1 -summability are not so tidy. Recall that a sequence of real numbers $\{A_n\}$ is R_1 -summable to A if

$$\lim_{t \rightarrow 0^+} \frac{2}{\pi} \sum_{n=1}^{\infty} n^{-1} A_n \sin nt = A. \quad (1)$$

It is well known [3] that this is not a regular summability method. More importantly from our point of view, it is a discrete to continuous method in that the limit in (1) is taken through a continuum of values. In a recent paper Cuzick and Lai [2] have shown that a sequence of i.i.d. random variables $X, X_1, X_2, \dots$ is R_1 -summable to EX if $E|X|\log^+\log^+|X| < \infty$ and also that $E|X| < \infty$ is a necessary condition for summability. The approach there, as also followed here, is to treat (1) as a random Fourier series and, when $EX = 0$, to establish uniform convergence. The purpose of the present note is threefold:

(i) to show that

$$E|X|\log^+\log^+|X| < \infty \quad (2)$$

is a sufficient condition for R_1 -summability,

(ii) to give examples which demonstrate that in general the condition $E|X| < \infty$ is the strongest possible necessary condition of moment type, i.e., given a monotone

Received by the editors November 19, 1979 and, in revised form, May 29, 1980.

1980 *Mathematics Subject Classification*. Primary 60F15, 40G99; Secondary 60G17, 60G50, 42A24.

Key words and phrases. Riemann summability, random Fourier series, almost sure convergence.

© American Mathematical Society

0002-9939/81/0000-0427/\$02.50

function $g(x)$ such that $\lim_{x \rightarrow \infty} g(x)/x = +\infty$ we can find an R_1 -summable random variable X with $Eg(|X|) = +\infty$, and

(iii) to show that under a weak regularity condition on the smoothness of the tails of the distribution function of X the condition (2) is in fact necessary and sufficient.

2. Sufficient conditions. As the constant sequence is R_1 -summable we may take $EX = 0$. Then R_1 -summability follows from the continuity of

$$S(t) = \sum_{n=1}^{\infty} n^{-1} X_n \sin nt \quad (3)$$

which in turn follows from the uniform convergence of the series in (3).

THEOREM 1. Assume $EX = 0$ and $E|X|\log^+ \log^+ < \infty$, where $\log^+ x = \max(0, \log x)$. Then the series in (3) is uniformly convergent and $\{X_n\}$ is R_1 -summable.

PROOF. The proof is a refinement of the techniques used in [2]. Let X have distribution function F and for $\beta > 1$ set

$$\begin{aligned} X'_n &= X_n I_{|X_n| \leq n/\log^\beta n}, & X''_n &= X_n I_{n/\log^\beta n < |X_n| \leq n}, \\ X'''_n &= X_n I_{|X_n| > n} \end{aligned}$$

and write

$$\begin{aligned} S(t) &= \sum_{n=1}^{\infty} (X'_n - EX'_n) n^{-1} \sin nt + \sum_{n=1}^{\infty} EX'_n n^{-1} \sin nt \\ &\quad + \sum_{n=1}^{\infty} X''_n n^{-1} \sin nt + \sum_{n=1}^{\infty} X'''_n n^{-1} \sin nt \\ &= S_1(t) + S_2(t) + S_3(t) + S_4(t), \quad \text{say.} \end{aligned}$$

As $E|X| < \infty$ it follows from the Borel-Cantelli lemma that S_4 has only a finite number of nonzero terms and hence is uniformly convergent. Uniform convergence of S_3 follows from the fact that

$$\begin{aligned} E \sum_{n=3}^{\infty} n^{-1} |X''_n| &= \sum_{n=3}^{\infty} n^{-1} \int_{n/\log^\beta n < |x| \leq n} |x| dF(x) \\ &\leq \text{Const.} \int_{-\infty}^{\infty} \left(\sum_{|x| \leq n < |x| \log^\beta |x|} n^{-1} \right) |x| dF(x) \\ &\leq \text{Const.} \int_{-\infty}^{\infty} |x| \log^+ \log^+ |x| dF(x) < \infty. \end{aligned}$$

The uniform convergence of $S_1(t)$ follows from a result of Marcus [7] (see also [2, remarks after Theorem 3]) by checking that

$$\sum_{n=3}^{\infty} \left[\left\{ \sum_{k=n}^{\infty} E(X'_k)^2 k^{-2} \right\}^{1/2} / n \log^{1/2} n \right] < \infty$$

which follows from the estimate

$$\begin{aligned}\sum_{k=n}^{\infty} E(X'_n)^2 k^2 &= \sum_{k=n}^{\infty} k^{-2} \int_{|x| < k/\log^{\beta} k} x^2 dF(x) \\ &= \int_{|x| < n/\log^{\beta} n} \left(\sum_{k=n}^{\infty} k^{-2} \right) x^2 dF(x) \\ &\quad + \int_{|x| > n/\log^{\beta} n} \left(\sum_{k/\log^{\beta} k > |x|} k^{-2} \right) x^2 dF(x) \\ &= O(\log n)^{-\beta} \quad \text{since } \beta > 1.\end{aligned}$$

As $EX = 0$, we may replace EX'_n by $EXI_{|x| > n/\log^{\beta} n}$ in S_2 . Interchanging summation and expectation as before we find the N th partial sum of $S_2(t)$ equals

$$\int_{-\infty}^{\infty} \left(\sum_{n/\log^{\beta} n < x}^N n^{-1} \sin nt \right) x dF(x). \quad (4)$$

If we compute

$$C(i, t) = \sup_j \left| \sum_{i < n < j} n^{-1} \sin nt \right|$$

and

$$C = \sup_{i, t} C(i, t),$$

it is easy to check that $C < \infty$ and $C(i, t) \downarrow 0$ as $i \uparrow \infty$ uniformly for t on compact sets not containing $\{2\pi n\}_{n=-\infty}^{\infty}$. From this it follows that $S_2(t)$ is uniformly convergent on such sets and we need only concern ourselves with uniform convergence in a neighborhood of the origin, i.e. given $\varepsilon > 0$, it is enough to find a $\delta > 0$ such that when $|t| < \delta$, (4) is less than ε for all N . Choose D so that

$$\int_{|x| > D} |x| dF(x) < \varepsilon/2C.$$

Then the integral (4) on the set $|x| > D$ is less than $\varepsilon/2$ and it suffices to show that for all N

$$\int_{|x| < D} \left(\sum_{N < n/\log^{\beta} n < x} n^{-1} \sin nt \right) x dF(x) < \varepsilon/2$$

for all t sufficiently small. This follows immediately from the convergence to zero of the integrand as $t \rightarrow 0$ uniformly for $|x| < D$, as the series contains only a finite number of terms for $|x|$ bounded.

3. Necessary conditions. Cuzick and Lai [2] noted that $E|X| < \infty$ was necessary for R_1 -convergence. The following example indicates that no stronger necessary condition of moment type can be found. Let $g(x)$ be any function such that $g(x)/x \uparrow \infty$ as $x \rightarrow \infty$. Choose any $0 < \varepsilon < 1$ and let n_k be the first $n > 2n_{k-1}$ and of the form $2^j n$ for some $j > 1$ such that $g(n_k)/n_k > k^{2+\varepsilon}$. Let X have mass $M/(n_k k^{2+\varepsilon})$ at $\pm n_k$ with M chosen so that the total mass is unity. Then $E|X| < \infty$ but $Eg(|X|) = +\infty$. As $E|X_n| < \infty$, by the Borel-Cantelli lemma we may replace

X_n by $X'_n = X_n I_{|X_n| < n}$ and change only a finite number of terms. Now for $n_k \leq 2^n < n_{k+1}$ compute

$$s_n = \left(\sum_{j=2^n+1}^{2^{n+1}} E(X'_j/j)^2 \right)^{1/2} = O\left(\frac{n_k}{2^n k^{2+\varepsilon}} \right)^{1/2}.$$

Then we may choose $s_n^* > s_n$, with s_n^* nonincreasing so that

$$\sum_{n=1}^{\infty} s_n^* = \sum_{k=1}^{\infty} \left(\sum_{n_k < 2^n < n_{k+1}} s_n^* \right) = O\left(\sum_{k=1}^{\infty} k^{-(1+\varepsilon/2)} \right) < \infty.$$

It follows from a result of Kahane [4, p. 65, Remark 2] that

$$\sum_{n=1}^{\infty} X'_n n^{-1} \sin nt$$

is uniformly convergent and thus $\{X_n\}$ is R_1 -summable.

This result relies heavily on the lacunary behavior of the distribution of X . If some mild conditions are placed on the regularity of the tails of the distribution of X , it is possible to show that (2) is also necessary for R_1 -summability.

THEOREM 2. Assume that for all n sufficiently large

$$\int_{2^n < |x| < 2^{n+1}} |x| dF(x) \text{ is nonincreasing as } n \uparrow \infty. \quad (5)$$

Then $\{X_n\}$ is Riemann R_1 -summable $\Leftrightarrow E|X| \log^+ \log^+ |X| < \infty$.

PROOF. We need only establish that under (5), condition (2) is necessary for summability. First assume X is symmetric. We shall need the results of §3 in [2]. Set $W_n = XI_{0 < X < n}$ and let G_n be the unique solution of the equation

$$nE(\min\{|W_n/G_n|, (W_n/G_n)^2\}) = 1. \quad (6)$$

If $\{X_n\}$ is summable it follows from Theorem 5 of [2] that

$$\sum_{n=1}^{\infty} \gamma_n < \infty \quad (7)$$

where $\gamma_n = G_{2^n}/2^n$. We aim to show that

$$\gamma_n = o\left(\frac{1}{n}\right) \quad (8)$$

which follows from (7) if we can establish that γ_n is "almost monotone" in the sense that there exists an n_0 and $K_1 > 0$ such that

$$\gamma_n > K_1 \gamma_m \quad \text{for all } m > n > n_0. \quad (9)$$

To prove this we need to verify the existence of an n_0 and $K_2 > 0$ such that for all $n > j > n_0$

$$2^{-n} \int_0^{2^n} x^2 dF(x) > K_2 2^{-(n+k)} \int_0^{2^{n+k}} x^2 dF(x), \quad \text{all } k > 0 \quad (10)$$

and

$$\int_{2^j}^{2^n} x dF(x) > K_2 \int_{2^{j+k}}^{2^{n+k}} x dF(x), \quad \text{all } k > 0. \quad (11)$$

Equation (11) follows immediately from (5) and (10) follows from (5) upon noting that

$$\begin{aligned} \int_{2^{n_0+1}}^{2^n} x^2 dF(x) &= \sum_{m=n_0}^n \int_{2^{m-1}}^{2^m} x^2 dF(x) \\ &> \sum_{m=n_0}^n 2^{m-1} \int_{2^{m-1}}^{2^m} x dF(x) > \sum_{m=n_0}^n 2^{m-1} \int_{2^{m+k-1}}^{2^{m+k}} x dF(x) \\ &\geq \sum_{m=n_0}^n 2^{k-2} \int_{2^{m+k-1}}^{2^{m+k}} x^2 dF(x) \geq \frac{1}{4} 2^{-k} \int_{2^{n_0+1}}^{2^{n+k}} x^2 dF(x). \end{aligned}$$

Now for $0 < G \leq 2^n$, define

$$g_n(G) = \int_0^G \frac{x^2}{G} dF(x) + \int_G^{2^n} x dF(x). \quad (12)$$

Then g_n is nonincreasing in G . Choose β_n of the form 2^i for some integer i such that $\frac{1}{2}\beta_n < G_{2^n} < \beta_n$. Then using (6)

$$\gamma_n = g_n(G_{2^n}) \geq g_n(\beta_n) \quad (13)$$

which from (10) and (11) is greater than or equal to $K_2 g_{n+k}(2^k \beta_n)$ for all $k \geq 0$. If $2^k \beta_n < G_{2^{n+k}}$ then

$$K_2 g_{n+k}(2^k \beta_n) \geq K_2 g_{n+k}(G_{2^{n+k}}) = K_2 \gamma_{n+k}$$

so that (9) holds. Of course (9) also holds when $2^k \beta_n > G_{2^{n+k}}$ since then $G_{2^n} > \frac{1}{2}\beta_n > 2^{-(k+1)}G_{2^{n+k}}$ so that $\gamma_n = G_{2^n}/2^n \geq \frac{1}{2}\gamma_{n+k}$.

Now define $G(x) = G_{2^n}$ for $x = n$ and by linear interpolation for nonintegral x . It is easily checked that G is nondecreasing. From (8) it follows that there exists $K_3 > 0$, such that for sufficiently large x

$$G^{-1}(x) \geq \log x + K_3 \log \log x. \quad (14)$$

It follows from (7), (13) and (12) that

$$\sum_{n=1}^{\infty} \int_{G(n)}^{2^n} x dF(x) < \infty$$

and a Fubini argument shows

$$\int_0^{\infty} [G^{-1}(x) - \log x] x dF(x) < \infty$$

from whence (2) follows with the help of (14). This establishes the theorem when X is symmetric. In general let m be a median of X and for $2^k < n < 2^{k+1}$ define

$$Y_n = (X_n - m)I_{|X_n - m| < 2^{k-1}}$$

and let Z_n be the symmetrized version of Y_n , i.e. $Z_n = Y_n - Y'_n$ where Y_n, Y'_n are i.i.d. Then $\{X_n\}$ -summable $\Rightarrow \{Z_n\}$ -summable

$$\Rightarrow \sum_{k=1}^{\infty} \gamma_k^Z < \infty \quad \text{as at (7)} \quad (15)$$

where $\gamma_k^Z = G_{2^k}^Z/2^k$ and $G_{2^k}^Z$ is the unique solution of (6) with W_n replaced by Z_{2^k} .

Define γ_k^Y and $G_{2^k}^Y$ similarly. Then integration by parts and the weak symmetrization inequality [6, p. 245] gives

$$\begin{aligned}\gamma_k^Z &= E(\min\{Z_{2^k}^2/G_{2^k}^Z, |Z_{2^k}|\}) = \int_0^\infty \min\{x^2/G_{2^k}^Z, x\} dF_{|Z_{2^k}|}(x) \\ &> \frac{1}{2} \int_0^\infty (1 - F_{|Z_{2^k}|}(x)) \min\{2x/G_{2^k}^Z, 2\} dx \\ &> \frac{1}{4} \int_0^\infty \left(1 - F_{|Y_{2^k}|}\left(\frac{x}{2}\right)\right) \min\{2x/G_{2^k}^Z, 2\} dx \\ &> \frac{1}{16} \int_0^\infty (1 - F_{|Y_{2^k}|}(x)) \min\{2x/G_{2^k}^Z, 2\} dx \\ &> \frac{1}{16} E(\min\{Y_{2^k}^2/G_{2^k}^Z, |Y_{2^k}|\}).\end{aligned}$$

From this it can be shown as in the argument following (13) that

$$\gamma_k^Z \geq \frac{1}{8} \gamma_k^Y. \quad (16)$$

It is easily verified from (6) that for k large there exists $K_3 > 0$ such that

$$G_{2^k}^Y > K_3 2^{k/2}. \quad (17)$$

Now for $G_{2^k}^Y > 2m$ we have

$$\begin{aligned}\gamma_k^Y &= E(\min\{Y_{2^k}^2/G_{2^k}^Y, |Y_{2^k}|\}) \\ &= E(\min\{(X - m)^2 I_{|X-m| < 2^{k-1}}/G_{2^k}^Y, |X - m| I_{|X-m| < 2^{k-1}}\}) \\ &> \frac{1}{2} E(\min\{X^2 I_{|X| < 2^k}/G_{2^k}^Y, |X| I_{|X| < 2^k}\} - m^2/G_{2^k}^Y) \\ &> \frac{1}{2} (\gamma_k^X - m^2/G_{2^k}^Y).\end{aligned}$$

Combining (15), (16) and (17) gives

$$\sum_{k=1}^{\infty} \gamma_k^X < \infty. \quad (18)$$

If we let $|V| \stackrel{\mathfrak{D}}{=} |X|$ and let V be symmetric, i.e. $V = \pm X$, then $\gamma_k^V = \gamma_k^X$ and, as V satisfies (5) and is symmetric, we may use the proof previously given to see that (18) implies $\{V_n\}$ is summable

$$\Rightarrow E|X| \log^+ \log^+ |X| = E|V| \log^+ \log^+ |V| < \infty$$

which completes the proof in general.

REFERENCES

1. Y. S. Chow, *Delayed sums and Borel summability of independent, identically distributed random variables*, Bull. Inst. Math. Acad. Sinica **1** (1973), 207–220.
2. Jack Cuzick and T. L. Lai, *On random Fourier series*, Trans. Amer. Math. Soc. **261** (1980), 53–80.
3. G. H. Hardy and W. W. Rogosinski, *Notes on Fourier series (V): Summability (R_1)*, Proc. Cambridge Philos. Soc. **45** (1949), 173–185.
4. J. P. Kahane, *Some random series of functions*, Heath, Lexington, Mass., 1968.
5. T. L. Lai, *Summability methods for independent, identically distributed random variables*, Proc. Amer. Math. Soc. **45** (1974), 253–261.
6. M. Loeve, *Probability theory*, 3rd ed., Van Nostrand, Princeton, N. J., 1963.
7. M. B. Marcus, *Continuity and the central limit theorem for random trigonometric series*, Z. Wahr. Verw. Gebiete **42** (1978), 35–56.