LIMIT SETS OF AUTOMORPHISM GROUPS OF A TREE

SU-SHING CHEN

ABSTRACT. Let X be an infinite and locally finite tree and G be an arbitrary automorphism group of X . The limit set $L(G)$ of G in the boundary B of X is studied.

1. Introduction. A tree is a nonempty connected graph X without circuit. One can associate a tree to SL_2 over the field Q_p as a special case of the Bruhat-Tits building which is the p -adic analogue of homogeneous symmetric spaces of real semisimple Lie groups [1], [2], [7], [8]. Other applications are the graph $\Gamma(G, P)$ of a group G and a subset P of G and algebraic geometry [7]. Automorphism groups and harmonic analysis on a tree have been investigated by Tits [8] and Cartier [1], [2] respectively.

In the theory of hyperbolic plane and fuchsian groups one has the definition of the limit set $L(G)$ of a subgroup $G \subseteq SL_2(R)$. This concept will be defined for an arbitrary automorphism group G on an infinite and locally finite tree X . We shall obtain some general facts about $L(G)$ such as the cardinality of $L(G)$ and the relative size of $L(G)$ in the boundary B of X . In [5] and [6], Kulkarni has considered this question for a general class of topological spaces. He has also investigated the connection with the theory of ends of Freudenthal and Hopf. Our main result is a theorem similar to that of Ahlfors for Kleinian groups [3].

2. Tree. Let X be an infinite and locally finite tree with vertex set S and edge set A . Let s and t be vertices and be connected. Then the edge $\{s, t\}$ is in A . (For simplicity we do not emphasize orientation here.) A path in X is a sequence $C = [s_0, s_1, \dots, s_n]$ of vertices such that s_{i-1} is connected to s_i for $1 < i < n$. The length of C is n . If all the vertices in a path C_0 are distinct, then this unique path C_0 is called a geodesic. The length of the geodesic joining vertices s to t is called the distance $d(s, t)$ between them. Let s be a vertex. The infinite path originated from s is an infinite sequence $C = [s_0, s_1, \dots, s_n, \dots]$ of vertices such that $s_0 = s$ and s_n and s_{n+1} are connected for each $n > 0$. C is called an infinite geodesic if $s_i \neq s_j$ for $i \neq j$. The relation that two infinite geodesics C and C' have infinitely many common vertices is an equivalence relation. Each equivalence class is called an end. The set B of all ends is the boundary to X . We can join each s in S and each b in B by an infinite geodesic. Moreover, each pair $b \neq b'$ in B can be joined by an infinite geodesic. With the topology described in Theorem 1.1 of [2], the space

Received by the editors October 31, 1979 and, in revised form, February 19, 1981.

1980 *Mathematics Subject Classification*. Primary 05C05, 05C25; Secondary 22E40.

© 1981 American Mathematical Society
 0002-9939/81/0000-0497/\$02.25

$\hat{S} = S \cup B$ is a compact and totally disconnected space such that S , with the discrete topology, is an open dense subspace of $\hat{S}$. Let s and s' be two vertices. If the vertex t moves along the infinite geodesic representing the end b , then $d(s, t) - d(s', t)$ is a finite constant $\delta_b(s, s')$. A horocycle H through s is the set of vertices s' such that $\delta_b(s, s') = 0$. There is a partition of S into a disjoint union of horocycles $\{H_n\}_{n=0}^\infty$.

Let g be an automorphism of X . Tits [8] has proved that one of the following holds:

- (1) there exists s in S such that $g(s) = s$,
- (2) there exist s and s' in S such that $\{s, s'\}$ is an edge and $g(s) = s'$ and $g(s') = s$,
- (3) there exist $\{s_n\}_{n=-\infty}^\infty$ such that $\{s_n, s_{n+1}\}$ is an edge and $g(s_n) = s_{n+k}$ for any n and some $k > 0$.

Furthermore, he has proved that if G is an automorphism group of X and G has no element of type (3), then G leaves a vertex s , an edge $\{s, s'\}$ or an end b (together with all horocycles associated to b) invariant. It is obvious that automorphisms of X are isometries with respect to the distance $d(s, s')$ defined above. Each automorphism g of X can be extended to act on the boundary B . The action is a homeomorphism.

3. Limit set. With the topology on $\hat{S} = S \cup B$, we define the limit set $L(G)$ of an automorphism group G to be the set of accumulation points in B of an orbit Gs of s . That is $L(G) = \overline{Gs} \cap B$. There is the question as to whether $L(G)$ is independent of the choice of the vertex s . Let s and s' be two vertices with finite distance $d(s, s') = d$. We have two orbits Gs and Gs' . Each b in $L(G)$ is the limit $\lim g_n s$ for some sequence g_n in G . Let b' be the limit $\lim g_n s'$. Since $d(s, s') = d(g_n s, g_n s') = d$ and $d(b, b') = \infty$, we have $b = b'$. Consequently, $L(G)$ is independent of the choice of s .

PROPOSITION 1. (i) $L(G)$ is a closed invariant subset of B under G .

(ii) Let A be any closed invariant subset of B under G and let A contain more than one point. Then $A \supset L(G)$ and $L(G)$ is minimal under the action of G on B .

(iii) The set of fixed points of elements of type (3) (if they exist) is dense in $L(G)$.

PROOF. It is clear that $L(G)$ is closed. The invariance of $L(G)$ is seen by $b = \lim g g_n(s) = g \lim g_n(s) = g(b)$. To prove (ii), let b and b' be in A . Let b'' be in $L(G)$ such that $\lim g_n s = b''$. The two infinite points b and b' can be joined by an infinite geodesic. We choose s to be on that geodesic. Then either $g_n b$ or $g_n b'$ must converge to b'' . Thus $A \supset L(G)$. (iii) follows from (ii) directly. In fact, each element of type (3) has two fixed points in B . The closure of fixed points of elements of type (3) in G can be taken as A .

PROPOSITION 2. The cardinality of $L(G)$ of an automorphism group G is 0, 1, 2 or ∞ .

PROOF. Either G contains no elements of type (3) or it does contain them. In the first case, G has a fixed point s or an invariant edge $\{s, s'\}$ or a fixed end b . If G

has a fixed point s or an invariant edge $\{s, s'\}$, then $L(G)$ is empty. If G has a fixed end b , then G leaves horocycles associated to b invariant.

The orbit Gs will converge to b , if we take s to be a vertex on a horocycle associated to b . The limit set $L(G)$ consists of one point. In the second case, $L(G)$ contains at least two points, because two ends b and b' of the infinite geodesic translated by an element g of type (3) will be limit points. If $L(G)$ has more than two points, then $L(G)$ has infinitely many points. In fact, if $L(G)$ has finitely many points, then the n th power g^n of any element g of G will leave every point in $L(G)$ fixed when n is sufficiently large. Then g^n has at least two fixed ends and must translate the infinite geodesic joining them. Thus g is of type (3) and $L(G)$ can have only two points.

REMARK. We may obtain a classification (see Corollary 3.5 of [8] also) of automorphism groups on an infinite and locally finite tree X . There are the following possibilities:

- (i) G has a common fixed point s in S or has a common invariant edge $\{s, t\}$ in A . $L(G)$ is empty.
- (ii) G has a common fixed end b in B and $L(G)$ consists of one point.
- (iii) G has two common fixed ends b_1 and b_2 in B and $L(G)$ consists of two points.
- (iv) G has an invariant proper subtree X' (not a vertex, an edge or an infinite geodesic) of X and $L(G)$ is the boundary B' of X' .
- (v) G acts minimally on B and $L(G) = B$.

From the above, we also obtain a classification of quotient graphs $G \backslash X$ of X if G acts freely on X . There are the following possibilities:

- (i) $G \backslash X$ is parabolic if $L(G)$ has one point and $G \backslash X$ is a union of graphs $G \backslash H_n$, which are the quotient graphs of horocycles H_n by G .
- (ii) $G \backslash X$ is axial if $L(G)$ consists of two points. There is an infinite geodesic C which is translated by G and $G \backslash C$ is the unique circuit such that the injection $G \backslash C \rightarrow G \backslash X$ is a homotopy equivalence [7].
- (iii) $G \backslash X$ is fuchsian if $L(G)$ consists of infinitely many points. This is the most interesting case and offers a great variety of quotient graphs.

4. Boundary measure. We use §§2.5, 3.3 and 3.4 of [2] to investigate the boundary measure of $L(G)$ of an automorphism group G . We refer to [1], [2] for details. Let X be an infinite and locally finite tree. For each finite path a , we assign a positive number $p(a)$ such that $p(aa') = p(a)p(a')$ if the composite path aa' is defined, and for $s, s' \in S$, $\sum p(ss') = 1$ for all s' connected to s . If h is a function on S , we define the function Ph on S by $Ph(s) = \sum p(ss')h(s')$ over the set of all vertices s' connected to s . Let $\Delta h = Ph - h$ be the Laplace operator and let $K(s, b)$ be the Martin kernel of X . There exists a positive measure μ of mass 1 on B such that

$$\int_B K(s, b)\mu(db) = 1$$

for each s in S . This measure is called the Poisson measure. Consider a random walk $\{X_0, X_1, \dots, X_n, \dots\}$ on S in the following manner. If $X_0 = 0$ is the original point and $X_n = s$, then the permissible values of X_{n+1} are vertices s' connected to s

with probability $p(s, s')$ from s to s' . By the Borel-Cantelli lemma, X_n converges to a point X_∞ in B with probability 1, with respect to the probability law given by the Poisson measure μ on B . If h is a bounded harmonic function, the Martingale theorem shows that the sequence of random variables $\{h(X_n)\}$ converges to a random variable Y with probability 1. There exists a bounded Borel function ϕ on B such that $Y = \phi(X_\infty)$ and one has the Poisson integral formula

$$h(s) = \int_B K(s, b)\phi(b)\mu(db), \quad \text{any } s \in S.$$

The radial convergence theorem of Fatou holds in the following sense: $\phi(b) = \lim h(s_n)$, almost everywhere in B with respect to μ , on an infinite geodesic $[s_0, s_1, \dots, s_n, \dots]$ joining the original point 0 to b .

A limit point b in $L(G)$ is said to be an approximation limit point if for any sequence $\{s_n\}$ in a geodesic approaching to b , there exists a sequence $\{g_n\}$ in G such that $g_n(s_n)$ lies in a finite set (compact set). This notion is originally due to Hedlund ([3], [6] for further references). Because $g_n(s_n)$ lies in a finite set, without loss of generality we may assume that they are all equal. Thus, we can find a sequence $\{s_k\}$ converging to b and $\{g_k\} \subseteq G$ such that $\{g_k(s_k)\}$ is a point in S . It is obvious that fixed points of elements of type (3) are approximation limit points. Also the set $A(G)$ of approximation limit points of G is G -invariant.

THEOREM. *Let X be an infinite and locally finite tree and G be an automorphism group of X . Then the Poisson measure $\mu(A(G))$ of the set $A(G)$ of approximation limit points of G is either 0 or 1. If $L(G)$ contains almost all approximation limit points, then the Poisson measure $\mu(L(G))$ is either 0 or 1.*

PROOF. The characteristic function ϕ on B of $A(G)$ is G -invariant and takes values 0 and 1. Let h be the harmonic function

$$h(s) = \int_B K(s, b)\phi(b)\mu(db).$$

Then h is also G -invariant. If the measure $\mu(A(G))$ is positive, then there is a point b in $A(G)$ such that Fatou's convergence theorem can be applied. There exists an infinite geodesic C approaching b such that the radial limit of $h(s)$ along C approaching b must exist and is equal to 1. Note that $0 < h(s) < 1$ for all s in S . Since each b in $A(G)$ is an approximation limit point, there is a sequence $\{s_n\}$ on C converging to b and there is a sequence $\{g_n\}$ in G such that $g_n(s_n)$ is a point s_0 in S . Then the G -invariance and continuity implies that $h(s_0) = 1$. But 1 is the upper bound for $h(s)$. Hence $h(s)$ attains its maximum at an interior point s_0 in S . $h(s)$ has to be the constant 1 and

$$\begin{aligned} \mu(A(G)) &= \int_{A(G)} K(s, b)\mu(db) \\ &= \int_B K(s, b)\phi(b)\mu(db) \quad (\phi \text{ is the characteristic function}) \\ &= \int_B K(s, b)\mu(db) \quad \left(\text{since } \phi(b) = \lim_{s \rightarrow b} h(s) = 1 \right) = 1. \end{aligned}$$

If $L(G)$ contains almost all approximation limit points (that is, except a set of measure zero), then we have $\mu(L(G))$ is 0 to 1 by the above result of $A(G)$.

REFERENCES

1. P. Cartier, *Géométrie et analyse sur les arbres*, Séminaire Bourbaki, 1971/72, Lecture Notes in Math., vol. 317, Springer-Verlag, Berlin and New York, 1973, pp. 123–140.
2. ———, *Fonctions harmoniques sur un arbre*, Symposia Math. **9** (1972), 203–270.
3. S. Chen, *A theorem of Ahlfors for hyperbolic spaces*, Trans. Amer. Math. Soc. **242** (1978), 401–406.
4. P. Eberlein and B. O'Neill, *Visibility manifolds*, Pacific J. Math. **46** (1973), 45–109.
5. R. Kulkarni, *Groups with domains of discontinuity*, Math. Ann. **237** (1978), 253–272.
6. ———, *Some topological aspects of Kleinian groups*, Amer. J. Math. **100** (1978), 897–911.
7. J. Serre, *Arbres, amalgames, SL_2* , Soc. Math. France Astérisque, No. 46, 1977.
8. J. Tits, *Sur le groupe des automorphismes d'un arbre*, Essays on Topology and Related Topics, Mémoires dédiés à G. de Rham, Springer-Verlag, Berlin and New York, 1970, pp. 188–211.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FLORIDA 32611