A NOTE ON STRICTLY CYCLIC WEIGHTED SHIFTS

HECTOR N. SALAS

ABSTRACT. We answer negatively Shields' Question 15 in [1].

The object of this note is to answer negatively one of the many questions raised by Allen L. Shields in his article [1]. Question 15 of [1] is: If T is a unilateral weighted shift with weights w_n decreasing to 1 and if $\sum_{n=0}^{\infty} \beta(n)^{-2} < \infty$ where $\beta(n) = w_0 w_1 \cdots w_{n-1}$, must T be strictly cyclic? If $\{w_n\}$ decreases to 0, must T be strictly cyclic? Strongly strictly cyclic?

By Proposition 32 of [1, p. 96], to answer the question negatively it will suffice to show that the supremum

$$(1) \quad \sup_n \sum_{k=0}^n \left(\frac{\beta(n)}{\beta(k)\beta(n-k)} \right)^2$$

is infinite.

For the first part of the question, we let $\{a_j\}$ be any sequence decreasing to 1 and we choose an increasing sequence of positive integers so that

$$(2) \quad [a_1^{n_1} a_2^{n_2 - n_1} \cdots a_j^{n_j - n_{j-1}}]^{-2} \left[\frac{a_{j+1}^2}{a_{j+1}^2 - 1} \right] < 2^{-j}$$

and

$$(3) \quad (1 + n_j) \left(\frac{a_j}{a_1} \right)^{2n_{j-1}} > j.$$

Such a choice is clearly possible. We set $w_n = a_1$, if $0 \leq n < n_1$, and we set $w_n = a_j$, if $n_{j-1} \leq n < n_j$. In order to verify that our selection works we use (2) to show that $\sum_{n \geq 0} \beta(n)^{-2} < a_1^2 / (a_1^2 - 1) + 1$ and (3) to show that the supremum (1) is infinite.

Consider (2) first. We have

$$(4) \quad \sum_{n=0}^{\infty} \frac{1}{\beta(n)^2} = \sum_{n=0}^{n_1} \frac{1}{\beta(n)^2} + \sum_{j=2}^{\infty} \left(\sum_{n_{j-1} \leq n < n_j} \frac{1}{\beta(n)^2} \right)$$

where $\beta(n)$ is given by the equations $\beta(n) = a_1^n$ if $n < n_1$, and $\beta(n) = a_1^{n_1} a_2^{n_2 - n_1} \cdots a_j^{n - n_{j-1}}$ if $n_{j-1} \leq n < n_j$.

Received by the editors March 12, 1981.

1980 *Mathematics Subject Classification*. Primary 47B37.

© 1981 American Mathematical Society
0002-9939/81/0000-0525/\$01.50

Therefore, $\sum_{n=0}^{n_1} \beta(n)^{-2} < \sum_{n=0}^{\infty} a_1^{-2n} = a_1^2/(a_1^2 - 1)$, while

$$\begin{aligned} \sum_{n_{j-1} < n \leq n_j} \frac{1}{\beta(n)^2} &= [a_1^{n_1} a^{n_2 - n_1} \cdots a_{j-1}^{n_{j-1} - n_{j-2}}]^{-2} \left(\sum_{n_{j-1} < n \leq n_j} \frac{1}{a_j^{2(n - n_{j-1})}} \right) \\ &< [a_1^{n_1} \cdots a_{j-1}^{n_{j-1} - n_{j-2}}]^{-2} \left[\frac{a_j^2}{a_j^2 - 1} \right] < 2^{-j+1}. \end{aligned}$$

This with (4) shows that $\sum_{n=0}^{\infty} \beta(n)^{-2} < a_1^2/(a_1^2 - 1) + 1$.

To show that (1) is violated we show that a lower bound for

$$(5) \quad \sum_{k=0}^{n_j} \left(\frac{\beta(n_j)}{\beta(k)\beta(n_j - k)} \right)^2$$

is j . Consider

$$\frac{\beta(n_j)}{\beta(k)\beta(n_j - k)} = \frac{w_k \cdots w_{n_j-1}}{w_0 \cdots w_{n_j-(k+1)}};$$

in both top and bottom there are no more than n_{j-1} weights different from a_j . Thus, we can say that $\beta(n_j)/\beta(k)\beta(n_j - k) > (a_j/a_1)^{n_{j-1}}$, which together with (3) gives (5).

Note that it is actually possible to choose the sequence $\{w_n\}$ strictly decreasing if desired. One might ask if it is possible to choose the weight sequence to be convex.

The second part of the question is answered with a similar construction. Simply choose $\{a_j\}$ decreasing to zero and choose $\{n_j\}$ so that (3) is satisfied. If $\{w_n\}$ is defined as before, then the supremum (1) is infinite, and T is not strictly cyclic.

REFERENCES

1. A. L. Shields, *Weighted shift operators and analytic function theory*, Math. Surveys No. 13, Amer. Math. Soc., Providence, R.I., 1974.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242