

## SHORTER NOTES

The purpose of this department is to publish very short papers of unusually elegant and polished character, for which there is no other outlet.

### THE RIEMANN INTEGRAL IN CONSTRUCTIVE MATHEMATICS

DAVID SPRING

**ABSTRACT.** A suitable Riemann integral in one variable is developed for use in constructive mathematics.

In this note we briefly define and prove the main properties of the Riemann integral (one variable) in a manner which is suitable for constructive mathematics [1], [2]. In particular we correct the development of the Riemann integral in Bishop [1]. Our definition generalizes easily to multiple Riemann integrals; however, we postpone to a subsequent paper the discussion of multiple integrals where also Stokes' theorem will be proved.

The main difficulty is that the constructive real numbers are not totally ordered. In particular, two partitions of an interval (defined classically) may not have a common refinement.

To begin, we first define  $\int_a^b f(t) dt$  where  $f$  is a real valued uniformly continuous map on the interval  $[a, b]$ ,  $a < b$ . We consider only partitions of  $[a, b]$  of the following form:  $P \equiv a_0 = a < a_1 < a_2 < a_3 < \cdots < a_n < b = a_{n+1}$ , where  $a_i$  is rational,  $i = 1, 2, 3, \dots, n$ . Note the strict inequality at the "extremities" of  $P$ . The point is that two such partitions admit a common refinement (the rationals are totally ordered).

*Notation.* (a) mesh  $P = \sup\{a_{i+1} - a_i \mid i = 0, 1, 2, 3, \dots, n\}$ ,

(b)  $S(f; P) = \sum_{i=0}^n f(a_i)(a_{i+1} - a_i)$ .

The standard classical arguments now apply to prove that  $\lim_{\text{mesh } P \rightarrow 0} S(f; P)$  exists (cf. [1]). Let  $\int_a^b f \equiv \int_a^b f(t) dt$  denote this limit.

**LEMMA 1.** *Let  $f: [a, b] \rightarrow R$  be uniformly continuous,  $a < b$ . Let  $c \in (a, b)$ . Then  $\int_a^b f = \int_a^c f + \int_c^b f$ .*

---

Received by the editors October 21, 1980.

1980 *Mathematics Subject Classification.* Primary 03F55.

*Key words and phrases.* Total order, partition.

© 1981 American Mathematical Society  
0002-9939/81/0000-0585/\$01.50

PROOF. The standard classical argument applies, subject to the following modification: Let  $P \equiv a < a_1 \leq a_2 \leq \dots \leq a_n < c$ ;  $Q \equiv c < c_1 \leq c_2 \leq \dots \leq c_m < b$ , be partitions of  $[a, c]$ ,  $[c, b]$ , respectively. Let  $d$  be a rational approximation to  $c$  such that  $a_n < d < c_1$ . Let  $R \equiv a < a_1 \leq a_2 \leq \dots \leq a_n < d < c_1 \leq c_2 \leq \dots \leq c_m < b$ . It is clear that for mesh  $P$ , mesh  $Q$ ,  $|d - c|$  sufficiently small,  $S(f; R)$  is arbitrarily close to  $\int_a^b f(t) dt$ . Thus Lemma 1 is proved. Let  $f$  be uniformly continuous on an open interval  $J$ . One fixes  $a, b \in J$ .

We define  $\int_a^b f$  in the following way: Let  $x \in J$ ,  $x < \inf\{a, b\}$  [recall that  $J$  is open].

DEFINITION.  $\int_a^b f = \int_x^b f - \int_x^a f$ .

LEMMA 2.  $\int_a^b f$  is well defined.

PROOF. Let  $x, y \in J$ ;  $x < \inf\{a, b\}$ ;  $y < \inf\{a, b\}$ . It is required to prove that

$$\int_x^b f - \int_x^a f = \int_y^b f - \int_y^a f.$$

To this end, let  $z \in J$ ,  $z < \inf\{x, y\}$ . Applying Lemma 1,

$$\int_x^b f = \int_z^b f - \int_z^x f; \quad \int_x^a f = \int_z^a f - \int_z^x f.$$

Consequently,

$$\int_x^b f - \int_x^a f = \int_z^b f - \int_z^a f.$$

Similarly,

$$\int_y^b f - \int_y^a f = \int_z^b f - \int_z^a f.$$

This completes the proof of Lemma 2.

COROLLARY. Let  $a, b, c \in J$ . Then  $\int_a^b f = \int_a^c f + \int_c^b f$ . [One chooses  $x \in J$ ,  $x < \inf\{a, b, c\}$ .]

REMARK. Evidently, the above corollary permits one to prove the relation  $d[\int_a^x f]/dx = f(x)$ .

#### BIBLIOGRAPHY

1. E. Bishop, *Foundations of constructive analysis*, McGraw-Hill, New York, 1967.
2. D. S. Bridges, *Constructive functional analysis*, Pitman, New York, 1979.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA M5S 1A1  
(Current address)