FIXED POINTS OF NONEXPANSIVE CONDENSING MULTI-VALUED MAPPINGS ON METRIC SPACES

CHI-LIN YEN¹

ABSTRACT. In this paper we consider the existence of fixed points of nonexpansive condensing multi-valued mappings from a certain kind of metric space into itself; the spaces, here, are neither linear nor compact. Our result generalizes a theorem of Dotson and also a theorem of Bose and Mukherjee in some respect.

In this note we consider the existence of fixed points of nonexpansive condensing multi-valued mappings from a certain kind of metric space into itself. This result generalizes a theorem of Dotson [2] and also a theorem of Bose and Mukherjee [1] in some respect.

Let (X, d) be a metric space. Denote by 2^X the set of all closed subsets of X and by H the Hausdorff metric on 2^X induced by metric d .

DEFINITION 1. If $F = \{f_A \mid A \in 2^X\}$ is a family of functions from $[0, 1]$ into 2^X satisfying the following conditions:

- (I) $f_A(1) = A$ for all $A \in 2^X$,
- (II) there is a function $\phi: [0, 1] \rightarrow [0, 1]$ with

$$H(f_A(t), f_B(t)) \leq \phi(t)H(A, B)$$

for all A, B in 2^X and $t \in [0, 1]$,

- (III) for all $A \in 2^X$, all $t_0 \in [0, 1]$,

$$\lim_{t \rightarrow t_0} H(f_A(t), f_A(t_0)) = 0,$$

and for all $t \in [0, 1]$,

- (IV) $f_A(t) \subset f_B(t)$ for all A, B in 2^X with $A \subset B$, and

- (V) $f_A(t)$ is compact whenever the set A is finite;

we call F an (S) -convex structure on X .

REMARK 1. If X is a starlike convex subset of a Banach space and if x_0 is a starlike center of X , then for each $A \subset X$, we define

$$f_A(t) = (1 - t)x_0 + tA, \quad t \in [0, 1].$$

$F = \{f_A \mid A \in 2^X\}$ is an (S) -convex structure.

Received by the editors October 9, 1979 and, in revised form, July 16, 1980, December 5, 1980 and August 18, 1981.

1980 *Mathematics Subject Classification.* Primary 54H25, 47H10; Secondary 54C60.

¹ This work was supported by NSC, Republic of China.

© 1982 American Mathematical Society
0002-9939/81/0000-1035/\$02.25

Let γ denote the measure of noncompact sets on X [4], that is, for each bounded subset A of X ,

$$\gamma(A) = \inf\{r > 0 \mid A \text{ can be covered by finitely many balls of radii } \leq r\}.$$

DEFINITION 2. Let $T: X \rightarrow 2^X$.

(1) T is said to be nonexpansive if for any $x \in X$ and any $y \in X$, we have

$$H(Tx, Ty) \leq d(x, y).$$

(2) A continuous mapping T of (X, d) into $(2^X, H)$ is said to be condensing if for any bounded set $A \subset X$, we have

$$\gamma(T(A)) < \gamma(A)$$

provided that $\gamma(A) \neq 0$.

Our result is the following:

THEOREM. *Let (X, d) be a bounded complete metric space with an (S) -convex structure F . Then every nonexpansive condensing multi-valued mapping T of X into 2^X has a fixed point.*

Before we prove our theorem, we first need some preliminary results related to measure of noncompact sets as follows:

LEMMA 1. *Let γ be defined as before. Then*

- (1) $\gamma(A) = 0$ if and only if $\bar{A}$ is compact,
- (2) $\gamma(\cup_{i=1}^n A_i) = \max\{\gamma(A_i) \mid 1 \leq i \leq n\}$,
- (3) $\gamma(A) \leq \gamma(B)$ whenever $A \subset B$,
- (4) $\gamma((A)_r) \leq \gamma(A) + r$, where $r \geq 0$ and $(A)_r = \{b \in X \mid d(b, A) \leq r\}$,
- (5) $\gamma(\bar{A}) = \gamma(A)$ for all bounded subsets A of X .

The proof of Lemma 1 is known (see Sadovskii [5]).

REMARK 2. It is easy to see that $r > \gamma(A)$ if and only if there is a finite set G such that $(G)_r \supset A$ if and only if there is a finite set J such that $H(J, A) \leq r$.

In order to prove our theorem we need the following three lemmas.

LEMMA 2. *Suppose that F satisfies (II) and (III) of Definition 1. Then for $\{A_n \mid n = 0, 1, 2, \dots\} \subset 2^X$ and $\{t_n \mid n = 0, 1, 2, \dots\} \subset [0, 1]$ with $\lim_{n \rightarrow \infty} H(A_n, A_0) = 0$ and $\lim_{n \rightarrow \infty} t_n = t_0$, we have*

$$\lim_{n \rightarrow \infty} H(f_{A_n}(t_n), f_{A_0}(t_0)) = 0.$$

PROOF. Since the triangle inequality is valid for the Hausdorff metric and (II) we have

$$\begin{aligned} H(f_{A_n}(t_n), f_{A_0}(t_0)) &\leq H(f_{A_n}(t_n), f_{A_0}(t_n)) + H(f_{A_0}(t_n), f_{A_0}(t_0)) \\ &\leq H(A_n, A_0) + H(f_{A_0}(t_n), f_{A_0}(t_0)); \end{aligned}$$

let n tend to ∞ ; it follows from (III) and the hypothesis that we have

$$\lim_{n \rightarrow \infty} H(f_{A_n}(t_n), f_{A_0}(t_0)) = 0.$$

In the rest of this paper we assume that F is an (S) -convex structure on the metric space (X, d) .

LEMMA 3. For $0 \leq t \leq 1$, the mapping $x \rightarrow f_{\{x\}}(t)$ is nonexpansive from X into 2^X . Moreover, for all $A \in 2^X$ and $t \in [0, 1]$

$$\gamma(A) \geq \gamma(f_A(t)).$$

PROOF. Due to (II) it is easy to see that the mapping $x \rightarrow f_{\{x\}}(t)$ is nonexpansive.

Next, if $r > \gamma(A)$, by Remark 2 there is a finite set G such that $H(G, A) \leq r$. By (V), $f_G(t)$ is compact, thus for any $c > 0$ there is a finite set J with $H(J, f_G(t)) < c$. By the triangle inequality and (II) we have

$$H(J, f_A(t)) \leq H(J, f_G(t)) + H(f_G(t), f_A(t)) < c + r.$$

By Remark 2, $\gamma(f_A(t)) < c + r$ for all $c > 0$ and $r > \gamma(A)$. Hence

$$\gamma(f_A(t)) \leq \gamma(A).$$

LEMMA 4. For any $A \subset X$, let

$$g(A) = \bigcup \{f_A^-(t) \mid 0 \leq t \leq 1\}.$$

Then $\gamma(g(A)) = \gamma(A)$.

PROOF. It is due to (5) in Lemma 1 and the definition of g , we may assume that $A \in 2^X$. Define a real-valued function h on $[0, 1] \times [0, 1]$ as follows:

$$h(s, t) = H(f_A(s), f_A(t)).$$

It follows from condition (III) that h is a continuous function on $[0, 1] \times [0, 1]$, and thus h is uniformly continuous. In particular, for $\varepsilon > 0$ there is a positive number δ such that

$$h(s, t) < \varepsilon \quad \text{for all } 0 \leq s, t \leq 1 \text{ and } |s - t| < \delta.$$

Hence for all $0 \leq s, t \leq 1$ and $|s - t| < \delta$ we have $f_A(s) \subset (f_A(t))_\varepsilon$. Take $P = \{t_0 = 0 < t_1 < t_2 < \dots < t_n = 1\}$ be a partition of $[0, 1]$ with

$$t_i - t_{i-1} < \delta \quad \text{for all } i = 1, 2, \dots, n,$$

then $g(A) \subset \bigcup_{i=1}^n (f_A(t_i))_\varepsilon$.

From Lemmas 1 and 3, we have

$$\begin{aligned} \gamma(g(A)) &\leq \gamma\left(\bigcup_{i=1}^n (f_A(t_i))_\varepsilon\right) \\ &= \text{Max}\{\gamma((f_A(t_i))_\varepsilon) \mid i = 1, 2, \dots, n\} \\ &\leq \gamma(A) + \varepsilon. \end{aligned}$$

Therefore, $\gamma(g(A)) \leq \gamma(A)$. The fact that $A \subset g(A)$ implies

$$\gamma(g(A)) = \gamma(A).$$

REMARK 3. It follows from (IV) and the definition of g that for $A \subset B \subset X$ we have

$$f_A^-(t) \subset f_B^-(t) \quad \text{for all } t \in [0, 1]$$

and then

$$g(A) \subset g(B).$$

Now we are going to prove our theorem.

PROOF OF THE THEOREM. Let $x_0 \in X$. We denote O_{x_0} as the set

$$\cup \{(gT)^n x_0 \mid n = 0, 1, 2, \dots\}$$

where $(gT)^0 x_0 = \{x_0\}$ and $(gT)^n x_0 = g(T((gT)^{n-1} x_0))$ for $n = 1, 2, \dots$. Since $T(gT)^n x_0 \subset (gT)^{n+1} x_0$, we have

$$T(\overline{O_{x_0}}) \subset \overline{O_{x_0}} \quad \text{and} \quad \gamma(T(\overline{O_{x_0}})) \leq \gamma(\overline{O_{x_0}}).$$

Moreover, by (IV) and Remark 3 that

$$g(T(O_{x_0})) = g\left(\bigcup_{n=0}^{\infty} T(gT)^n x_0\right) \supset (gT)^{k+1} x_0$$

for all $k = 0, 1, 2, \dots$; hence

$$\{x_0\} \cup gT(O_{x_0}) \supset \{x_0\} \cup \left(\bigcup_{n=0}^{\infty} (gT)^{n+1} x_0\right) = O_{x_0}$$

and thus

$$\gamma(\overline{O_{x_0}}) = \gamma(O_{x_0}) \leq \gamma(gT(\overline{O_{x_0}})) = \gamma(T(\overline{O_{x_0}})).$$

That is,

$$\gamma(\overline{O_{x_0}}) = \gamma(T(\overline{O_{x_0}})).$$

Since T is a condensing mapping, we have that $\overline{O_{x_0}}$ is a compact subset of X . Now we let $\{r_n\}$ be a strictly increasing sequence of positive numbers with $\lim_{n \rightarrow \infty} r_n = 1$ and let $K = \overline{O_{x_0}}$ and

$$T_n x = f_{Tx}(r_n) \quad \text{for all } x \in K.$$

For $y \in O_{x_0}$ we have $y \in (gT)^k x_0$ for some $k = 0, 1, 2, \dots$. It follows from the definitions of T_n and O_{x_0} and Remark 3 that

$$T_n y = f_{Ty}(r_n) \subset g(Ty) \subset (gT)^{k+1} x_0 \subset O_{x_0}.$$

Then for any $x \in K$ there is a sequence $\{y_i\}$ in O_{x_0} which converges to x . By (II) we have

$$H(T_n y_i, T_n x) = H(f_{Ty_i}(r_n), f_{Tx}(r_n)) \leq H(Ty_i, Tx) \leq d(y_i, x);$$

hence $\lim_{i \rightarrow \infty} H(T_n y_i, T_n x) = 0$, or $T_n x \subset \overline{O_{x_0}} = K$, and thus T_n maps K into 2^K . Moreover, if x, y are in K we have

$$H(T_n x, T_n y) = H(f_{Tx}(r_n), f_{Ty}(r_n)) \leq \phi(r_n) H(Tx, Ty) \leq \phi(r_n) d(x, y);$$

that is, T_n is a contraction mapping of K into 2^K . It is due to a theorem of Nadler [3] that T_n has a fixed point x_n in K , or $x_n \in f_{Tx_n}(r_n)$. By the compactness of K there is a

convergent subsequence $\{x_{n_k}\}$ of $\{x_n\}$, with y_0 as its limit. Hence, by Lemma 2, we have that

$$\lim_{k \rightarrow \infty} H(f_{Tx_{n_k}}(r_{n_k}), f_{Ty_0}(1)) = 0$$

and therefore $y_0 \in f_{Ty_0}(1) = Ty_0$.

It follows from our theorem and Remark 1 that we have the following:

COROLLARY. *Let X be a starlike convex, bounded, closed subset of a Banach space E . Then every nonexpansive condensing multi-valued mapping of X into 2^X has a fixed point.*

REMARK 4. If X is compact then every continuous mapping of X into 2^X is condensing, hence Dotson's theorem and its Corollary 2 are special cases of our theorem and corollary, respectively.

ACKNOWLEDGMENT. The author wishes to express his appreciation to the referee who has made a lot of valuable suggestions for this paper.

REFERENCES

1. R. K. Bose and R. N. Mukherjee, *On fixed points of nonexpansive set-valued mappings*, Proc. Amer. Math. Soc. **72** (1978), 97–98.
2. W. G. Dotson, Jr., *On fixed points of nonexpansive mappings in nonconvex sets*, Proc. Amer. Math. Soc. **38** (1973), 155–156.
3. S. B. Nadler, Jr., *Multi-valued contraction mappings*, Pacific J. Math. **30** (1969), 473–488.
4. B. N. Sadovskii, *A fixed point principle*, Funktional. Anal. i Prilozhen. **1** (1967), 74–76.
5. ———, *Limit compact and condensing operators*, Russian Math. Surveys **127** (1972), 85–155.

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN NORMAL UNIVERSITY, TAIPEI, REPUBLIC OF CHINA