ON THE FIRST FACTOR OF THE CLASS NUMBER OF A CYCLOTOMIC FIELD

KEQIN FENG

ABSTRACT. Let p be an odd prime. $h_1(p)$ is the first factor of the class number of field $Q(\zeta_p)$. We proved that

$$h_1(p) \leq \begin{cases} 2p \left(\frac{p-1}{8(2^{l/2}+1)^{4/l}} \right)^{(p-1)/4}, & \text{if } l \text{ is even,} \\ 2p \left(\frac{p-1}{8(2^l-1)^{2/l}} \right)^{(p-1)/4}, & \text{if } l \text{ is odd.} \end{cases}$$

From that we obtain $h_1(p) \leq 2p((p-1)/31.997158\dots)^{(p-1)/4}$ which is better than Carlitz's and Metsänkylä's results. For the fields $Q(\zeta_{2^n})$ and $Q(\zeta_{p^n})$ ($n \geq 2$), we get the similar results.

Let p be an odd prime, $\zeta_p = e^{(2\pi i)/p}$, $h(p)$ and $h_2(p)$ be the class number of the cyclotomic field $Q(\zeta_p)$ and its maximal real subfield $Q(\zeta_p + \zeta_p^{-1})$ respectively. In 1850 Kummer [2] proved that $h_1(p) = h(p)/h_2(p)$ is a rational integer which is called the first factor of the class number of $Q(\zeta_p)$. Kummer conjectured that

$$h_1(p) \sim 2p \left(\frac{p}{4\pi^2} \right)^{(p-1)/4} = 2p \left(\frac{p}{39.4784\dots} \right)^{(p-1)/4} \quad (\text{when } p \rightarrow \infty).$$

In 1961 L. Carlitz [1] proved that $h_1(p) < (p-1)((p-1)/2)^{(p-1)/4}$. In 1974 T. Metsänkylä proved that $h_1(p) < 2p((p-1)(p-2)/24p)^{(p-1)/4}$. In this short paper I will show that

$$(1) \quad h_1(p) < 2p \left(\frac{p-1}{31.997158\dots} \right)^{(p-1)/4}.$$

Exactly speaking, I will show the following theorem.

THEOREM. Let l be the order of 2 (mod p). Then

$$h_1(p) \leq \begin{cases} 2p \left(\frac{p-1}{8(2^{l/2}+1)^{4/l}} \right)^{(p-1)/4}, & \text{if } l \text{ is even,} \\ 2p \left(\frac{p-1}{8(2^l-1)^{2/l}} \right)^{(p-1)/4}, & \text{if } l \text{ is odd.} \end{cases}$$

Received by the editors August 1, 1980.

1980 *Mathematics Subject Classification.* Primary 12A35, 12A50.

Key words and phrases. Class number, cyclotomic field.

© 1982 American Mathematical Society
0002-9939/81/0000-1087/\$02.00

FROM THE THEOREM TO (1). For even l , from the theorem we know that

$$h_1(p) \leq 2p \left(\frac{p-1}{8(2^{l/2} + 1)^{4/l}} \right)^{(p-1)/4} \leq 2p \left(\frac{p-1}{32} \right)^{(p-1)/4}.$$

For odd l , it is easy to see that if $p > 127$ then $l \geq 11$ and

$$(2^l - 1)^{2/l} \geq (2^{11} - 1)^{2/11} = 3.9996448 \dots$$

Thus

$$h_1(p) \leq 2p \left(\frac{p-1}{8 \times 3.9996448 \dots} \right)^{(p-1)/4} = 2p \left(\frac{p-1}{31.997158} \right)^{(p-1)/4}.$$

For $p \leq 127$ we can verify (1) directly.

To prove the theorem we need

LEMMA. Let χ be an odd Dirichlet character with the conductor f . If $f \not\equiv 2 \pmod{4}$, then

$$\sum_{1 \leq a < f} \chi(a)a = \frac{f}{\bar{\chi}(2) - 2} \sum_{1 \leq a < f/2} \chi(a).$$

PROOF. Long [3] proved this lemma for quadratic character χ . His method is available for any character satisfying the conditions of the lemma. We restate it for sake of completeness.

When $2 \nmid f$ then, on one hand,

$$\begin{aligned} - \sum_{0 < a < f} \chi(a)a &= - \sum_{\substack{0 < a < f \\ 2|a}} (\chi(a)a + \chi(f-a)(f-a)) \\ &= - \sum_{0 < a < f/2} \chi(2a)2a + \sum_{0 < a < f/2} \chi(2a)(f-2a) \\ &= \chi(2) \sum_{0 < a < f/2} (f-4a)\chi(a). \end{aligned}$$

Thus

$$(2) \quad -\bar{\chi}(2) \sum_{0 < a < f} \chi(a)a = \sum_{0 < a < f/2} (f-4a)\chi(a).$$

On the other hand,

$$\begin{aligned} - \sum_{0 < a < f} \chi(a)a &= - \sum_{0 < a < f/2} \chi(a)a - \sum_{0 < a < f/2} \chi(f-a)(f-a) \\ (3) \quad &= - \sum_{0 < a < f/2} \chi(a)a + \sum_{0 < a < f/2} \chi(a)(f-a) \\ &= \sum_{0 < a < f/2} (f-2a)\chi(a). \end{aligned}$$

From (2) and (3) we obtain

$$\sum_{0 < a < f} \chi(a)a = \frac{f}{\bar{\chi}(2) - 2} \sum_{0 < a < f/2} \chi(a).$$

When $4 \mid f$ then $\chi(2) = 0$ and $\chi(x + f/2) = -\chi(x)$. Thus

$$\begin{aligned} \sum_{0 < a < f} \chi(a)a &= \sum_{0 < a < f/2} \chi(a)a + \sum_{0 < a < f/2} \chi\left(a + \frac{f}{2}\right)\left(a + \frac{f}{2}\right) \\ &= \sum_{0 < a < f/2} \chi(a)a - \sum_{0 < a < f/2} \chi(a)\left(a + \frac{f}{2}\right) \\ &= -\frac{f}{2} \sum_{0 < a < f/2} \chi(a). \end{aligned}$$

PROOF OF THE THEOREM. Starting from the analytic formula of $h_1(p)$,

$$\begin{aligned} h_1(p) &= 2p \prod_{\chi(-1)=-1} \sum_{a=1}^{p-1} \left(-\frac{1}{2p}\right) \chi(a)a \\ &= (2p)^{-(p-1)/2+1} \prod_{\chi(-1)=1} \left| \sum_{a=1}^{p-1} \chi(a)a \right|, \end{aligned}$$

where χ is taken to be the $(p-1)/2$ odd characters of mod p . Let g be a primitive root of mod p and $\chi_0(a) = \exp(2\pi i)/(p-1) \cdot \text{ind}_g a$ ($1 \leq a \leq p-1$). Then

$$h_1(p) = (2p)^{-(p-1)/2+1} \prod_{\lambda=1}^{(p-1)/2} |\chi_0^{2\lambda-1}(a)a|.$$

From the above lemma we also have

$$(4) \quad h_1(p) = (2p) \cdot 2^{-(p-1)/2} \left(\prod_{\lambda=1}^{(p-1)/2} |2 - \chi_0^{2\lambda-1}(2)|^{-1} \right)^{(p-1)/2} \left| \sum_{a=1}^{(p-1)/2} \chi_0^{2\lambda-1}(a) \right|.$$

Using the orthogonal relations of odd characters of mod p ,

$$\begin{aligned} \sum_{\lambda=1}^{(p-1)/2} \chi_0^{2\lambda-1}(a) \bar{\chi}_0^{2\lambda-1}(b) &= \sum_{\lambda=1}^{(p-1)/2} \exp\left(\frac{2\pi i \cdot \text{ind}_g \left(\frac{a}{b}\right)}{p-1} \cdot (2\lambda-1)\right) \\ &= \begin{cases} (p-1)/2, & \text{if } a \equiv b \pmod{p}, \\ -(p-1)/2, & \text{if } a \equiv -b \pmod{p}, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

For $1 \leq a, b \leq p-1$, we have

$$\begin{aligned} \prod_{\lambda=1}^{(p-1)/2} \left| \sum_{a=1}^{(p-1)/2} \chi_0^{2\lambda-1}(a) \right|^{4/(p-1)} &= \prod_{\lambda=1}^{(p-1)/2} \left(\sum_{a,b=1}^{(p-1)/2} \chi_0^{2\lambda-1}(a) \bar{\chi}_0^{2\lambda-1}(b) \right)^{2/(p-1)} \\ &\leq \frac{2}{p-1} \sum_{\lambda=1}^{(p-1)/2} \sum_{a,b=1}^{(p-1)/2} \chi_0^{2\lambda-1}(a) \bar{\chi}_0^{2\lambda-1}(b) \\ &= \frac{2}{p-1} \sum_{a,b=1}^{(p-1)/2} \sum_{\lambda=1}^{(p-1)/2} \chi_0^{2\lambda-1}(a) \bar{\chi}_0^{2\lambda-1}(b) \\ &= \frac{2}{p-1} \cdot \frac{p-1}{2} \cdot \frac{p-1}{2} = \frac{p-1}{2}. \end{aligned}$$

Thus

$$(5) \quad \prod_{\lambda=1}^{(p-1)/2} \left| \sum_{a=1}^{(p-1)/2} \chi_0^{2\lambda-1}(a) \right| \leq \left(\frac{p-1}{2} \right)^{(p-1)/4}.$$

On the other hand, we can assume $\text{ind}_g 2 = (p-1)/l$ without loss of generality. So

$$\begin{aligned} \prod_{\lambda=1}^{(p-1)/2} (2 - \chi_0^{2\lambda-1}(2)) &= \prod_{\lambda=1}^{(p-1)/2} \left(2 - \exp\left(\frac{2\pi i \cdot \text{ind}_g 2}{p-1}(2\lambda-1)\right) \right) \\ &= \prod_{\lambda=1}^{(p-1)/2} \left(2 - \exp\left(\frac{2\pi i(2\lambda-1)}{l}\right) \right) \\ (6) \quad &= \begin{cases} \left(\prod_{\mu=0}^{l-1} (2 - e^{2\pi i \mu/l}) \right)^{(p-1)/2l} = (2^l - 1)^{(p-1)/2l}, & \text{if } l \text{ is odd,} \\ \left(\frac{\prod_{\mu=0}^{l-1} (2 - e^{2\pi i \mu/l})}{\prod_{\mu=0}^{l/2-1} (2 - e^{2\pi i \mu/(l/2)})} \right)^{(p-1)/l} = \left(\frac{2^l - 1}{2^{l/2} - 1} \right)^{(p-1)/l} \\ &= (2^{l/2} + 1)^{(p-1)/l}, & \text{if } l \text{ is even.} \end{cases} \end{aligned}$$

Combining (4), (5), and (6) we obtain the theorem.

REMARK. Let $h_1(p^n)$ be the first factor of class number of field $Q(\zeta_{p^n})$. Using the similar argument and the same lemma we can get that

(i) $h_1(2^n) \leq 2^{n+(n-6)2^{n-3}}$ ($n \geq 2$),

(ii) For odd prime p

$$h_1(p^n) \leq \begin{cases} 2p^n \left(\frac{\varphi(p^n)}{8(2^{l/2} + 1)^{4/l}} \right)^{\varphi(p^n)/4}, & \text{if } l \text{ is even,} \\ 2p^n \left(\frac{\varphi(p^n)}{8(2^l - 1)^{2/l}} \right)^{\varphi(p^n)/4}, & \text{if } l \text{ is odd,} \end{cases}$$

where l is the order of 2 (mod p^n).

REFERENCES

1. L. Carlitz, *A generalization of Maillet's determinant and a bound for the first factor of the class number*, Proc. Amer. Math. Soc. **12** (1961), 256–261.
2. E. E. Kummer, *Bestimmung der Anzahl nicht Äquivalenter Klassen für die aus λ -ten Wurzeln der Einheit gebildeten complexen Zählen und die ideale Faktoren derselben*, J. Reine Angew. Math. **40** (1850), 93–116.
3. R. Long, *Algebraic number theory*, Dekker, New York, 1977.
4. T. Metsänkylä, *Class number and μ -invariants of cyclotomic fields*, Proc. Amer. Math. Soc. **43** (1974), 199–200.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MARYLAND 20742

Current address: Department of Mathematics, Chinese University of Science and Technology, Hefei, Anhui, The People's Republic of China