DISTANCE BETWEEN NORMAL OPERATORS

V. S. SUNDER¹

ABSTRACT. Lidskii and Wielandt have proved independently that if A and B are selfadjoint operators on an n -dimensional space H , with eigenvalues $\{\alpha_k\}_{k=1}^n$ and $\{\beta_k\}_{k=1}^n$ respectively (counting multiplicity), then,

$$\|A - B\| \geq \min_{\sigma \in S_n} \|\text{diag}(\alpha_k - \beta_{\sigma(k)})\|$$

for any unitarily invariant norm on $L(H)$. In this note an example is given to show that this result is no longer true if A and B are only required to be normal (even unitary). It is also shown that the above inequality holds in the operator norm, if A is selfadjoint and B is skew-self-adjoint.

Introduction. In [2], Weyl proved the following: if A and B are Hermitian operators on a finite-dimensional Hilbert space, and if $\text{sp } A = \{\alpha_k\}_{k=1}^n$, $\text{sp } B = \{\beta_k\}_{k=1}^n$ (counting multiplicity), then

$$(*) \quad \|A - B\| \geq \min_{\sigma \in S_n} \|\text{diag}(\alpha_k - \beta_{\sigma(k)})\|.$$

(Here, $\text{sp } A$ denotes the spectrum of A , and $\text{diag}(\gamma_k)$ denotes any normal operator with spectrum $\{\gamma_k\}_{k=1}^n$.) The norm used above is the operator norm. It was later shown by Lidskii [1] and Wielandt [3] that $(*)$ is valid in any norm on $L(H)$ that is unitarily invariant. (Throughout this note, H denotes a finite-dimensional Hilbert space.)

In this short note, we first give an example to show that the analogue of Lidskii's theorem for general normal operators is false. In the example given, A is Hermitian and unitary, while B is skew-Hermitian and unitary. We show, however (cf. Theorem 2) that Weyl's theorem remains valid if A is Hermitian and B is skew-Hermitian.

EXAMPLE 1. Let

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

Then A is Hermitian, B is skew-Hermitian and $\text{sp } A = \{\alpha_1, \alpha_2\}$, $\text{sp } B = \{\beta_1, \beta_2\}$ where $\alpha_1 = 1$, $\alpha_2 = -1$, $\beta_1 = i$, $\beta_2 = -i$. (Note that A and B are unitary.) If $\|\cdot\|_p$ is the Schatten p -norm, then it is easily checked that

$$\|A - B\|_p = 2, \quad \|\text{diag}(\alpha_k - \beta_{\sigma(k)})\|_p = 2^{1/2+1/p},$$

for any σ in S_2 . Thus, if $1 \leq p < 2$,

$$\|A - B\|_p < \min_{\sigma \in S_2} \|\text{diag}(\alpha_k - \beta_{\sigma(k)})\|_p. \quad \square$$

Received by the editors May 11, 1981.

1980 *Mathematics Subject Classification*. Primary 15A60, 15A57, 65F35; Secondary 15A42, 47A55.

¹The author is currently a Visiting Fellow at the Tata Institute of Fundamental Research.

© 1982 American Mathematical Society
0002-9939/81/0000-1088/\$01.50

In the above example, the operator norm of $A - B$ is larger than $\min_{\sigma \in S_2} \|\text{diag}(\alpha_k - \beta_{\sigma(k)})\|$. That this is not a coincidence is the content of the next theorem.

THEOREM 2. Let $A = A^*$, $B = B^* \in L(H)$. Let $\text{sp } A = \{\alpha_k\}_{k=1}^n$, $\text{sp } B = \{\beta_k\}_{k=1}^n$. Assume the numbering to satisfy $|\alpha_1| \geq \cdots \geq |\alpha_n|$, $|\beta_1| \leq \cdots \leq |\beta_n|$. Then,

$$\|A + iB\| \geq \|\text{diag}(\alpha_k + i\beta_k)\| = \max_k |\alpha_k + i\beta_k|.$$

PROOF. Let $\{e_k\}_{k=1}^n$ and $\{f_k\}_{k=1}^n$ be orthonormal bases of H such that $Ae_k = \alpha_k e_k$ and $Bf_k = \beta_k f_k$ for all k . Let $M_k = \vee \{e_l: l \leq k\}$ and $N_k = \vee \{f_l: l \geq k\}$. Then, since $\dim M_k = k$ and $\dim N_k = n - k + 1$, it follows that $\dim(M_k \cap N_k) \geq 1$. Let x_k be a unit vector in $M_k \cap N_k$. The ordering of the α_k 's and the β_k 's ensures that $\|Ax_k\| \geq |\alpha_k|$, $\|Bx_k\| \geq |\beta_k|$. Hence,

$$\begin{aligned} |\alpha_k + i\beta_k|^2 &= |\alpha_k|^2 + |\beta_k|^2 \leq \|Ax_k\|^2 + \|Bx_k\|^2 \\ &= \frac{1}{2} [\|(A + iB)x_k\|^2 + \|(A - iB)x_k\|^2] \\ &\leq \frac{1}{2} [\|A + iB\|^2 + \|A - iB\|^2] = \|A + iB\|^2. \end{aligned}$$

(The first equality follows from the parallelogram identity, while the last equality is a consequence of the fact that $A - iB = (A + iB)^*$.) Hence,

$$\|A + iB\| \geq \max_k |\alpha_k + i\beta_k|. \quad \square$$

The following is just a reformulation of Theorem 2: if $T \in L(H)$, then there exists a normal operator N on H such that (i) the real and imaginary parts of N are, respectively, unitarily equivalent to the real and imaginary parts of T , and (ii) $\|N\| \leq \|T\|$.

BIBLIOGRAPHY

1. V. Lidskii, *The proper values of the sum and product of symmetric matrices*, Dokl. Akad. Nauk SSSR **75** (1950), 760–772.
2. H. Weyl, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen*, Math. Ann. **71** (1912), 441–479.
3. H. Wielandt, *An extremum property of sums of eigenvalues*, Proc. Amer. Math. Soc. **6** (1955), 106–110.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400 005, INDIA

Current address: Indian Statistical Institute, 7, S. J. S. Sansanwal Marg, New Delhi 110029, India