SPHERICAL HARMONICS GENERATING BOUNDED BIHARMONICS

BRADLEY BEAVER, LEO SARIO AND CECILIA WANG

ABSTRACT. Let $H^2B(R)$ be the family of bounded nonharmonic biharmonic functions on a Riemannian manifold R . On the punctured Euclidean N -space $E_0^N = \{x = (x^1, \dots, x^N) \mid 0 < |x| < \infty\}$, H^2B is void for $N > 3$, whereas for $N = 2, 3$, it is generated by certain fundamental spherical harmonics. It is also known that H^2B remains void on the Riemannian manifold E_α^N , $N > 3$, obtained by endowing E_0^N with the non-Euclidean metric $ds_\alpha = r^\alpha |dx|$, $\alpha \in \mathbf{R}$.

The purpose of the present paper is to show that the fundamental spherical harmonics continue generating $H^2B(E_\alpha^3)$, despite the distorting metric ds_α . An analogous result holds for E_α^2 .

Let $H^2B(R)$ be the family of bounded nonharmonic biharmonic functions on a Riemannian manifold R . It was shown in Sario and Wang [2] that on the punctured Euclidean N -space $E_0^N = \{x = (x^1, \dots, x^N) \mid 0 < |x| < \infty\}$, H^2B is void for $N > 3$, whereas for $N = 2, 3$, it is generated by certain fundamental spherical harmonics.

In Sario and Wang [3], it was proved that H^2B remains void on the Riemannian manifold E_α^N , $N > 3$, obtained by endowing E_0^N with the non-Euclidean metric $ds_\alpha = r^\alpha |dx|$, $\alpha \in \mathbf{R}$. The cases $N = 2, 3$ were left open.

The purpose of the present paper is to show that the fundamental spherical harmonics continue generating $H^2B(E_\alpha^3)$, despite the distorting metric ds_α . An analogous result holds for E_α^2 .

1.1. Consider the space

$$E_\alpha^N = \{x = (r, \theta^1, \dots, \theta^{N-1}) \mid 0 < r < \infty, ds_\alpha = r^\alpha |dx|, \alpha \in \mathbf{R}\}$$

with polar coordinates $(r, \theta) = (r, \theta^1, \dots, \theta^{N-1})$. We first state some facts from [3].

LEMMA 1. *There exist no H^2B functions on E_{-1}^N for $N \geq 2$.*

Thus when we are searching for generators of H^2B functions, we may ignore the case of $\alpha = -1$.

LEMMA 2. *For any $\alpha \neq -1$, and $N \geq 2$, every $h \in H(E_\alpha^N)$ has an expansion*

$$h = \sum_{n=1}^{\infty} \sum_{m=1}^{m_n} (a_{nm} r^{p_n} + b_{nm} r^{q_n}) S_{nm} + a\sigma(r) + b,$$

Received by the editors February 18, 1981.

1980 *Mathematics Subject Classification.* Primary 31A30, 31B30.

©1982 American Mathematical Society
0002-9939/81/0000-0794/\$02.50

where

$$\sigma(r) = \begin{cases} \log r, & \text{if } N = 2, \\ r^{-(N-2)(\alpha+1)}, & \text{if } N > 2. \end{cases}$$

LEMMA 3. Let u be a biharmonic function on E_α^N , $N \geq 2$, $\alpha \neq -1$, with

$$\Delta u = \sum_{n=1}^{\infty} \sum_{m=1}^{m_n} (a_{nm} r^{p_n} + b_{nm} r^{q_n}) S_{nm} + a\sigma(r) + b.$$

Then

$$\begin{aligned} u = & \frac{-1}{4(\alpha+1)} \left[\sum_{n \neq \mu} \sum_{m=1}^{m_n} \frac{1}{P_n} a_{nm} r^{p_n+2\alpha+2} S_{nm} + \sum_{n \neq \nu} \sum_{m=1}^{m_n} \frac{1}{Q_n} b_{nm} r^{q_n+2\alpha+2} S_{nm} \right] \\ & - \frac{1}{2(\alpha+1)} r^{(2-N/2)(\alpha+1)} \log r \cdot \left(\sum_{m=1}^{m_\mu} a_{\mu m} S_{\mu m} + \sum_{m=1}^{m_\nu} b_{\nu m} S_{\nu m} \right) \\ & + \sum_{n=1}^{\infty} \sum_{m=1}^{m_n} (c_{nm} r^{p_n} + d_{nm} r^{q_n}) S_{nm} + a\tau(r) + bs(r) + c\sigma(r) + d, \end{aligned}$$

where

$$\begin{aligned} p_n &= \frac{1}{2} \left\{ -(N-2)(\alpha+1) + [(N-2)^2(\alpha+1)^2 + 4n(n+N-2)]^{1/2} \right\}, \\ q_n &= \frac{1}{2} \left\{ -(N-2)(\alpha+1) - [(N-2)^2(\alpha+1)^2 + 4n(n+N-2)]^{1/2} \right\}, \\ P_n &= \frac{1}{2} N(\alpha+1) + p_n, \quad Q_n = \frac{1}{2} N(\alpha+1) + q_n, \end{aligned}$$

μ and ν are defined by $P_\mu = 0$ and $Q_\nu = 0$ respectively, $\sigma(r)$ is as above,

$$s(r) = \frac{-1}{2N(\alpha+1)^2} r^{2\alpha+2},$$

and

$$\tau(r) = \begin{cases} s(r) \left[\log r - \frac{1}{\alpha+1} \right], & N = 2, \\ -\frac{1}{2\alpha+2} \log r, & N = 4, \\ \frac{1}{2(N-4)(\alpha+1)^2} r^{-(N-4)(\alpha+1)}, & N \neq 2, 4. \end{cases}$$

1.2. Generators of H^2B -functions on E_α^2 . We claim

THEOREM 1. $H^2B(E_\alpha^2)$, where $\alpha = -1 \pm k/2$, $k \in \mathbb{Z}^+$, is generated by 1, $\sin k\theta$, and $\cos k\theta$.

PROOF. We shall establish the theorem for the case $\alpha = -1 - k/2$, the other case being entirely similar. Let $S_{n1} = \cos n\theta$ and $S_{n2} = \sin n\theta$. From Lemma 3 we calculate

$$\begin{aligned} p_n &= \frac{1}{2} \left\{ -(N-2)(\alpha+1) + [(N-2)^2(\alpha+1)^2 + 4n(n+N-2)]^{1/2} \right\} \\ &= \frac{1}{2} [(4n^2)]^{1/2} = n, \\ q_n &= \frac{1}{2} \left\{ -(N-2)(\alpha+1) - [(N-2)^2(\alpha+1)^2 + 4n(n+N-2)]^{1/2} \right\} \\ &= -\frac{1}{2} [(4n^2)]^{1/2} = -n. \end{aligned}$$

$P_n = 0$ implies $\frac{1}{2}N(\alpha+1) + p_n = 0$, which yields $n = \frac{k}{2}$. Thus $\mu = \frac{k}{2}$ if k is even. Since μ must be an integer, there is no μ term if k is odd.

$Q_n = 0$ gives $\frac{1}{2}N(\alpha+1) + q_n = 0$, whence $-n = \frac{k}{2}$, which is impossible, since both n and k are positive. Thus there is no ν term.

We may represent u by

$$\begin{aligned} u &= \sum_{n \neq k/2} \sum_{m=1}^2 A_{nm} r^{n-k} S_{nm} + \sum_{n=1}^{\infty} \sum_{m=1}^2 B_{nm} r^{-n-k} S_{nm} \\ &\quad + r^{-k/2} \log r \cdot \sum_{m=1}^2 a_{(k/2)m} S_{(k/2)m} + \sum_{n=1}^{\infty} \sum_{m=1}^2 (c_{nm} r^n + d_{nm} r^{-n}) S_{nm} \\ &\quad + a\tau(r) + bs(r) + c\sigma(r) + d. \end{aligned}$$

We wish to show that all coefficients are zero except possibly A_{k1} , A_{k2} and d .

Suppose $c_{nm} \neq 0$ for some (n, m) . Choose $\rho(r) \in C_0^\infty$, $\rho \geq 0$, $\text{supp } \rho \subset (0, 1)$. Define $\rho_t(r) = \rho(r-t)$ for $0 < t < \infty$, and $\varphi_t(r, \theta) = \rho_t S_{nm}$. Then $\text{supp } \varphi_t \subset (t, t+1)$, and by the orthogonality of the $\{S_{nm}\}$,

$$\begin{aligned} (u, \varphi_t) &= \int_{E_a^2} u \rho_t S_{nm} r^{-k-1} d\theta dr \\ &= c \int_t^{t+1} [A_{nm} r^{n-k} + B_{nm} r^{-n-k} \\ &\quad + a_{(k/2)m} r^{-k/2} \log r + c_{nm} r^n + d_{nm} r^{-n}] \rho_t r^{-k-1} dr, \end{aligned}$$

where the third term in the integrand appears only if $n = k/2$; also,

$$(1, |\varphi_t|) = c \int_t^{t+1} \rho_t r^{-k-1} dr,$$

the c 's denoting constants, not necessarily the same.

Certainly $\varphi_t \in L^1(E_a^2)$, and thus we have the inequality

$$|(u, \varphi_t)| \leq (\sup |u|)(1, |\varphi_t|).$$

But as $t \rightarrow \infty$, $(u, \varphi_t) \sim ct^{n-k-1}$ and $(1, |\varphi_t|) \sim ct^{-k-1}$, thus forcing $c_{nm} = 0$ for all n and m .

If $A_{nm} \neq 0$ for $n > k$, the same estimate with the same test function as in the first step gives, as $t \rightarrow \infty$,

$$(u, \varphi_t) \sim ct^{n-2k-1}, \quad (1, |\varphi_t|) = O(t^{-k-1}).$$

Therefore $A_{nm} = 0$ for $n > k$.

Now suppose $B_{nm} \neq 0$ for some (n, m) . Choose $\rho(r) \in C_0^\infty$, $\rho \geq 0$, $\text{supp } \rho \subset (\beta, \gamma)$, $0 < \beta < \gamma < 1$. Define $\rho_t(r) = \rho(r/t)$ for $0 < t < 1$, and $\varphi_t(r, \theta) = \rho_t S_{nm}$. Then

$$(u, \varphi_t) = c \int_{\beta t}^{\gamma t} [A_{nm} r^{n-k} + B_{nm} r^{-n-k} + a_{(k/2)m} r^{-k/2} \log r + d_{nm} r^{-n}] \rho_t r^{-k-1} dr,$$

where the first term in the integrand appears only if $n < k$, and the third only if $n = k/2$. On the other hand,

$$(1, |\varphi_t|) = c \int_{\beta t}^{\gamma t} \rho_t r^{-k-1} dr.$$

As $t \rightarrow 0$,

$$(u, \varphi_t) \sim ct^{-n-2k}, \quad (1, |\varphi_t|) = O(t^{-k}),$$

and again we have a contradiction to $\varphi_t \in L^1(E_a^2)$. Therefore, $B_{nm} = 0$ for all m and n .

Suppose $d_{mn} \neq 0$ for some $n \geq k$. Using the same test function as above we obtain

$$(u, \varphi_t) = c \int_{\beta t}^{\gamma t} d_{nm} r^{-n} \rho_t r^{-k-1} dr \sim ct^{-n-k}, \quad \text{as } t \rightarrow 0,$$

$$(1, |\varphi_t|) = c \int_{\beta t}^{\gamma t} \rho_t r^{-k-1} dr = O(t^{-k}), \quad \text{as } t \rightarrow 0.$$

A fortiori, $d_{nm} = 0$ for $n \geq k$.

We have reduced u to

$$\begin{aligned} u = & A_{k1} S_{k1} + A_{k2} S_{k2} + d + \sum_{\substack{n=1 \\ n \neq k/2}}^{k-1} \sum_{m=1}^2 A_{nm} r^{n-k} S_{nm} \\ & + r^{-k/2} \log r \sum_{m=1}^2 a_{(k/2)m} S_{(k/2)m} + \sum_{n=1}^{k-1} \sum_{m=1}^2 d_{nm} r^{-n} S_{nm} \\ & + a\tau(r) + b\sigma(r) + c\sigma(r). \end{aligned}$$

Since u is bounded, we have, on letting $r \rightarrow \infty$, $A_{nm} = d_{nm} = 0$ for $n = 1, \dots, k-1$, and $a_{(k/2)1} = a_{(k/2)2} = a = b = c = 0$. We infer that

$$u = A_{k1} S_{k1} + A_{k2} S_{k2} + d = A_{k1} \cos k\theta + A_{k2} \sin k\theta + d.$$

The proof of Theorem 1 is complete.

In the case $k = 2$, we have $\alpha = 0$, and therefore

COROLLARY [2]. $H^2 B(E_0^2)$ is generated by 1, $\sin 2\theta$, and $\cos 2\theta$.

1.3. Generators of H^2B functions on E_α^3 . We proceed to show

THEOREM 2. *If $\alpha = -1 \pm \sqrt{\frac{1}{2}k(k+1)}$, $k \in \mathbb{Z}^+$, then $H^2B(E_\alpha^3)$ is generated by 1 and $\{S_{km}\}_{m=1}^k$, the set of fundamental spherical harmonics of order k .*

PROOF. Again we will give the proof only for $\alpha = -1 - \sqrt{\frac{1}{2}k(k+1)}$, the other case being similar. We start with

$$\begin{aligned} p_n &= \frac{1}{2} \left\{ -(3-2)(\alpha+1) + \left[(3-2)^2(\alpha+1)^2 + 4n(n+3-2) \right]^{1/2} \right\}, \\ &= \frac{1}{4} \left\{ \sqrt{2k(k+1)} + \sqrt{2k(k+1) + 16n(n+1)} \right\}, \\ q_n &= \frac{1}{4} \left\{ \sqrt{2k(k+1)} - \sqrt{2k(k+1) + 16n(n+1)} \right\}. \end{aligned}$$

If $P_n = 0$, then $\frac{1}{2}N(\alpha+1) + p_n = 0$ or $n(n+1) = \frac{3}{8}k(k+1)$. For fixed k this is a quadratic in n which has at most one positive integral solution. If such a solution exists, we denote it by $n = \mu$. Observe that $\mu < k$.

If $Q_n = 0$, then

$$-\frac{3}{4}\sqrt{2k(k+1)} = -\frac{1}{4}\sqrt{2k(k+1)} + \frac{1}{4}\sqrt{2k(k+1) + 16n(n+1)},$$

which has no solution, so that there is no ν term in $u \in H^2B(E_\alpha^3)$.

Thus u has the representation

$$\begin{aligned} u &= \sum_{n \neq \mu} \sum_{m=1}^{m_n} A_{nm} r^{p_n+2\alpha+2} S_{nm} + \sum_{n=1}^{\infty} \sum_{m=1}^{m_n} B_{nm} r^{q_n+2\alpha+2} S_{nm} \\ &\quad + r^{(2-N/2)(\alpha+1)} \log r \cdot \sum_{m=1}^{\mu} a_{\mu m} S_{\mu m} + \sum_{n=1}^{\infty} \sum_{m=1}^{m_n} (c_{nm} r^{p_n} + d_{nm} r^{q_n}) S_{nm} \\ &\quad + a\tau(r) + bs(r) + c\sigma(r) + d. \end{aligned}$$

The remainder of the proof will be divided into Steps I–V.

I. For any given n , the highest order term is obviously $c_{nm} r^{p_n}$; we eliminate this first.

Choose $\rho \in C_0^\infty$, $\rho \geq 0$, $\text{supp } \rho \subset (0, 1)$. Define $\rho_t(r) = \rho(r-t)$, and $\varphi_t = \rho_t S_{nm}$. Then

$$\begin{aligned} (u, \varphi_t) &= c \int_t^{t+1} \left[A_{nm} r^{p_n+2\alpha+2} + B_{nm} r^{q_n+2\alpha+2} \right. \\ &\quad \left. + a_{\mu m} r^{-1/4(2k(k+1))^{1/2}} \log r + c_{nm} r^{p_n} + d_{nm} r^{q_n} \right] \\ &\quad \cdot \rho_t r^{-1-3/2(2k(k+1))^{1/2}} dr, \\ (1, |\varphi_t|) &= c \int_t^{t+1} \rho_t r^{-1-3/2(2k(k+1))^{1/2}} dr = O(t^{-1-3/2(2k(k+1))^{1/2}}), \end{aligned}$$

whereas

$$(u, \varphi_t) \sim ct^{p_n-1-3/2(2k(k+1))^{1/2}}, \quad \text{as } t \rightarrow \infty.$$

Since $p_n > 0$ for all n , it follows that $c_{nm} = 0$ for all n and m .

II. For $n > k$, $p_n + 2\alpha + 2 > 0$, $p_n + 2\alpha + 2 > q_n$, and $p_n + 2\alpha + 2 > \mu$. After Step I above, the latter statements assure us that, for a fixed $n > k$, $A_{nm}r^{p_n+2\alpha+2}$ is the dominant term as $r \rightarrow \infty$. If $A_{nm} \neq 0$ for $n > k$, the test function from Step I gives

$$(u, \varphi_t) = c \int_t^{t+1} [A_{nm}r^{p_n+2\alpha+2} + \dots] \rho_t r^{-1-3/2(2k(k+1))^{1/2}} dr \\ \sim ct^{p_n+2\alpha+2-1-3/2(2k(k+1))^{1/2}}, \quad \text{as } t \rightarrow \infty.$$

Thus $A_{nm} = 0$ for $n > k$.

III. Suppose $B_{nm} \neq 0$ for some n . Choose $\rho \in C_0^\infty$, $\rho \geq 0$, $\text{supp } \rho \subset (\beta, \gamma)$, $0 < \beta < \gamma < 1$. Define $\rho_t(r) = \rho(r/t)$, $\varphi_t = \rho_t S_{nm}$. Then

$$(u, \varphi_t) = c \int_{\beta t}^{\gamma t} [B_{nm}r^{q_n+2\alpha+2} + \dots] \rho_t r^{-1-3/2(2k(k+1))^{1/2}} dr \\ \sim ct^{q_n+2\alpha+2-3/2(2k(k+1))^{1/2}}, \quad \text{as } t \rightarrow 0, \\ (1, |\varphi_t|) = c \int_{\beta t}^{\gamma t} \rho_t r^{-1-3/2(2k(k+1))^{1/2}} dr = O(t^{-3/2(2k(k+1))^{1/2}}), \quad \text{as } t \rightarrow 0.$$

The relation $q_n + 2\alpha + 2 < 0$ for all n implies $B_{nm} = 0$ for all n and m .

IV. As in Step II, assume $n > k$. In view of the above, the test function from Step III gives

$$(u, \varphi_t) = c \int_{\beta t}^{\gamma t} d_{nm} r^{q_n} \rho_t r^{-1-3/2(2k(k+1))^{1/2}} dr \sim ct^{q_n-3/2(2k(k+1))^{1/2}}$$

and

$$(1, |\varphi_t|) = c \int_{\beta t}^{\gamma t} \rho_t r^{-1-3/2(2k(k+1))^{1/2}} dr = O(t^{-3/2(2k(k+1))^{1/2}}), \quad \text{as } t \rightarrow 0.$$

Inasmuch as $q_n < 0$ for all n , $d_n = 0$ for all $n > k$.

V. We have now reduced u to

$$u = \sum_{\substack{n=1 \\ n \neq \mu}}^k \sum_{m=1}^{m_n} A_{nm} r^{p_n+2\alpha+2} S_{nm} + r^{-1/4(2k(k+1))^{1/2}} \log r \cdot \sum_{m=1}^{m_\mu} a_{\mu m} S_{\mu m} \\ + \sum_{n=1}^k \sum_{m=1}^{m_n} d_{nm} r^{q_n} S_{nm} + a\tau(r) + bs(r) + c\sigma(r) + d.$$

As in the proof of Theorem 1, we may eliminate a finite number of linearly independent unbounded terms and are left with

$$u = \sum_{m=1}^{m_k} A_{km} S_{km} + d.$$

The proof of Theorem 2 is complete.

In particular, when $k = 1$, we have $\alpha = 0$, and the S_{1m} are $\cos \theta \cos \psi$, $\sin \theta \sin \psi$ and $\cos \theta$, where θ is the angle between the vector and the z -axis.

COROLLARY [2]. $HB^2(E_0^3)$ is generated by 1, $\cos \theta \cos \psi$, $\sin \theta \sin \psi$, and $\cos \theta$.

REFERENCES

1. C. Müller, *Spherical harmonics*, Lecture Notes in Math., vol. 17, Springer-Verlag, Berlin-Heidelberg-New York, 1966.
2. L. Sario and C. Wang, *Generators of the space of bounded biharmonic functions*, Math. Z. **127** (1972), 273–280.
3. ———, *Riemannian manifolds of dimension $N \geq 4$ without bounded biharmonic functions*, J. London Math. Soc. (2) **7** (1974), 635–644.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA 90024

DEPARTMENT OF MATHEMATICS, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA 85281