COMBINATORIAL EQUIVALENCE BETWEEN GROUP PRESENTATIONS

SUSHIL JAJODIA

ABSTRACT. Let

$$\mathcal{P} = (x_1, \dots, x_n: W_1, \dots, W_m) \quad \text{and} \quad \mathcal{R} = (x_1, \dots, x_n: R_1, \dots, R_m)$$

be two presentations, with the same generators, for a group π . In this note, we give a necessary and sufficient criterion which insures the existence of a combinatorial equivalence between $\mathcal{P}$ and $\mathcal{R}$ requiring only replacement operations.

I. Introduction. Let

$$(1) \quad \mathcal{P} = (x_1, \dots, x_n: W_1, \dots, W_m)$$

and

$$(2) \quad \mathcal{R} = (x_1, \dots, x_n: R_1, \dots, R_m)$$

be two presentations, with the same generators, for a group π (i.e., the normal closures of $W_1, \dots, W_m$ and of $R_1, \dots, R_m$ in the free group F on $x_1, \dots, x_n$ coincide). In [8, p. 95], A. J. Sieradski makes the following conjecture: Suppose there exists a combinatorial equivalence between $\mathcal{P}$ and $\mathcal{R}$ inducing the identity isomorphism. Then there exists a combinatorial equivalence between $\mathcal{P}$ and $\mathcal{R}$ requiring only replacement operations. We are unable to prove this conjecture; however, we give a necessary and sufficient criterion which insures its existence.

II. Definition of combinatorial equivalence. Let π be any group, and let

$$\mathcal{R} = (x_1, \dots, x_n: R_1, \dots, R_m) = (x_i: R_j)$$

be a finite presentation for π . We have the following *combinatorial operations* on the presentation $\mathcal{R}$:

(1) *The expansion operation.* It appends to the presentation $\mathcal{R}$ a new generator $x \notin \{x_i\}$ and a relator Wx^{-1} where W is an element in the free group F on generators $x_1, \dots, x_n$. Therefore

$$(x_i: R_j) \rightarrow (x_i, x: R_j, Wx^{-1}).$$

(2) *The contraction operation.* It simultaneously deletes a generator x and a relator Wx^{-1} provided that W and the remaining relators R_j are elements in the free group F generated by the remaining generators $x_1, \dots, x_n$. Therefore $(x_i, x: R_j, Wx^{-1}) \rightarrow (x_i: R_j)$.

(3) *The replacement operation.* It replaces a single relator R by a new relator S provided that S and $R^{\pm 1}$ are conjugates modulo the normal subgroup $N(R_j)$ of F

Received by the editors June 10, 1981.

1980 *Mathematics Subject Classification.* Primary 20E06.

© 1982 American Mathematical Society
0002-9939/82/0000-0072/\$01.75

generated by the other relators R_j which are kept unchanged. Hence $(x_i: R_j, R) \rightarrow (x_i: R_j, S)$.

Let $\mathcal{P}$ and $\mathcal{R}$ be two presentations for the same group π . We say $\mathcal{P}$ and $\mathcal{R}$ are *combinatorially equivalent* if there is a sequence

$$\mathcal{P} = \mathcal{P}_0 \rightarrow \mathcal{P}_1 \rightarrow \cdots \rightarrow \mathcal{P}_{s-1} \rightarrow \mathcal{P}_s = \mathcal{R}$$

of combinatorial operations which begin with $\mathcal{P}$ and end with $\mathcal{R}$.

Note that we do not permit the Tietze operation of adding trivial relators to the presentation. This restriction is motivated by topological considerations. We urge the reader to read [8] for details.

III. Necessary and sufficient criterion. Suppose the presentations (1) and (2), with the same deficiency and the same generators, present the same group π . We will now formulate the necessary and sufficient conditions.

Let F denote the free group generated by $x_1, \dots, x_n$ and R the normal closure in F of $R_1, \dots, R_m$. Let $\bar{F}$ denote the free group on symbols $r_1, \dots, r_m$. Since each $W_i \in R$, we can write

$$W_i = \prod_{k=1}^{m_i} (x_{ik} R_{ik} x_{ik}^{-1})^{\epsilon_{ik}}$$

where $R_{ik} = R_j$ for some j , $1 \leq j \leq m$, $x_{ik} \in F$, and $\epsilon_{ik} = \pm 1$. Let

$$w_i = \prod_{k=1}^{m_i} (x_{ik} r_{ik} x_{ik}^{-1})^{\epsilon_{ik}}$$

where if $R_{ik} = R_j$ in W_i , then $r_{ik} = r_j$ in w_i .

Let J denote the $m \times m$ matrix

$$J = \begin{vmatrix} \frac{\partial w_1}{\partial r_1} & \cdots & \frac{\partial w_1}{\partial r_m} \\ \vdots & & \vdots \\ \frac{\partial w_m}{\partial r_1} & \cdots & \frac{\partial w_m}{\partial r_m} \end{vmatrix}$$

where $\partial/\partial r_j: ZF \rightarrow ZF$ denotes the j th free partial derivative [2].

THEOREM. *Notation as above. Then the following are equivalent:*

- (1) *There is a combinatorial equivalence between $\mathcal{P}$ and $\mathcal{R}$ requiring only replacement operations.*
- (2) *$\{x_1, \dots, x_n, w_1, \dots, w_m\}$ forms a generating set for the free product $F * \bar{F}$.*
- (3) *J has a right inverse.*

PROOF. (1) $\Rightarrow$ (2) Suppose $\mathcal{P}$ is combinatorially equivalent to $\mathcal{R}$ using only replacement operations. Now all replacement operations are composites of these three replacement operations [8, p. 69]:

- (a) Replace a single relator R_i by its inverse R_i^{-1} .
- (b) Replace a relator R_i by the product $R_i R_j$ or $R_j R_i$ with a different relator R_j .
- (c) Replace a relator R_i by $x^\epsilon R_i x^{-\epsilon}$ by a generator x , $\epsilon = \pm 1$.

This means that we can convert $x_1, \dots, x_n, r_1, \dots, r_m$ into $x_1, \dots, x_n, w_1, \dots, w_m$ using transformations which leave $x_1, \dots, x_n$ fixed and on the remaining generators the following operations:

(a') Replace r by r' .

(b') Replace r by $r\bar{r}$ or $\bar{r}r$ where $\bar{r}$ is a different generator.

(c') Replace r by $x^\epsilon r x^{-\epsilon}$ where $x = x_i, 1 \leq i \leq n$.

But these are the elementary Nielsen transformations [3, p. 130]. Hence $x_1, \dots, x_n, w_1, \dots, w_m$ forms a generating set for $F * \bar{F}$.

(2) $\Rightarrow$ (1) Suppose $\{x_1, \dots, x_n, w_1, \dots, w_m\}$ generates $F * \bar{F}$. Then by a result of Rapaport [7, §5], it follows that one can obtain $x_1, \dots, x_n, w_1, \dots, w_m$ from $x_1, \dots, x_n, r_1, \dots, r_m$ using transformations which leave $x_1, \dots, x_n$ fixed and on the remaining generators the operations (a'), (b'), and (c') described above. But then one can construct a combinatorial equivalence between $\mathcal{P}$ and $\mathcal{R}$ using only replacement operations.

(2) $\Rightarrow$ (3) Suppose $\{x_1, \dots, x_n, w_1, \dots, w_m\}$ forms a generating set for $F * \bar{F}$. Then by the Inverse Function Theorem [1, p. 635], the Jacobian which has the form

$$\begin{vmatrix} I_n & 0 \\ A & J \end{vmatrix}$$

has a right inverse B . By a result of Kaplansky [6], B is a two-sided inverse. Therefore J has a right inverse.

(3) $\Rightarrow$ (2) Suppose J has a right inverse H . Then the Jacobian which has the form

$$\begin{vmatrix} I_n & 0_m \\ A_{m \times n} & J_m \end{vmatrix}$$

has a right inverse

$$\begin{vmatrix} I_n & 0_m \\ -H_m A_{m \times n} & H_m \end{vmatrix}.$$

The result now follows from the Inverse Function Theorem [1, p. 635]. This completes the proof.

EXAMPLE. Consider the two presentations

$$\mathcal{P} = (x, y, z: W_1 = x^5, W_2 = y^5, W_3 = z^5, W_r = [x, y], W_5 = [x, z], W_6 = [y, z])$$

and

$$\mathcal{R} = (x, y, z: R_1 = x^5, R_2 = y^5, R_3 = z^5, R_4 = [x^4, y], R_5 = [x, z], R_6 = [y, z])$$

of the group $Z_5 \times Z_5 \times Z_5$. We can write

$$W_4 = xyR_1^{-1}y^{-1}R_4^{-1}R_1x^{-1}.$$

Therefore, $w_4 = xyr^{-1}y^{-1}r_4^{-1}r_1x^{-1}$.

Clearly $\{x, y, z, w_1, w_2, w_3, w_4, w_5\}$ forms a generating set for $F * \bar{F}$ and the matrix J is invertible. Thus there is a combinatorial equivalence between $\mathcal{P}$ and $\mathcal{R}$ using only replacement operations.

REMARK. In [4 and 5], examples are given of presentations which are combinatorially equivalent, but there does not exist a combinatorial equivalence between them requiring replacement operations only; however, these presentations do not have the same generators.

REFERENCES

1. J. S. Birman, *An inverse function theorem for free groups*, Proc. Amer. Math. Soc. **41** (1973), 634–638.
2. R. H. Fox, *Free differential calculus. I. Derivation in the free group ring*, Ann. of Math. (2) **57** (1953), 547–560.
3. W. Magnus, A. Karrass and D. Solitar, *Combinatorial group theory*, Interscience, New York, 1966.
4. J. McCool and A. Pietrowski, *On free products with amalgamation of two infinite cyclic groups*, J. Algebra **18** (1971), 377–383.
5. W. Metzler, *Über den homotopietyp zweidimensionaler CW-Komplexe und Elementartransformationen bei definierende relatumen*, J. Reine Angew. Math. **285** (1976), 7–23.
6. M. S. Montgomery, *Left and right inverses in group algebras*, Bull. Amer. Math. Soc. **75** (1969), 539–540.
7. E. S. Rapaport, *Groups of order 1: Some properties of presentations*, Acta Math. **121** (1968), 127–150.
8. A. J. Sieradski, *Combinatorial isomorphisms and combinatorial homotopy equivalences*, J. Pure Appl. Algebra **7** (1976), 59–95.

COMPUTER SCIENCE DEPARTMENT, 304 MATHEMATICAL SCIENCES BUILDING, UNIVERSITY OF MISSOURI-COLUMBIA, COLUMBIA, MISSOURI 65211