CLEAR VISIBILITY AND THE DIMENSION OF KERNELS OF STARSHAPED SETS

MARILYN BREEN

ABSTRACT. This paper will use the concept of clearly visible to obtain a Krasnosel'skii-type theorem for the dimension of the kernel of a starshaped set, and the following result will be proved: For each k and n , $1 \leq k \leq n$, let $f(n, n) = n + 1$ and $f(n, k) = 2n$ if $1 \leq k \leq n - 1$. Let S be a nonempty compact set in R^n . Then for a k with $1 \leq k \leq n$, $\dim \ker S \geq k$ if and only if every $f(n, k)$ points of $\text{bdry } S$ are clearly visible from a common k -dimensional subset of S . If $k = 1$ or $k = n$, the result is best possible. Moreover, if S is a compact, connected, nonconvex set in R^2 , then $\text{bdry } S$ may be replaced by $\text{inc } S$ in the theorem.

1. Introduction. We begin with some definitions from [1]. Let S be a compact set in R^n . A point s in S is said to be a *point of local convexity* of S if and only if there is some neighborhood N of s such that $N \cap S$ is convex. If S fails to be locally convex at q in S , then q is called a *point of local nonconvexity* (inc point) of S . For points x and y in S , we say x *sees* y *via* S (x is *visible* from y *via* S) if and only if the segment $[x, y]$ lies in S . Similarly, x is *clearly visible* from y *via* S if and only if there is some neighborhood N of x such that y sees each point of $N \cap S$ *via* S . Finally, set S is *starshaped* if and only if there is some point p in S such that p sees each point of S *via* S , and the set of all such points p is called the (convex) *kernel* of S , denoted $\ker S$.

A theorem of Krasnosel'skii [6] states that if S is a nonempty compact set in R^n , then S is starshaped if and only if every $n + 1$ points of S are visible from a common point of S . (A stronger result may be obtained by replacing points of S with boundary points of S .) In [2], an analogue of the Krasnosel'skii theorem was proved for the dimension of this kernel: For each k and n , $1 \leq k \leq n$, let $f(n, n) = n + 1$ and $f(n, k) = 2n$ if $1 \leq k \leq n - 1$. If S is a compact set in R^n , then $\ker S$ has dimension at least k if and only if for some $\epsilon > 0$, every $f(n, k)$ points of S see *via* S a common k -dimensional ϵ -neighborhood. Unfortunately, the uniform lower bound ϵ is necessary by an example in [3].

In order to obtain a theorem for the dimension of the kernel independent of this cumbersome ϵ -bound, we turn to the notion of clearly visible, previously appearing in work by Stavrakas [8] and Falconer [4]. In [1], analogues of the Krasnosel'skii theorem were obtained by replacing the concept of visible with that of clearly visible and by replacing points of S with inc points of S . A similar approach proves helpful here, and we have the following result: For $f(n, k)$ defined above, $1 \leq k \leq n$, and for S a nonempty compact set in R^n , $\ker S$ has dimension at least k if and only if every $f(n, k)$ boundary points of S are clearly visible from a common k -dimensional

Received by the editors October 16, 1981.

1980 *Mathematics Subject Classification*. Primary 52A30, 52A35; Secondary 52A20.

Key words and phrases. Starshaped sets, convex kernel.

© 1982 American Mathematical Society
0002-9939/82/0000-0088/\$02.25

subset of S . If $k = 1$ or $k = n$, the result is best possible. Moreover, if S is a compact, connected, nonconvex set in R^2 , then the boundary of S may be replaced by the lnc points of S .

The following terminology will be used: $\text{conv } S$, $\text{bdry } S$, and $\text{ker } S$ will denote the convex hull, boundary, and kernel, respectively, for set S . $\text{lnc } S$ will be the set of points of local nonconvexity of S , and if S is convex, $\dim S$ will be the dimension of S . Finally, σ will represent the Hausdorff metric defined on the collection of compact, convex subsets of R^n . The reader is referred to Valentine [9] for a discussion of these concepts and to Nadler [7] for further information on the Hausdorff metric.

2. The results. The first lemma is a variation of a result in [2].

LEMMA 1. *For each k and n , $1 \leq k \leq n$, let $f(n, n) = n + 1$ and $f(n, k) = 2n$ if $1 \leq k \leq n - 1$. Let $\mathcal{B}$ be a uniformly bounded collection of compact convex sets in R^n which is closed with respect to the Hausdorff metric. Then for a k with $1 \leq k \leq n$, $\dim \bigcap \{B : B \text{ in } \mathcal{B}\} \geq k$ if and only if every $f(n, k)$ members of $\mathcal{B}$ contain a common k -dimensional set.*

PROOF. We need only establish the sufficiency of the condition. Observe that if every $f(n, k)$ members of $\mathcal{B}$ contain a common k -dimensional set, then by Helly's theorem in R^n , $\bigcap \{B : B \text{ in } \mathcal{B}\} \neq \emptyset$. If $k = n$, suppose on the contrary that $\dim \bigcap \{B : B \text{ in } \mathcal{B}\} < n$. Then by a theorem of Falconer [4, Theorem 1], there exist r sets $B_1, \dots, B_r$ in $\mathcal{B}$ such that $\dim(B_1 \cap \dots \cap B_r) = q$, where $q < n$ and $r \leq 2(n - q)$. However, by a result of Katchalski [5, Theorem a], every finite subfamily of $\mathcal{B}$ has an n -dimensional intersection. We have a contradiction, our supposition is false, and the result is established when $k = n$.

To establish the result for arbitrary k , $1 \leq k \leq n$, we adapt an argument from [2, Lemma]. The inductive proof is sketched below. If $n = 1$, then $k = 1$, and we have our result. Assume the result is true for natural numbers less than n , $2 \leq n$. If $k = n$, again the proof is immediate, so assume $1 \leq k < n$. If $\dim \bigcap \{B : B \text{ in } \mathcal{B}\} = n$, there is nothing to prove, so suppose that $\dim \bigcap \{B : B \text{ in } \mathcal{B}\} < n$. Then by Falconer's result, there are r sets $B_1, \dots, B_r$ in $\mathcal{B}$ with $\dim(B_1 \cap \dots \cap B_r) = q$, for some $q < n$ and $r \leq 2(n - q)$. Using Katchalski's theorem, $q \geq k$.

Let G denote the q -dimensional flat in R^n determined by $B_1 \cap \dots \cap B_r$. Using our hypothesis and the fact that $1 \leq k \leq q < n$, it is not hard to show that every $2q$ sets in $\mathcal{B}$ meet in a k -dimensional set in G . Hence every $f(q, k)$ members of $\{B \cap G : B \text{ in } \mathcal{B}\}$ meet in a k -dimensional set, and by our induction hypothesis, $\dim \bigcap \{B \cap G : B \text{ in } \mathcal{B}\} \geq k$, finishing the induction and completing the proof of the lemma.

The following definitions will be helpful.

DEFINITION. Let S be a compact set in R^n and let $q \in \text{bdry } S$. We define

$$S_q \equiv \{x : [x, q] \subseteq S\},$$

$$A_q \equiv \{x : q \text{ is clearly visible from } x \text{ via } S\},$$

and

$$C_q \equiv \bigcap \{H : H \text{ a closed halfspace with } A_q \subseteq H \text{ and } q \in \text{bdry } H\}.$$

Using these definitions, we establish two more lemmas.

LEMMA 2. Let S be a compact set in R^n . If B is in the σ -closure of $\mathcal{C} \equiv \{C_q \cap \text{conv } S : q \in \text{bdry } S\}$, then B contains some set A_q .

PROOF. Let $\{C_{q_n} \cap \text{conv } S : n \geq 1\}$ be a sequence in $\mathcal{C}$ converging to B . Then since S is bounded, some subsequence $\{q'_n\}$ of $\{q_n\}$ converges to a point q , and clearly $q \in \text{bdry } S$. Since $\{C_{q'_n} \cap \text{conv } S\}$ also converges to B , for convenience of notation we may assume that $\{q_n\}$ converges to q . We assert that $A_q \subseteq B$ for this particular q . Let $x \in A_q$. Then q is clearly visible via S from x , so for some neighborhood N of q , x sees via S each point of $N \cap S$. There is some integer M such that whenever $n > M$, $q_n \in N$. Hence for $n > M$, q_n is clearly visible via S from x , and so $x \in A_{q_n} \subseteq C_{q_n} \cap \text{conv } S$ for $n > M$. Since $\{C_{q_n} \cap \text{conv } S : n > M\}$ converges to B , $x \in B$. Thus $A_q \subseteq B$ and the lemma is established.

COROLLARY. Let S be a compact set in R^n , and let k and j be fixed, $0 \leq k \leq n$. If every j sets in $\{A_q : q \in \text{bdry } S\}$ have at least a k -dimensional intersection, then every j sets in the σ -closure of $\mathcal{C} \equiv \{C_q \cap \text{conv } S : q \in \text{bdry } S\}$ have at least a k -dimensional intersection.

LEMMA 3. Let S be a compact set in R^n , and let $\mathcal{C} \equiv \{C_q \cap \text{conv } S : q \in \text{bdry } S\}$. Then $\ker S = \bigcap \{B : B \text{ in the } \sigma\text{-closure of } \mathcal{C}\}$.

PROOF. For convenience, let $\mathcal{B}$ denote the σ -closure of $\mathcal{C}$. Using Lemma 2, clearly $\ker S \subseteq \bigcap \{A_q : q \in \text{bdry } S\} \subseteq \bigcap \{B : B \text{ in } \mathcal{B}\}$, so we need to establish only the reverse inclusion. Let x belong to $R^n \sim \ker S$. By Krasnosel'skii's Lemma [9, Lemma 6.2], there is a z in $\text{bdry } S$ and a closed halfspace H such that $z \in \text{bdry } H$, $S_z \subseteq H$, and $x \notin H$. Since $A_z \subseteq S_z$, we have $A_z \subseteq H$, $x \notin C_z$, and $x \notin \bigcap \{B : B \text{ in } \mathcal{B}\}$. Thus the lemma is proved.

THEOREM 1. For each k and n , $1 \leq k \leq n$, let $f(n, n) = n + 1$ and $f(n, k) = 2n$ if $1 \leq k \leq n - 1$. Let S be a nonempty compact set in R^n . Then for a k with $1 \leq k \leq n$, $\dim \ker S \geq k$ if and only if every $f(n, k)$ points of $\text{bdry } S$ are clearly visible from a common k -dimensional subset of S . If $k = 1$ or $k = n$, the result is best possible.

PROOF. The necessity of the condition is obvious. To establish its sufficiency, assume that every $f(n, k)$ points of $\text{bdry } S$ are clearly visible from a common k -dimensional subset of S . Using our hypothesis, every $f(n, k)$ members of $\{A_q : q \in \text{bdry } S\}$ have at least a k -dimensional intersection. Therefore, by the corollary to Lemma 2, every $f(n, k)$ sets in the σ -closure $\mathcal{B}$ of $\mathcal{C} \equiv \{C_q \cap \text{conv } S : q \in \text{bdry } S\}$ have at least a k -dimensional intersection. Standard arguments involving the Hausdorff metric [7, 9] show that $\mathcal{B}$ is a uniformly bounded collection of compact convex sets in R^n . Hence by Lemma 1, $\dim \bigcap \{B : B \text{ in } \mathcal{B}\} \geq k$. Using Lemma 3, $\ker S = \bigcap \{B : B \text{ in } \mathcal{B}\}$, so $\dim \ker S \geq k$, and the theorem is proved.

To see that the number $f(n, 1) = 2n$ is best possible, consider the following example, adapted from a construction by Katchalski [5, Theorem b, Case 1].

EXAMPLE 1. For convenience of notation, use the component representation $(z_1, \dots, z_n)$ for a point in R^n , and let D denote the closed unit ball centered at the origin. Construct a compact set S in R^n such that $\text{Int } S$ is a union of $2n$ disjoint convex sets $K_1, \dots, K_{2n}$, each of dimension $n - 2$, and having the following properties:

- (1) For x, y in K_i , $C_x = C_y$.

(2) For x in K_i , $1 \leq i \leq n$, $D \cap \{(z_1, \dots, z_n): z_i \geq 0\} \subseteq A_x$ and $A_x \cap \{(z_1, \dots, z_n): z_i < 0\} = \emptyset$.

For x in K_{i+n} , $1 \leq i \leq n$, $D \cap \{(z_1, \dots, z_n): z_i \leq 0\} \subseteq A_x$ and $A_x \cap \{(z_1, \dots, z_n): z_i > 0\} = \emptyset$.

(3) $K_i \cap K_{i+n} \subseteq \{(z_1, \dots, z_n): z_i = 0\}$, $1 \leq i \leq n$.

Then for $q_i \in K_i$, $1 \leq i \leq 2n$, the corresponding sets A_{q_i} have only the origin in common. However, every $2n - 1$ of the sets in $\{A_q: q \in \text{bdry } S\}$ share a common interval in D . Thus the number $f(n, 1) = 2n$ is best possible.

Figure 1 below illustrates the example in R^2 .

FIGURE 1

Similarly, our second example shows that the number $f(n, n) = n + 1$ is best.

EXAMPLE 2. Let S be a compact set in R^n , with $\text{inc } S$ a union of $n + 1$ disjoint convex sets $K_1, \dots, K_{n+1}$, each of dimension $n - 2$, and satisfying the following properties:

(1) For $x, y \in K_i$, $C_x = C_y$.

(2) Every n members of $\{C_{q_i}: q_i \in K_i, 1 \leq i \leq n + 1\}$ meet in an n -dimensional set in S , yet $\bigcap \{C_{q_i}: q_i \in K_i, 1 \leq i \leq n\} = \emptyset$.

This may be done so that every n members of $\{A_q: q \in \text{bdry } S\}$ meet in an n -dimensional set as well. Since $\bigcap \{A_q: q \in \text{bdry } S\} = \emptyset$, the number $f(n, 1) = n + 1$ is best possible.

We close with a stronger version of Theorem 1 which holds in the plane.

THEOREM 2. Let S be a compact, connected, nonconvex set in R^2 . Then for $k = 1$ or $k = 2$, $\dim \ker S \geq k$ if and only if every $g(k) = \max\{3, 6 - 2k\}$ inc points of S are clearly visible from a common k -dimensional subset of S . The result is best possible.

PROOF. By [1, Lemma 4], if S is a closed connected set in R^2 , then $\ker S = \bigcap \{C_q \cap \text{conv } S: q \in \text{inc } S\}$. Using this result, it is not hard to show that Lemmas 2 and 3 above hold when S is a compact, connected, nonconvex set in R^2 and when $\text{bdry } S$ is replaced by $\text{inc } S$. An easy adaptation of the proof of Theorem 1 completes the argument.

Examples 1 and 2 of [1] show that the result is best possible.

REFERENCES

1. Marilyn Breen, *A Krasnosel'skii-type theorem for points of local nonconvexity*, Proc. Amer. Math. Soc. **85** (1982), 261–266.
2. ———, *K-dimensional intersections of convex sets and convex kernels*, Discrete Math. **36** (1981), 233–237.
3. ———, *The dimension of the kernel of a planar set*, Pacific J. Math. **82** (1979), 15–21.
4. K. J. Falconer, *The dimension of the kernel of a compact starshaped set*, Bull. London Math. Soc. **9** (1977), 313–316.
5. Meir Katchalski, *The dimension of intersections of convex sets*, Israel J. Math. **10** (1971), 465–470.
6. M. A. Krasnosel'skii, *Sur un critère pour qu'un domaine soit étoilé*, Mat. Sb. (61) **19** (1946), 309–310.
7. S. Nadler, *Hyperspaces of sets*, Dekker, New York, 1978.
8. Nick M. Stavrakas, *The dimension of the convex kernel and points of local nonconvexity*, Proc. Amer. Math. Soc. **34** (1972), 222–224.
9. F. A. Valentine, *Convex sets*, McGraw-Hill, New York, 1964.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OKLAHOMA
73019