MIDDLE NUCLEUS = CENTER IN SEMIPRIME JORDAN ALGEBRAS

KEVIN MCCRIMMON¹ AND NG SEONG NAM²

ABSTRACT. A. A. Albert showed that the middle nucleus and center coincide for a simple Jordan algebra finite-dimensional over a field of characteristic $\neq 2$. E. Kleinfeld extended this to arbitrary simple Jordan algebras of characteristic $\neq 2$. Recently this result has played a crucial role in the structure theory of E. Zelmanov. In this note we extend the result to linear Jordan algebras with no derivation-invariant trivial ideals.

The *left*, *middle*, and *right nucleus* $N_l(A)$, $N_m(A)$, $N_r(A)$ of any nonassociative algebra A consists of the elements $n \in A$ with $[n, A, A] = 0$, $[A, n, A] = 0$, $[A, A, n] = 0$ respectively, where the *associator* is given by $[x, y, z] = (xy)z - x(yz)$. The *nucleus* $N(A)$ consists of the elements in all three nuclei simultaneously, and the *center* $C(A)$ consists of the nuclear elements which commute with A , $[n, A] = 0$. A *derivation* of A is a linear transformation D satisfying

$$(1) \quad D(xy) = D(x)y + xD(y)$$

and hence necessarily

$$(2) \quad D([x, y, z]) = [D(x), y, z] + [x, D(y), z] + [x, y, D(z)]$$

as well. From this we see each nucleus is derivation-invariant, $D(N_i(A)) \subset N_i(A)$ for all D (the same is true of the center). When A is commutative we have

$$(3) \quad [x, y, z] = -[z, y, x],$$

$$(4) \quad [x, y, z] + [y, z, x] + [z, x, y] = 0,$$

so $C(A) = N(A) = N_l(A) = N_r(A) \subset N_m(A)$. Our goal is to show conversely that for semiprime Jordan algebras the reverse inclusion holds as well. This has become important in Zelmanov's work [4], where expressions $[x_3, [x_1, x_2]^2, x_4]$ play a crucial role and one wants to know that if these vanish for all x_3, x_4 (i.e. $[x_1, x_2]^2$ is middle-nuclear) then $[x_1, x_2]^2$ is actually central.

We restrain our quadratic sympathies and work entirely with (nonunital) *linear Jordan algebras* J over a ring of scalars Φ containing $\frac{1}{2}$. Thus J has product xy satisfying the Jordan axioms

$$(5) \quad xy = yx, \text{ i.e. } [x, y] = 0,$$

$$(6) \quad (x^2y)x = x^2(yx), \text{ i.e. } [x^2, y, x] = 0.$$

In addition to the left multiplication operator $L_x(z) = xz$, the U -operator

$$(7) \quad U_x(z) = (2L_x^2 - L_{x^2})(z)$$

plays an important role even in the linear theory. It satisfies the identity

Received by the editors August 28, 1981.

1980 *Mathematics Subject Classification*. Primary 17C10.

¹Research partially supported by NSF Grant MCS-80-02319

²The second author wishes to thank the Department of Pure Mathematics at the University of Sydney for its hospitality, and to thank Dr. K. G. Choo for his assistance and encouragement.

© 1982 American Mathematical Society
 0002-9939/81/0000-0177/\$01.75

$$(8) L_y U_x + U_x L_y = U_{yx,x} (U_{x,y} = U_{x+y} - U_x - U_y).$$

The inner derivations $D_{x,y}$ are defined by

$$(9) D_{x,y}(z) = [L_x, L_y](z) = -[x, z, y];$$

any derivation D satisfies

$$(10) D(U_x y) = U_{D(x),x} y + U_x D(y), [D, D_{x,y}] = D_{D(x),y} + D_{x,D(y)}.$$

A Jordan algebra is *semiprime* if it contains no *trivial* ideals B ($U_B B = 0$) and is *D-semiprime* if it contains no nonzero trivial ideals B invariant under all derivations.

THEOREM. *The middle nucleus and center coincide, $N_m(J) = C(J)$, for any D-semiprime linear Jordan algebra J .*

PROOF. We must show $n \in N_m(J) \Rightarrow n \in N_l(J)$, i.e. $[J, n, J] = 0 \Rightarrow [n, J, J] = 0$. Following Albert [1] we introduce an ideal

$$(11) B = [n, J, J]$$

which measures the failure of n to lie in $N_l(J)$. Here B is spanned by all associators $[n, x, y]$ for $x, y \in J$, where

$$(12) [n, x, y] = [n, y, x]$$

since $0 = [n, x, y] + [x, y, n] + [y, n, x] = [n, x, y] - [n, y, x] + 0$ by (4), (3), and the definition of $n \in N_m(J)$. Thus $[n, x, y] = \frac{1}{2}([n, x, y] + [n, y, x])$ results from linearizing $[n, x, x]$, so

$$(13) B \text{ is spanned by all } [n, x, x] \text{ for } x \in J.$$

To see B is actually an ideal, note JB is spanned by all $y[n, x, x] = -x[n, y, x] + [n, xy, x]$ (by (1), (9)) $= -x[n, x, y] + [n, xy, x]$ (by (12)) $= -\frac{1}{2}[n, x^2, y] + [n, xy, x]$ (by (1)) $\in [n, J, J] = B$.

Now by (13), $[n, B, J]$ is spanned by elements $[n, [n, x, x], y] = -[[n, x, x], y, n] - [y, n, [n, x, x]]$ (by (4)) $= D_{n, D_{n,x}(x)}(y) + 0$ (by (9)) $= \{[D_{n,x}, D_{n,x}] - D_{D_{n,x}(n),x}\}(y)$ (by (10)) $= 0$ (since $D_{J,J}(n) = 0$), so

$$(14) [n, B, J] = 0.$$

Then $0 = [n, B, J] \supset [n, U_B J, J] = U_{[n,B,J]} B J + U_B [n, J, J]$ (by (9), (10)) $= 0 + U_B B$ implies $U_B B = 0$. If J is semiprime then $B = [n, J, J] = 0$ by (11) and $n \in N_l(J)$.

If J is merely *D-semiprime* we must work a little harder. We claim $C = [N_m(J), J, J] = \sum [n, J, J]$ is a trivial *D*-invariant ideal. C is an ideal since each $[n, J, J]$ is by (11), and it is *D*-invariant by (2) and the fact that $N_m(J)$ is *D*-invariant. As in (14) we have

$$(14') [n, C, J] = 0$$

for any $n \in N_m(J)$ since by (12) this is spanned by all $[n, [n', x, x], y] = [[n, n', y], x, x] + [n', [n, x, y], x] + [n', x, [n, x, y]]$ (by (9), (2)) $= 0 + 2[n', [n, x, y], x]$ (by (12) and $n' \in N_m(J)$) $= 2\{n'[n, y, x]\}x - 2n'\{[n, x, y]x\}$ (by (12)) $= [n, x^2, n'y] - [n, n'y, x^2]$ (by (1), (9)) $= 0$ (by (12)). Then $0 = [n, C, J] \supset [n, C J, J] = [n, C, J] J + C[n, J, J]$ (by (1)) $= 0 + C[n, J, J]$ (by (14')); this holds for all $n \in N_m(J)$, so $0 = C C$ and C is trivial as well as *D*-invariant. If J is *D-semiprime* this forces $C = 0$, so $[N_m(J), J, J] = 0$ and $N_m(J) \subset N_l(J)$. $\square$

It is easy to give examples to show that the semiprimeness hypothesis is needed. If we take $J = F/K$ for F the free Jordan algebra on x, y , K the ideal generated by all $[a, y, b]$ for $a, b \in F$, then $n = y + K$ is middle-nuclear in J by construction but is not central: $[n, J, J] \neq 0$ since $[y, x, x] \notin K$ (K is graded, vanishing in total degrees < 3 , and in degree x^2y^1 it is spanned by $[x, y, x] = 0$, so $[y, x, x] \notin K$).

We can also easily give special examples $J = A^+$ for A an associative algebra. Here $n \in N_m(J) \Leftrightarrow [[J, J], n] = 0$ and $n \in N_l(J) \Leftrightarrow [[n, J], J] = 0$. If $A = \Phi E_{11} + \Phi E_{12} + \Phi E_{22}$ consists of all upper triangular 2×2 matrices over Φ , then $[J, J] = \Phi E_{12}$, so $n = E_{12}$ lies in $N_m(J)$, yet $n \notin N_l(J)$ since $[[n, E_{11}], E_{11}] = -[E_{12}, E_{11}] = E_{12} \neq 0$. Note that n falls in the trivial ideal $B = \Phi E_{12}$.

Another example is the algebra $J(Q, c)$ determined by a quadratic form Q ; here $J = \Phi c \oplus V$ where $[v_1, v_2, v_3] = Q(v_1, v_2)v_3 - Q(v_2, v_3)v_1$, so $B = \text{Rad } Q \subset N_m$ but $B \not\subset N_r$ if $Q \neq 0$.

Without reference to middle nuclei we can establish

THEOREM. *If J is a semiprime linear Jordan algebra then $C(I) \subset C(J)$ for any ideal $I \triangleleft J$. More generally, $C(J) \subset C(\tilde{J})$ as soon as $J \triangleleft \tilde{J}$ and J contains no trivial ideals of $\tilde{J}$ invariant under all derivations of $\tilde{J}$ which map J into itself.*

PROOF. The first assertion follows from the second, since by a result of Slin'ko [4] if J is semiprime so is any $I \triangleleft J$. Assume J is D -semiprime in $\tilde{J}$ in the above sense; for convenience we may assume that $\tilde{J}$ is unital. We must show that if $c \in C(J)$ then $c \in C(\tilde{J}) = N_l(\tilde{J})$, i.e. $[c, \tilde{J}, \tilde{J}] = 0$.

All the derived ideals $J^{(n)}$ (where $J^{(0)} = J, J^{(n+1)} = U_{J^{(n)}}J^{(n)}$) are invariant under the indicated derivations and remain ideals in $\tilde{J}$, as are their annihilators $(J^{(n)})^\perp$ (if $B \triangleleft J$ is invariant so is $B^\perp = \{z \in \tilde{J} | \{z, B, \tilde{J}\} = 0\}$, and $B^\perp \triangleleft \tilde{J}$ since $\{\tilde{J}z, B, \tilde{J}\} = \tilde{J}\{z, B, \tilde{J}\} - \{z, B, \tilde{J}\tilde{J}\} + \{z, \tilde{J}B, \tilde{J}\} \subset \tilde{J}\{z, B, \tilde{J}\} = 0$). Moreover, $B = J \cap J^{(n)\perp}$ are solvable: $B^{(n)} \subset J^{(n)} \cap J^{(n)\perp}, B^{(n+1)} = 0$. If J contains no trivial invariant ideals then it contains no solvable invariant ideals of $\tilde{J}$, so $B = 0$:

$$(15) \quad \text{if } J \text{ is } D\text{-semiprime in } \tilde{J} \text{ then } J \cap J^{(n)\perp} = 0.$$

If $D(J^{(n)}) = 0$ then $D(\tilde{J}) \subset J^{(n+1)\perp}$ since $\{D(\tilde{J}), J^{(n+1)}, \tilde{J}\} \subset \{D(\tilde{J}), J^{(n)}, J^{(n)}\}$ (because $\{d, U_x y, \tilde{a}\} = \{d, x, \{y, x, \tilde{a}\}\} - \{d, U_x \tilde{a}, y\}$ where $x, y \in J^{(n)} \triangleleft \tilde{J} = D(\{\tilde{J}, J^{(n)}, J^{(n)}\}) - \{\tilde{J}, D(J^{(n)}), J^{(n)}\} - \{\tilde{J}, J^{(n)}, D(J^{(n)})\} = 0$ (because $J^{(n)} \triangleleft \tilde{J}, D(J^{(n)}) = 0$), so from (15) we see

$$(16) \quad \begin{aligned} &\text{if } J \text{ is } D\text{-semiprime in } \tilde{J} \text{ then } D(J^{(n)}) = 0 \Rightarrow D(\tilde{J}) = 0 \\ &\text{for any derivation of } \tilde{J} \text{ into } J. \end{aligned}$$

In particular, for $D = D_{c, J}$ as in (9) we see $[c, J, J] = 0$ (by $c \in N_l(J)$) implies $[c, \tilde{J}, J] = 0$, hence for $D = D_{c, \tilde{J}}$ we see $[c, J^2, \tilde{J}] = -[J^2, \tilde{J}, c] - [\tilde{J}, c, J^2]$ (by (4)) $\subset [c, \tilde{J}, J^2] + [J, c, J\tilde{J} + \tilde{J}J]$ (by (3) and linearized (6)) $\subset [c, \tilde{J}, J] + [J, c, J]$ (by $J \triangleleft \tilde{J} = 0$ (by the above and $c \in N_m(J)$) implies $[c, \tilde{J}, \tilde{J}] = 0$ as desired. $\square$

REFERENCES

1. A. A. Albert, *On the nuclei of a simple Jordan algebra*, Proc. Nat. Acad. Sci. U.S.A. **50** (1963), 446–447.
2. N. Jacobson, *Structure theory of Jordan algebras*, University of Arkansas Lecture Notes, Fayetteville, 1981.
3. E. Kleinfeld, *Middle nucleus=center in a simple Jordan ring*, J. Algebra **1** (1964), 40–42.
4. E. I. Zelmanov, *Jordan division rings*, Algebra i Logika **18** (1979), 286–310.
5. Ng Seong Nam, *Middle nucleus of semiprime Jordan rings*, Nanta Mathematica **9** (1976), 1–3.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE,
VIRGINIA 22903

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITI SAINS MALAYSIA, PENANG,
MALAYSIA