UNIQUENESS OF SOLUTIONS OF PARTIAL DIFFERENTIAL EQUATIONS WHEN THE INITIAL SURFACE IS CHARACTERISTIC AT A POINT

LETITIA J. KORBLY¹

ABSTRACT. Uniqueness in the Cauchy problem for hyperbolic operators degenerate at a point on the initial surface depends on values of the coefficients of the lower order terms. If the operator P is doubly characteristic at the origin with respect to the $t = 0$ line, P has uniqueness for functions which are smooth enough if the coefficient of the D_t term does not lie in a certain discrete set of numbers.

Introduction. We consider operators of the form

$$(1) \quad P = A(x, t)D_t^2 + 2B(x, t)D_x D_t - C(x, t)D_x^2 + \lambda(x, t)D_t + R_1$$

where all the coefficients are real C^∞ functions, A is nonnegative, C is positive, R_1 is a first order linear operator in D_x ,

$$D_x = \partial/\partial x \quad \text{and} \quad D_t = \partial/\partial t.$$

When P is doubly characteristic at the origin with respect to the surface given by $t = 0$, uniqueness of the solution to the problem:

$$\begin{aligned} Pu &= F, & (x, t) \in \Omega, \text{ a neighborhood of the origin, } t \geq 0, \\ u &= u_0(x), & \text{if } t = 0, \\ D_t u &= u_1(x), & \text{if } t = 0, \end{aligned}$$

is shown to depend on the values of the lower order terms, principally λ . If λ lies outside a certain discrete set of values the solutions to the problem will be unique whenever μ is smooth enough.

If λ lies in this discrete set of values, uniqueness will depend on R_1 . The dependence on the lower order terms is very complex. There is always uniqueness when

$$(2) \quad \lambda_0 = \lambda - D_x B - D_t A/2 \geq 0.$$

As λ_0 becomes more negative, u must be assumed to be smoother to get the same result.

There are some values of the coefficients for which no uniqueness results can be obtained. If $c > 0$,

$$(3) \quad Pu = x^2 D_t^2 u - c^2 D_x^2 u - (2n + 1)c D_t u$$

Received by the editors December 10, 1981.

1980 *Mathematics Subject Classification*. Primary 35L80, 35A07, 35L70; Secondary 35L15.

Key words and phrases. Hyperbolic PDE, Cauchy problem, doubly characteristic.

¹Part of this work is based on the author's thesis written under F. Trèves at Rutgers University

© 1982 American Mathematical Society
0002-9939/82/0000-0209/\$02.50

and

$$\tilde{X}u = xD_t u - cD_x u,$$

then

$$P\tilde{X}^N h(ct - x^2/2) = 0, \quad h \in C_c^\infty(R).$$

The assertion is proved by taking the support of h to be the positive halfline. Finally, if P is as in (3), $P + \gamma$ is shown to have uniqueness if γ is a nonzero function.

The operators. P is characteristic with respect to the initial surface at the origin when A vanishes at the origin, and doubly characteristic with respect to this surface when both B and $\text{grad } A$ also vanish there.

In order to use the concatenation technique of Višik and Grušin, [12], it is necessary to factor the principal part of the operator in a neighborhood of the origin which we will take to be Ω . The following hypotheses are assumed:

(Ha) There is a C^∞ function, a , such that

$$A = a^2 \quad \text{and} \quad D_x a = a_x > 0.$$

(Hb) There is a C^∞ function, b , such that

$$B = ba.$$

Then

$$P = a^2 D_t^2 + 2ab D_t D_x + b^2 D_x^2 - (C + b^2) D_x^2 + \lambda D_t + R_1.$$

Let

$$c^2 = C + b^2, \quad \text{where } c > 0.$$

Define

$$\tilde{Y} = aD_t + bD_x + cD_x \quad \text{and} \quad \tilde{X} = aD_t + bD_x - cD_x.$$

So

$$(4) \quad P = \tilde{Y}\tilde{X} + \lambda' D_t + R_2,$$

where R_2 is a first order operator in D_x and

$$\lambda' = \lambda - (b + c)a_x.$$

This form is used in the concatenations. The commutator, $[\tilde{Y}, \tilde{X}]$, of $\tilde{Y}$ and $\tilde{X}$ acting on a function is

$$(5) \quad [\tilde{Y}, \tilde{X}]u = (\tilde{Y}\tilde{X} - \tilde{X}\tilde{Y})u = (2a_x c D_t + R_3)u,$$

where R_3 is a first order operator in D_x . The hypotheses (1) and (Ha) guarantee $a_x c > 0$.

DEFINITION 1. The following norms are used:

$$\|u\|^2 = \|u\|_0^2 = \int \int_{\Omega} |u|^2 dx dt, \quad u \in \mathcal{H}^2(\Omega).$$

Then

$$\|u\|_{N+1}^2 = \|D_t u\|_N^2 + \sigma^2 \|u\|_N^2 + \|D_x u\|_N^2.$$

The following norm is taken from [1]:

$$\|u\|_{1,1}^2 = \|aD_t u\|_0^2 + \sigma^2 \|au\|_0^2 + \|D_x u\|_0^2.$$

Note that for all u in $C_c^\infty(\Omega)$, and $\sigma > \sigma_0$:

$$\|u\|_{1,1} \leq \|u\|_1.$$

Also, if $u \in C_c^\infty(\Omega)$,

$$\|(D_t + \sigma)u\|^2 = \|D_t u\|^2 + \sigma^2 \|u\|^2.$$

Similar statements are true when other norms are used.

DEFINITION 2. An operator, P , has $(UCP)_N$ in an open set, Ω , if, whenever u is an element of $H^N(\Omega)$, and $\text{supp } u \cap \{t \leq 0\}$ is compact, then $Pu = 0$ in Ω , implies $u = 0$ in a neighborhood of $\text{supp } u \cap \{t \leq 0\}$.

Let $\Omega^- = \Omega \cap \{t \leq 0\}$. The following lemma is only stated because its proof is an easy modification of that in [8].

LEMMA 1. If there are constants c and σ_0 such that for every function v in $C_c^\infty(\Omega)$, and every $\sigma > \sigma_0$, the Carleman estimate

$$(6) \quad C \|e^{-\sigma t} P v\|_N \geq \|e^{-\sigma t} v\|_0$$

holds, then P has $(UCP)_{N+m}$. Here m is the order of P .

Define

$$P_\phi u = \exp(-\sigma t) P(\exp(\sigma t) u).$$

The substitution

$$v = \exp(\sigma t) u$$

shows that (6) is equivalent to the following: there is a constant c , such that

$$c \|P_\phi u\|_N \geq \|u\|_0, \quad u \in C_c^\infty(\Omega), \quad \sigma > \sigma_0.$$

This is the estimate used in proving the theorem.

Proof of the theorem when λ_0 is positive. Let A_t be the partial of A with respect to t , B_x , the partial of B with respect to x , and C_t and C_x be the partials of C with respect to t and x respectively.

LEMMA 2. If P is as in (1) where A is nonnegative and C is positive, and

$$0 < L < \lambda_0 = \lambda(0, 0) - B_x(0, 0) - A_t(0, 0)/2,$$

then there are $c > 0$ and σ_0 , depending only on L and the coefficients of P such that if Ω is a sufficiently small neighborhood of the origin,

$$\langle P_\phi u, (D_t + \sigma)u \rangle \geq c \sigma \|u\|_{1,1}^2 + L \|u\|_1^2, \quad u \in C_c^\infty(\Omega), \quad \sigma > \sigma_0.$$

PROOF. Integration by parts in D_t shows

$$(7) \quad \langle A(D_t + \sigma)u, (D_t + \sigma)u \rangle = \sigma \|A^{1/2}(D_t + \sigma)u\|^2 - \langle A_t/2(D_t + \sigma)u, (D_t + \sigma)u \rangle$$

whenever u is in $C_c^\infty(\Omega)$.

Integration by parts in D_x shows that if $u \in C_c^\infty(\Omega)$,

$$(8) \quad \langle 2BD_x(D_t + \sigma)u, (D_t + \sigma)u \rangle = -\langle B_x(D_t + \sigma)u, (D_t + \sigma)u \rangle.$$

An integration by parts in D_t followed by another in D_x shows if $u \in C_c^\infty(\Omega)$,

$$(9) \quad \begin{aligned} & -\langle CD_x^2 u, (D_t + \sigma)u \rangle \\ & = \sigma \|C^{1/2} D_x u\|^2 - \langle (C_t/2) D_x u, D_x u \rangle + \langle (C_x/2) D_x u, (D_t + \sigma)u \rangle. \end{aligned}$$

We use (1), (7), (8), and the last equation, (9), to get if $u \in C_c^\infty(\Omega)$,

$$(10) \quad \langle P_\phi u, (D_t + \sigma)u \rangle = \sigma \|a(D_t + \sigma)u\|^2 + \|(\sigma C - C_x/2)^{1/2} D_x u\|^2 \\ + \|(\lambda_0)^{1/2} (D_t + \sigma)u\|^2 + \langle R_4 u, (D_t + \sigma)u \rangle$$

where R_4 is a first order operator in D_x .

Schwarz's inequality implies that for some R and any positive ϵ ,

$$\langle R_4 u, (D_t + \sigma)u \rangle \leq \epsilon \|(D_t + \sigma)u\|^2 + 1/(2\epsilon) \|R_4 u\|^2, \quad u \in C_c^\infty(\Omega).$$

Fix ϵ so that $L = \lambda_0 - \epsilon > 0$ after shrinking Ω if necessary. Then choose σ_0 so that $\sigma_0 C - C_x/2 - R/2\epsilon$ is positive. This implies that the first three terms on the right in (10) dominate the last term on the right, which proves the lemma. $\square$

COROLLARY. *If P is as in (1) and (Ha) and (Hb) hold, the conclusion of Lemma 2 is true if*

$$\lambda - a_x b > L > 0.$$

PROOF. Both ab_x and $A_t/2 = aa_t$ vanish at the origin.

Proof of the theorem when λ_0 is negative. It is easy to show that $P_\phi u$ dominates $Y^N u$ if λ_0 is not less than $-2Na_x c$, where

$$Yu = e^{-\sigma t} \tilde{Y}(e^{\sigma t} u), \quad u \in C_c^\infty(\Omega),$$

and

$$Y^{N+1}u = YY^N u,$$

But no inequality which is uniform in σ bounds Yu away from 0 in a neighborhood of the origin. To prove the theorem when $\lambda_0 < 0$, more structure is necessary. The next lemma handles the commutators between Y^N and P_ϕ . Let

$$Xu = e^{-\sigma t} X(e^{\sigma t} u), \quad u \in C_c^\infty(\Omega).$$

LEMMA 3. *If P is as in (1) and (Ha) and (Hb) are satisfied, then*

$$(11) \quad Y^N P_\phi u = (P_\phi + N[Y, X])Y^N u + \sum_{k < N} L_k Y^k u, \quad u \in C_c^\infty(\Omega).$$

Here L_k is a first order operator in D_x and D_t .

PROOF. The proof is by induction. The lemma is clearly true when $N = 0$. The induction hypothesis is (11). Then if $u \in C_c^\infty(\Omega)$,

$$Y^{N+1} P_\phi u = Y \left[(XY + \lambda'(D_t + \sigma) + R_2 + N[Y, X])Y^N u + \sum_{k < N} L_k Y^k u \right].$$

Y is commuted with the rest of the right-hand side above. The commutator satisfies

$$[Y, XY] = [Y, X]Y.$$

All the other commutators are first order. This recovers the inductive hypothesis. The lemma follows. $\square$

COROLLARY. *If P is as in (1) and satisfies (Ha) and (Hb), and if*

$$\lambda(0, 0) - a_x(0, 0)b(0, 0) = \lambda_0(0, 0) > -2Na_x(0, 0)c(0, 0)$$

and

$$P_\phi^N = P_\phi + N[Y, X]$$

then there is some neighborhood Ω of the origin where P^N satisfies the hypotheses of Lemma 2.

PROOF. Define

$$(12) \quad \lambda_N = \lambda_0 + 2Na_x c$$

then the hypothesis of the corollary implies that in some neighborhood of the origin λ_N is strictly positive, and must be bounded away from 0 in some neighborhood of the origin which is taken to be Ω . But

$$P_\phi^N = XY + \lambda_N D_t + R_N$$

where R_N is a first order operator in D_x . This proves the corollary. $\square$

Now the main theorem will be proved.

THEOREM 1. *If P is as in (1) and (Ha) and (Hb) are satisfied then P has $(UCP)_{N+2}$ if*

$$(h1) \quad \lambda' + 2ka_x(0,0)c(0,0) \neq 0 \text{ for any } k = 1, 2, 3, \dots$$

and

$$(h2) \quad \lambda_0 + 2Na_x(0,0)c(0,0) > 0.$$

PROOF. Lemma 3 implies that there are constants c , r , and σ_0 such that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$(13) \quad c\|P_\phi u\|_N \geq \|Y^N P_\phi u\|_0 \geq \|P_\phi^N Y^N u\|_0 - S_0.$$

Here

$$S_J = \sigma^J \sum_{k < N-J} r_k \|Y^k u\|_1.$$

The corollary to Lemma 3 and (h2) establish that P^N satisfies the hypothesis of Lemma 2, so there are constants c and σ_0 , such that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$(14) \quad c\langle P_\phi^N Y^N u, (D_t + \sigma)Y^N u \rangle \geq \sigma \|Y^N u\|_{1,1}^2 + \|Y^N u\|_1^2.$$

Since

$$\|(D_t + \sigma)v\|_0 \leq \|v\|_1$$

and because the 1,1 norm is smaller than the 1 norm, (14) implies there are constants, c and σ_0 , such that if $u \in C_c^\infty(\Omega)$, $\sigma > \sigma_0$ and $N \geq 1$,

$$c\|P_\phi u\|_N \geq \sigma \|Y^N u\|_{1,1} + \|Y^N u\|_1.$$

Substitution of the last inequality into (13) implies there is a constant c such that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$c\|P_\phi u\|_N \geq \sigma \|Y^N\|_{1,1} + \|Y^N u\|_1 - S_0.$$

This inequality is the first step in an inductive argument. The inductive hypothesis is that there are constants c , r , and σ_0 , depending on J , N , and P , but not on $u \in C_c^\infty(\Omega)$ or $\sigma > \sigma_0$, such that

$$c\|P_\phi u\|_N \geq \sigma^J (\sigma \|Y^{N-J} u\|_{1,1} + \|Y^{N-J} u\|_1) - S_J.$$

We now try to recover the inductive hypothesis for $J+1$. The inductive hypothesis for J implies there are constants c' , r' , and σ_0 such that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$(16) \quad c'\|P_\phi u\|_N \geq \sigma^{J+1} \|XY Y^{N-J-1} u\|_0 - S_J.$$

But the definition of the norms implies there is a constant c'' such that

$$c'' \|P_\phi u\|_N \geq \sigma^{J+1} \|Y^{N-J-1} P_\phi u\|_0, \quad u \in C_c^\infty(\Omega), \sigma > \sigma_0.$$

Lemma 3 applied to the right-hand side of the inequality above implies the existence of constants c''' , r'' , and σ_0 such that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$(17) \quad c''' \|P_\phi u\|_N \geq \sigma^{J+1} \|P_\phi^{N-J-1} Y^{N-J-1} u\|_0 - S_{J+1}.$$

Subtracting (16) from the last inequality gives the following for $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$: if $\underline{c} = c' + c'''$, and λ_{N-J-1} is as in (12),

$$\underline{c} \|P_\phi u\|_N \geq \sigma^{J+1} \|\lambda_{N-J-1} (D_t + \sigma) Y^{N-J-1} u\|_0 - \sigma^J r' \|Y^{N-J-1} u\|_1 - S_{J+1}.$$

Here r'' and σ_0 have been increased to absorb all the terms of S_J but the $(N-J-1)$ st. The $(N-J-1)$ st term is the one which must be dominated to recover the inductive hypothesis.

Since (h2) guarantees that $\lambda_{N-J-1} = 0$ in Ω , we can add a multiple, m , of the last inequality to (17) to get for each $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$\underline{c}' \|P_\phi u\|_N \geq \sigma^{J+1} \|P_\phi^g Y^{N-J-1} u\|_0 - \sigma^J m r' \|Y^{N-J-1} u\|_1 - S_{N+1},$$

where $\underline{c}' = m \underline{c} + c'''$. The multiple m is chosen so that P_ϕ^g satisfies the hypothesis of Lemma 2. This now implies that for some new $\underline{c}''$ and σ_0 , if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$\begin{aligned} \underline{c}'' \|P_\phi u\|_N &\geq \sigma^{J+1} (\sigma \|Y^{N-J-1} u\|_{1,1} + \|Y^{N-J-1} u\|_1) \\ &\quad - \sigma^J r \|Y^{N-J-1} u\|_1 - S_{N+1}. \end{aligned}$$

If σ_0 is taken bigger than r , then the inductive hypothesis is recovered. This proves the estimate. The theorem follows from Lemma 1. $\square$

With the introduction of more machinery, as in [1 and 2], a slightly stronger theorem can be proved. It is possible to show that the conditions of Theorem 1 imply P has $(UCP)_{N+1}$. It is probably not possible to conclude that P has uniqueness for functions with continuous second partials when λ_0 is negative. It is also possible to strengthen the conclusion of Theorem 2 slightly, but the gain is small compared to the increase in difficulty of the proof.

COROLLARY. *If P is as in (1) and (Ha) and (Hb) are satisfied, then P has $(UCP)_{N+2}$ if*

$$(h1)' \quad \lambda \neq (b - (2k-1)c)a_x, \quad k = 1, 2, 3, \dots,$$

and

$$(h2)' \quad \lambda > (b - 2Nc)a_x.$$

PROOF. The proof that $(h2)'$ is equivalent to (h2) is exactly as in the corollary to Lemma 2. The equivalence of $(h1)'$ to (h1) follows directly from (2), the definition of λ_0 .

COROLLARY 2. *If P is as in (1) and (Ha) and (Hb) hold, then P has $(UCP)_{N+2}$ if for every positive integer k , when $t = 0$,*

$$(h1)'' \quad \lambda(0,0) \neq \lim_{x \rightarrow 0} \{A_x [B - (2k-1)(AC + B^2)^{1/2}] / (2A)\}$$

and when $t = 0$,

$$(h2)'' \quad \lambda(0,0) > \lim_{x \rightarrow 0} \{A_x [B - 2N(AC + B^2)^{1/2}] / (2A)\}.$$

PROOF. This follows directly from the definitions of a , b , and c . We have

$$a_x = A_x/(2A^{1/2}) \quad \text{and} \quad ba_x = A_x B/(2A).$$

Also

$$ca_x = A_x(AC + B^2)^{1/2}/(2A). \quad \square$$

The final theorem shows the complex dependence of uniqueness on the terms R_N when λ violates (h1).

THEOREM 2. Let γ be a nonvanishing function,

$$Pu = (XY + N[Y, X] + \gamma)u,$$

$$Y\gamma = 0 \quad \text{and} \quad [Y, [Y, X]] = 0.$$

Then P has $(UCP)_{N+4}$.

PROOF. Application of Lemma 2 shows there are constants c and σ_0 , depending on P but not on $u \in C_c^\infty(\Omega)$ or $\sigma > \sigma_0$, such that

$$c\|P_\phi u\|_{N+2} \geq \|Y^{N+1}P_\phi u\|_1 = \|P_\phi^{N+1}Y^{N+1}u\|_1.$$

Here P^{N+1} satisfies the hypothesis of Lemma 2. If for $u \in C_c^\infty(\Omega)$,

$$\|u\|_{2,1}^2 = \|u_t\|_{1,1}^2 + \|u_x\|_{1,1}^2 + \sigma^2\|u\|_{1,1}^2,$$

an easy extension of Lemma 2 shows there are constants c and σ_0 such that

$$c\|P_\phi u\|_{N+2} \geq \sigma\|Y^{N+1}u\|_{2,1} + \|Y^{N+1}u\|_2, \quad u \in C_c^\infty(\Omega), \quad \sigma > \sigma_0.$$

This implies there are c' and σ_0 such that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$c'\|P_\phi u\|_{N+2} \geq \sigma\|XY^{N+1}u\|_1.$$

But the definitions of the norms imply that for some c and σ_0 , if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$

$$(18) \quad c''\|P_\phi u\|_{N+2} \geq \sigma\|Y^N P_\phi u\|_1 = \sigma\|P_\phi^N Y^N u\|_1.$$

Here a whole derivative has been lost. Subtracting (11) from (18) implies that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$(c'' + c')\|P_\phi u\|_{N+2} \geq \sigma|\gamma(0,0)/2|\|Y^N u\|_1.$$

The induction starts here. The inductive hypothesis is: there are c and σ_0 such that if $u \in C_c^\infty(\Omega)$ and $\sigma > \sigma_0$,

$$c\|P_\phi u\|_{N+2} \geq \sigma^{J+1}\|Y^{N-J}u\|_1.$$

The induction is similar to that in Theorem 1. The hypotheses of this theorem guarantee all the commutators except $[Y, X]$ vanish. When $J = N$, an estimate which proves the theorem is obtained. $\square$

REFERENCES

1. L. Boutet de Monvel and F. Treves, *On a class of pseudodifferential operators with double characteristics*, Invent. Math. **24** (1974), 1–34.
2. ———, *On a class of systems of pseudodifferential operators with double characteristics*, Comm. Pure Appl. Math. **27** (1974), 59–89.
3. M. S. Baouendi and C. Goulaouic, *Cauchy problems with characteristic initial surface*, Comm. Pure Appl. Math. **26** (1973), 555–575.
4. A. Gilioli and F. Treves, *An example in the solvability theory of linear PDE's*, Amer. J. Math. **96** (1974), 367–385.
5. L. Hormander, *Linear partial differential operators*, 3rd printing, Springer-Verlag, Berlin and New York, 1969.
6. A. Menikoff, *Uniqueness of the Cauchy problem for a class of partial differential operators with double characteristics*, Indiana Univ. Math. J. **25** (1976), 1–21.
7. F. Treves, *Concatenations of second order evolution equations applied to local solvability and hypoellipticity*, Comm. Pure Appl. Math. **26** (1973), 201–250.
8. ———, *Linear partial differential equations with constant coefficients*, Gordon & Breach, New York, 1956.
9. ———, *A link between local solvability of pseudodifferential operators and uniqueness in the Cauchy problem*, Amer. J. Math. **94** (1972), 167–188.
10. ———, *Discrete phenomena in uniqueness in the Cauchy problem*, Proc. Amer. Math. Soc. **46** (1974), 229–233.
11. M. Protter, *The Cauchy problem for a hyperbolic second order equation*, Canad. J. Math. **6** (1954), 542–553.
12. M. I. Višik and V. V. Grušin, *Elliptic pseudodifferential operators on a closed manifold which degenerate on a submanifold*, Soviet Math. Dokl. **10** (1969), 1316–1320.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALABAMA, BIRMINGHAM, ALABAMA 35294