STABLE HOMOTOPY TYPES OF STUNTED REAL PROJECTIVE SPACES

TEIICHI KOBAYASHI¹

ABSTRACT. The purpose of this note is to improve partly necessary conditions for two stunted real projective spaces to be of the same stable homotopy type given in [4].

1. Introduction. Let RP^n be the n -dimensional real projective space. If $k \geq 0$, we have natural inclusion $RP^{n-1} \subset RP^{n+k}$, and denote by RP_n^{n+k} the quotient space RP^{n+k}/RP^{n-1} . S. Feder, S. Gitler and M. E. Mahowald in [4] have obtained necessary and sufficient conditions for the stunted real projective spaces RP_n^{n+k} and RP_m^{m+k} to be of the same stable homotopy type for a great number of values of n , m and k . Let $\phi(k)$ be the number of integers s such that $0 < s \leq k$ and $s \equiv 0, 1, 2$ or $4 \pmod{8}$, and let $A_k = 2^{\phi(k)}$.

THEOREM 1. *Let k and n be integers such that $n \equiv 0 \pmod{A_k/2}$. If $n \not\equiv m \pmod{A_k}$, then the spaces RP_n^{n+k} and RP_m^{m+k} are not of the same stable homotopy type.*

Theorem 1 partly improves Theorem 1.1 in [4].² For the converse of the result, it is well known (cf. [1–3]) that if $n \equiv m \pmod{A_k}$, then the spaces RP_n^{n+k} and RP_m^{m+k} are of the same stable homotopy type.

2. Proof of Theorem 1. In case $k \leq 8$, we can prove without difficulty that if RP_n^{n+k} and RP_m^{m+k} are of the same stable homotopy type, then $n \equiv m \pmod{A_k}$, using naturality of the squaring operations.

Consider the case $k > 8$. Suppose $m < n$ and there is a homotopy equivalence $f: S^t RP_n^{n+k} \rightarrow S^{t+n-m} RP_m^{m+k}$ for some integer $t \geq 0$. Then $n \equiv m \pmod{A_k/2}$, by Theorem 1.1 in [4] (cf. also [5 and 6]). We may assume $t \equiv 0 \pmod{8}$, and so $t + n - m \equiv 0 \pmod{8}$. Put $t = 8u$ and $t + n - m = 8v$ and let

$$I^u: \widetilde{KO}(RP_n^{n+k}) \rightarrow \widetilde{KO}(S^t RP_n^{n+k}), \quad I^v: \widetilde{KO}(RP_m^{m+k}) \rightarrow \widetilde{KO}(S^{8v} RP_m^{m+k})$$

Received by the editors January 27, 1982.

1980 *Mathematics Subject Classification.* Primary 55P10; Secondary 55N15.

Key words and phrases. Stunted real projective space, stable homotopy type, KO -theory, Adams operation.

¹The author wishes to express his thanks to the referee for his kind suggestion.

²Theorem 1 was known to the authors of [4], but the result was not published.

©1983 American Mathematical Society
0002-9939/82/0000-0442/\$01.50

be the isomorphisms defined by the Bott periodicity. If $l \equiv 0 \pmod{4}$, we have by [1, Theorem 7.4],

$$\widetilde{KO}(RP^{l+k}_+) \cong Z \oplus \widetilde{KO}(RP^{l+k}_{+1}) \cong Z \oplus Z_{A_k}.$$

Denote by $\nu^{(l)}$ and $\lambda^{(l+1)}$ the generators of the first summand and the second summand, respectively. We may put

$$f^*I^v\nu^{(m)} = \varepsilon I^u\nu^{(n)} + aI^u\lambda^{(n+1)}, \quad f^*I^v\lambda^{(m+1)} = (2b + \varepsilon)I^u\lambda^{(n+1)},$$

where a and b are some integers and $\varepsilon = \pm 1$. Let Ψ^3 be the third Adams operation. Then by [1, Theorem 5.1, Corollary 5.3 and Theorem 7.4], we have

$$\begin{aligned} \Psi^3 f^* I^v \nu^{(m)} &= f^* \Psi^3 I^v \nu^{(m)} = 3^{4v} f^* I^v \Psi^3 \nu^{(m)} \\ &= 3^{4v} f^* I^v \{3^{m/2} \nu^{(m)} + 2^{-1}(3^{m/2} - 1)\lambda^{(m+1)}\} \\ &= 3^{4v} \{3^{m/2}(\varepsilon I^u \nu^{(n)} + a I^u \lambda^{(n+1)}) + 2^{-1}(3^{m/2} - 1)(2b + \varepsilon) I^u \lambda^{(n+1)}\} \\ &= \varepsilon 3^{4u+n/2} I^u \nu^{(n)} \\ &\quad + 3^{4u} [3^{n-m/2} a + 3^{(n-m)/2} \{(3^{m/2} - 1)b + 2^{-1}(3^{m/2} - 1)\varepsilon\}] I^u \lambda^{(n+1)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \Psi^3 f^* I^v \nu^{(m)} &= \varepsilon \Psi^3 I^u \nu^{(n)} + a \Psi^3 I^u \lambda^{(n+1)} = \varepsilon 3^{4u} I^u \Psi^3 \nu^{(n)} + a 3^{4u} I^u \Psi^3 \lambda^{(n+1)} \\ &= \varepsilon 3^{4u} I^u \{3^{n/2} \nu^{(n)} + 2^{-1}(3^{n/2} - 1)\lambda^{(n+1)}\} + 3^{4u} a I^u \lambda^{(n+1)} \\ &= \varepsilon 3^{4u+n/2} I^u \nu^{(n)} + 3^{4u} \{2^{-1}(3^{n/2} - 1)\varepsilon + a\} I^u \lambda^{(n+1)}. \end{aligned}$$

We have, therefore,

$$3^{n/2}a + 3^{(n-m)/2}\{(3^{m/2} - 1)b + 2^{-1}(3^{m/2} - 1)\varepsilon\} \equiv 2^{-1}(3^{n/2} - 1)\varepsilon + a \pmod{A_k},$$

because the order of the element $I^u\lambda^{(n+1)}$ is equal to $A_k = 2^{\phi(k)}$. Since $n \equiv m \equiv 0 \pmod{A_k/2}$, $3^{n/2} - 1 \equiv 3^{m/2} - 1 \equiv 0 \pmod{A_k}$ by [1, Lemma 8.1]. Hence we obtain $3^{(n-m)/2} - 1 \equiv 0 \pmod{2A_k}$, and so we get $n - m \equiv 0 \pmod{A_k}$ by [1, Lemma 8.1].

3. Remark. Combining Theorem 1 with known results, we have

COROLLARY 2. Let $k \leq 8$, or $k > 8$ and $n \equiv 0 \pmod{A_k/2}$. Then RP^{n+k}_n and RP^{m+k}_m are of the same stable homotopy type if and only if $n \equiv m \pmod{A_k}$.

REFERENCES

1. J. F. Adams, *Vector fields on spheres*, Ann. of Math. (2) **75** (1962), 603–632.
2. ———, *On the groups $J(X)$* , II, Topology **3** (1965), 137–171.
3. M. F. Atiyah, *Thom complexes*, Proc. London Math. Soc. **11** (1961), 291–310.
4. S. Feder, S. Gitler and M. E. Mahowald, *On the stable homotopy type of stunted projective spaces*, Bol. Soc. Mat. Mexicana **22** (1977), 1–5.
5. T. Kobayashi, *On the odd order nonsingular immersions of real projective spaces*, J. Sci. Hiroshima Univ. Ser. A–I **33** (1969), 197–207.
6. T. Kobayashi and M. Sugawara, *On stable homotopy types of stunted lens spaces*, Hiroshima Math. J. **1** (1971), 287–304.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KOCHI UNIVERSITY, KOCHI 780, JAPAN