F_p CLASSES AND HYPERGEOMETRIC SERIES

DANIEL S. MOAK

ABSTRACT. For $\operatorname{Re}(p) \geq 0$, let $F_p = \{f: f(z) = \int_{|x|=1} (1-xz)^{-p} d\mu(x), |z| < 1, \mu \text{ a probability measure on } |x|=1\}$, and let $F_p \cdot F_q = \{f \cdot g: f \in F_p, g \in F_q\}$. Brickman, Hallenbeck, MacGregor and Wilken proved that $p > 0$ and $q > 0$, then $F_p \cdot F_q \subseteq F_{p+q}$. Kent Pearce recently proved a converse result: if $F_p \cdot F_q \subseteq F_{p+q}$, then $p > 0$ and $q > 0$, or $p = q = 1 + it$ for some real t . The case $p = q = 1 + it, t \neq 0$, will be excluded. Consequently a full converse of the above theorem holds.

Introduction. Let $F_p = \{f: f(z) = \int_{|x|=1} (1-xz)^{-p} d\mu(x), |z| < 1, \mu \text{ a probability measure on } |x|=1\}$, where $\operatorname{Re} p > 0$. Also let $F_p \cdot F_q = \{f \cdot g: f \in F_p, g \in F_q\}$, where $\cdot$ denotes the Hadamard convolution. Pearce [1] proved that if $F_p \cdot F_q \subseteq F_{p+q}$ then $p > 0$ and $q > 0$, or $p = q = 1 + it, t$ real. In this paper, the case $p = q = 1 + it, t \neq 0$, will be excluded, using an identity involving hypergeometric functions.

Pierce [1] used the following theorem in establishing his result.

THEOREM A. If $F_p \cdot F_q \subseteq F_{p+q}$ then

$$(1) \quad \operatorname{Re} \left(\frac{1}{1-xz} - {}_2F_1 \left(\begin{matrix} 1, q \\ p+q \end{matrix}; \frac{(y-x)z}{1-xz} \right) \right) > \frac{1}{2}$$

for all x, y and z such that $|x|=|y|=1$ and $|z| < 1$.

Here the hypergeometric function ${}_2F_1 \left(\begin{smallmatrix} a, b \\ c \end{smallmatrix}; z \right)$ is defined for $|z| < 1$ by

$$(2) \quad {}_2F_1 \left(\begin{matrix} a, b \\ c \end{matrix}; z \right) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k k!} z^k,$$

where $(a)_n = a(a+1)(a+2) \cdots (a+n-1), n > 0$, and $(a)_0 = 1$.

If $\operatorname{Re}(c) > \operatorname{Re}(b) > 0$, then the ${}_2F_1$ has an integral representation

$$(3) \quad {}_2F_1 \left(\begin{matrix} a, b \\ c \end{matrix}; z \right) = \frac{\Gamma(c)}{\Gamma(c-b)\Gamma(b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-tz)^{-a} dt.$$

This provides an analytic continuation of the ${}_2F_1$ to $\mathbb{C} - [1, \infty)$ under the restriction $\operatorname{Re}(c) > \operatorname{Re}(b) > 0$. Using some recurrence relations involving the parameters, the ${}_2F_1$ can be analytically continued to $\mathbb{C} - [1, \infty)$ for all a, b , and c such that $c \neq 0, -1, \dots$ (see Slater [3]).

Received by the editors February 16, 1982.

1980 *Mathematics Subject Classification*. Primary 33A30, 46J15.

Key words and phrases. F_p spaces, hypergeometric functions.

©1983 American Mathematical Society
0002-9939/82/0000-0865/\$01.50

Also we note the simple identity

$$(4) \quad {}_2F_1\left(\begin{matrix} a, b \\ a \end{matrix}; z\right) = (1-z)^{-b}, \quad a \neq 0, -1, \dots$$

The main result.

THEOREM 1. *If $F_{1+it} \cdot F_{1+it} \subseteq F_{2+2it}$, t real, then $t = 0$.*

PROOF. By Theorem A we must have

$$(5) \quad \operatorname{Re}\left(\frac{1}{1-xz} {}_2F_1\left(\begin{matrix} 1, 1+it \\ 2+2it \end{matrix}; \frac{(y-x)z}{1-xz}\right)\right) > \frac{1}{2}, \quad |x|=|y|=1, |z|<1.$$

We will assume $t \neq 0$ and obtain a contradiction. First we state a lemma proved by Slater [3, p. 34].

$$(6) \quad {}_2F_1\left(\begin{matrix} a, b \\ c \end{matrix}; z\right) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} {}_2F_1\left(\begin{matrix} a, b \\ a+b-c+1 \end{matrix}; 1-z\right) \\ + (1-z)^{c-a-b} \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} {}_2F_1\left(\begin{matrix} c-a, c-b \\ c-a-b+1 \end{matrix}; 1-z\right).$$

$c \neq 0, -1, -2, \dots$, $a+b-c \neq 0, \pm 1, \pm 2, \dots$, $|\operatorname{Arg}(z)| < \pi$, $|\operatorname{Arg}(1-z)| < \pi$.

The lemma is useful in that it reveals the behavior of a ${}_2F_1$ when $z \rightarrow 1$ with rather relaxed restrictions on the parameters a, b and c . It is clear from (6) that if $\operatorname{Re}(c-a-b) > 0$ then

$${}_2F_1\left(\begin{matrix} a, b \\ c \end{matrix}; z\right) \rightarrow \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} \quad \text{as } z \rightarrow 1.$$

If $\operatorname{Re}(c-a-b) < 0$ then the ${}_2F_1\left(\begin{smallmatrix} a, b \\ c \end{smallmatrix}; z\right)$ goes to infinity like $(1-z)^{c-a-b}$ as $z \rightarrow 1$. The case where $\operatorname{Re}(c-a-b) = 0$ can also be treated by (6) and that is what will make the proof of this theorem work.

Now let $z = r\bar{y}$, $0 < r < 1$, and $x\bar{y} = \xi \neq 1$. Then by (6),

$$(7) \quad {}_2F_1\left(\begin{matrix} 1, 1+it \\ 2+2it \end{matrix}; \frac{(y-x)z}{1-xz}\right) = {}_2F_1\left(\begin{matrix} 1, 1+it \\ 2+2it \end{matrix}; \frac{r(1-\xi)}{1-\xi r}\right) \\ = \frac{\Gamma(2+2it)\Gamma(it)}{\Gamma(1+2it)\Gamma(1+it)} {}_2F_1\left(\begin{matrix} 1, 1+it \\ 1-it \end{matrix}; \frac{1-r}{1-\xi r}\right) \\ + \left(\frac{1-r}{1-\xi}\right)^{it} \frac{\Gamma(2+2it)\Gamma(-it)}{\Gamma(1)\Gamma(1+it)} {}_2F_1\left(\begin{matrix} 1+2it, 1+it \\ 1+it \end{matrix}; \frac{1-r}{1-\xi r}\right).$$

Here the powers are computed using the principal value of the arguments, for this was assumed in (6). In order to simplify (7) we use (4) and the following identities [2, pp. 12-24].

$$(8) \quad \begin{aligned} & \text{(i)} \quad \Gamma(z+1) = z\Gamma(z), \quad z = 0, -1, -2, \dots, \\ & \text{(ii)} \quad \sqrt{\pi} \Gamma(2z) = 2^{2z-1} \Gamma(z), \quad z \neq 0, -\frac{1}{2}, -1, -\frac{3}{2}, \dots, \\ & \text{(iii)} \quad \Gamma(1/2) = \sqrt{\pi}. \end{aligned}$$

We then obtain

$$\begin{aligned}
 (9) \quad & \frac{1}{1-xz} {}_2F_1\left(\begin{matrix} 1, 1+it \\ 2+2it \end{matrix}; \frac{(y-x)z}{1-xz}\right) \\
 &= \frac{2-i/t}{1-\xi r} {}_2F_1\left(\begin{matrix} 1, 1+it \\ 1-it \end{matrix}; \frac{1-r}{1-\xi r}\right) \\
 &\quad + (1-r)'' \Gamma\left(\frac{3}{2}+it\right) \Gamma(-it) \left(\frac{1}{2}(1-\xi)r\right)^{-1-2it} (1-\xi r)'',
 \end{aligned}$$

$\xi = x\bar{y}$, $z = r\bar{y}$, $0 < r < 1$. The right side of (9) has the form

$$(10) \quad A(r, \xi) + (1-r)'' B(r, \xi),$$

where

$$\lim_{r \rightarrow 1^-} A(r, \xi) = \frac{2-i/t}{1-\xi}$$

and

$$(11) \quad \lim_{r \rightarrow 1^-} B(r, \xi) = \left(\frac{1}{2}(1-\xi)\right)^{-1-2it} (1-\xi)'' \Gamma\left(\frac{3}{2}+it\right) \Gamma(-it).$$

Note that as $r \rightarrow 1^-$, $(1-r)'' = \exp(it \operatorname{Ln}(1-r))$ is a complex number which revolves around the origin more and more rapidly while the modulus is one. Consequently, we must have

$$(12) \quad \operatorname{Re} A(r, \xi) > \frac{1}{2}, \quad 0 < r < 1, |\xi| = 1, \xi \neq 1.$$

Letting $r \rightarrow 1^-$, we have

$$(13) \quad \operatorname{Re}\left(\frac{2-i/t}{1-\xi}\right) > \frac{1}{2}.$$

But if we let $\xi = e^{i\theta}$

$$\operatorname{Re}\left(\frac{2-i/t}{1-\xi}\right) = 1 - \frac{(1+\cos\theta)}{2t \sin\theta}$$

and it is clear that a nonzero value of θ can be chosen so that the above expression is less than $\frac{1}{2}$. We have obtained a contradiction and the theorem is proved. $\square$

An obvious consequence of this theorem is

COROLLARY. *If $F_p \cdot F_q \subseteq F_{p+q}$ then $p > 0$ and $q > 0$.*

REFERENCES

1. K. Pearce, *A product theorem for $\mathfrak{S}_p$ classes and an application*, Proc. Amer. Math. Soc. 84 (1982), 509–515.
2. E. Rainville, *Special functions*, Macmillan, New York, 1960.
3. L. Slater, *Generalized hypergeometric functions*, Cambridge Univ. Press, Cambridge, 1966.

DEPARTMENT OF MATHEMATICS, MICHIGAN TECH UNIVERSITY, HOUGHTON, MICHIGAN 49931