THE STABILITY OF THE SINE EQUATION

PIOTR W. CHOLEWA

ABSTRACT. Let δ be a positive real constant and let G be an abelian group (written additively) in which division by 2 is uniquely performable. Every unbounded complex-valued function f on G satisfying the inequality

$$|f(x+y)f(x-y) - f(x)^2 + f(y)^2| \leq \delta \quad \text{for all } x, y \in G$$

has to be a solution of the sine functional equation

$$f(x+y)f(x-y) = f(x)^2 - f(y)^2 \quad \text{for all } x, y \in G.$$

A stability problem in functional equation theory seems to have been first raised by S. M. Ulam in [4] and D. H. Hyers who proved the stability of a linear functional equation (see [3]). Generalizations of this result appeared then in many papers. In 1979 J. A. Baker, J. Lawrence and F. Zorzitto showed in [2] that the equation $f(x+y) = f(x)f(y)$ has a different kind of stability. Exactly, they proved that if f is a real-valued function defined on a rational vector space V and satisfies the inequality $|f(x+y) - f(x)f(y)| \leq \delta$ for all $x, y \in V$ and some real $\delta > 0$, then either f is bounded or $f(x+y) = f(x)f(y)$ for all $x, y \in V$. This result was then generalized by J. A. Baker (see [1]) who also had obtained a similar result for the cosine equation $f(x+y) + f(x-y) = 2f(x)f(y)$. The main purpose of this paper is to prove an analogous result for the sine functional equation

$$(1) \quad f(x+y)f(x-y) = f(x)^2 - f(y)^2.$$

We shall start from some general assumptions. Let G be an abelian group (written additively) in which division by 2 is uniquely performable (i.e. for each $x \in G$ there exists a unique $y \in G$ such that $y+y=x$; such a y will be denoted by $x/2$). Let f be a complex-valued function defined on G and such that the inequality

$$(2) \quad |f(x+y)f(x-y) - f(x)^2 + f(y)^2| \leq \delta$$

holds for all $x, y \in G$ and some real $\delta > 0$. Moreover, assume that f is an unbounded function.

For the proof we need the following three lemmas.

LEMMA 1.

$$(3) \quad f(0) = 0.$$

Received by the editors August 9, 1982.

1980 *Mathematics Subject Classification*. Primary 39B40; Secondary 39B70, 39C05.

Key words and phrases. Functional equation, sine equation, stability.

©1983 American Mathematical Society
0002-9939/82/0000-1314/\$01.75

PROOF. Put $x = y$ and $u = 2x$ in inequality (2). Then

$$\delta \geq |f(u)f(0) - f(u/2)^2 + f(u/2)^2| = |f(u)| |f(0)|$$

and since $|f(u)|$ can be as large as desired, we must have

$$f(0) = 0. \quad \text{Q.E.D.}$$

Now put $x = (u + v)/2$, $y = (u - v)/2$ in (2). Then $x + y = u$, $x - y = v$ and inequality (2) assumes the form

$$(4) \quad \left| f(u)f(v) - f\left(\frac{u+v}{2}\right)^2 + f\left(\frac{u-v}{2}\right)^2 \right| \leq \delta \quad \text{for all } u, v \in G.$$

Since f is an unbounded function, there exists an $a \in G$ such that $|f(a)| \geq 4$. Let $g: G \rightarrow \mathbb{C}$ ($\mathbb{C}$ denotes here the field of all complex numbers) be the function defined by

$$(5) \quad g(x) = \frac{f(x+a) - f(x-a)}{2f(a)} \quad \text{for all } x \in G.$$

LEMMA 2. For all $x, y \in G$ the following inequality is satisfied:

$$(6) \quad |f(x+y) + f(x-y) - 2f(x)g(y)| \leq \delta.$$

PROOF. Let $x, y \in G$. Using (4) and (5) we get

$$\begin{aligned} & |f(x+y) + f(x-y) - 2f(x)g(y)| \\ &= \frac{1}{|f(a)|} |f(x+y)f(a) + f(x-y)f(a) - 2f(a)f(x)g(y)| \\ &\leq \frac{1}{|f(a)|} \left| f(x+y)f(a) - f\left(\frac{x+y+a}{2}\right)^2 + f\left(\frac{x+y-a}{2}\right)^2 \right| \\ &\quad + \frac{1}{|f(a)|} \left| f(x-y)f(a) - f\left(\frac{x-y+a}{2}\right)^2 + f\left(\frac{x-y-a}{2}\right)^2 \right| \\ &\quad + \frac{1}{|f(a)|} \left| f\left(\frac{x+y+a}{2}\right)^2 - f\left(\frac{x-y-a}{2}\right)^2 - f(x)f(y+a) \right| \\ &\quad + \frac{1}{|f(a)|} \left| f(x)f(y-a) - f\left(\frac{x+y-a}{2}\right)^2 + f\left(\frac{x-y+a}{2}\right)^2 \right| \\ &\quad + \left| 2f(x) \frac{f(y+a) - f(y-a)}{2f(a)} - 2f(x)g(y) \right| \\ &\leq \frac{4\delta}{|f(a)|} \leq \delta. \quad \text{Q.E.D.} \end{aligned}$$

LEMMA 3. The following equation is satisfied for all $x, y \in G$:

$$(7) \quad f(x+y) + f(x-y) = 2f(x)g(y).$$

PROOF. Let x, y be two arbitrarily fixed points of G . Then, using (4) and (6), for all $z \in G$ we have

$$\begin{aligned}
 & |f(z)| |f(x+y) + f(x-y) - 2f(x)g(y)| \\
 &= |f(z)f(x+y) + f(z)f(x-y) - 2f(x)f(z)g(y)| \\
 &\leq \left| f(z)f(x+y) - f\left(\frac{z+x+y}{2}\right)^2 + f\left(\frac{z-x-y}{2}\right)^2 \right| \\
 &\quad + \left| f(z)f(x-y) - f\left(\frac{z+x-y}{2}\right)^2 + f\left(\frac{z-x+y}{2}\right)^2 \right| \\
 &\quad + \left| f\left(\frac{z+x+y}{2}\right)^2 - f\left(\frac{z-x+y}{2}\right)^2 - f(z+y)f(x) \right| \\
 &\quad + \left| f\left(\frac{z+x-y}{2}\right)^2 - f\left(\frac{z-x-y}{2}\right)^2 - f(z-y)f(x) \right| \\
 &\quad + |[f(z+y) + f(z-y)]f(x) - 2f(z)g(y)f(x)| \\
 &\leq 4\delta + \delta|f(x)| = \delta(4 + |f(x)|).
 \end{aligned}$$

Therefore

$$|f(x+y) + f(x-y) - 2f(x)g(y)| \leq (4 + |f(x)|)\delta/|f(z)|,$$

and since f is assumed to be an unbounded function and x is a fixed element, $(4 + |f(x)|)\delta/|f(z)|$ is as small as we wish. Hence we must have

$$|f(x+y) + f(x-y) - 2f(x)g(y)| = 0,$$

whence

$$f(x+y) + f(x-y) = 2f(x)g(y).$$

Since x and y were arbitrarily fixed elements of G , equation (7) holds for all $x, y \in G$. Q.E.D.

Now we are able to prove the main result of our paper.

THEOREM. Every unbounded function $f: G \rightarrow \mathbb{C}$ satisfying inequality

$$|f(x+y)f(x-y) - f(x)^2 + f(y)^2| \leq \delta \quad \text{for all } x, y \in G$$

has to be a solution of the equation

$$f(x+y)f(x-y) = f(x)^2 - f(y)^2.$$

PROOF. Let f be an unbounded solution of inequality (2). Put $x = 0$ in (7). Then, using (3), we obtain

$$f(y) + f(-y) = 0 \quad \text{for all } y \in G,$$

i.e.

$$(8) \quad f(y) = -f(-y) \quad \text{for all } y \in G.$$

Now, put $x = (u+v)/2$, $y = (u-v)/2$ in (7). Hence

$$(9) \quad f(u) + f(v) = 2f\left(\frac{u+v}{2}\right)g\left(\frac{u-v}{2}\right) \quad \text{for all } u, v \in G.$$

Then, from (3) and (9), we infer that

$$(10) \quad f(x+y) = f(x+y) + f(0) = 2f\left(\frac{x+y}{2}\right)g\left(\frac{x+y}{2}\right) \quad \text{for all } x, y \in G,$$

and

$$(11) \quad f(x-y) = f(x-y) + f(0) = 2f\left(\frac{x-y}{2}\right)g\left(\frac{x-y}{2}\right) \quad \text{for all } x, y \in G.$$

Using (8) and (9) we also have

$$(12) \quad f(x) - f(y) = f(x) + f(-y) = 2f\left(\frac{x-y}{2}\right)g\left(\frac{x+y}{2}\right) \quad \text{for all } x, y \in G.$$

Now, using (9)–(12) we obtain

$$\begin{aligned} f(x+y)f(x-y) &= \left[2f\left(\frac{x+y}{2}\right)g\left(\frac{x+y}{2}\right)\right] \left[2f\left(\frac{x-y}{2}\right)g\left(\frac{x-y}{2}\right)\right] \\ &= \left[2f\left(\frac{x+y}{2}\right)g\left(\frac{x-y}{2}\right)\right] \left[2f\left(\frac{x-y}{2}\right)g\left(\frac{x+y}{2}\right)\right] \\ &= [f(x) + f(y)][f(x) - f(y)] = f(x)^2 - f(y)^2 \end{aligned}$$

for all $x, y \in G$. Q.E.D.

REMARK 1. J. A. Baker proved in [1] that, concerning the stability of the cosine equation, a function g satisfying inequality

$$|g(x+y) + g(x-y) - 2g(x)g(y)| \leq \delta$$

is either a solution of the cosine equation or it is bounded by a constant depending on δ only. The crucial step of the proof was that the function satisfying the above inequality is either bounded by such a constant or unbounded. It is not the case for the sine equation; indeed, the (bounded) functions

$$f_N(x) = N \sin x + 1/N$$

satisfy inequality (2) with $\delta = 3$,

$$|f_N(x+y)f_N(x-y) - f_N(x)^2 + f_N(y)^2| \leq 3$$

for all real numbers x, y and all positive integers N . Nevertheless, for each positive real number M , the inequality $|f_N(x)| \leq M$ fails to hold for certain x and N .

REMARK 2. We must emphasize that our main result is proved for complex-valued functions only. Essential difficulties appear while considering functions with values in the algebra of quaternions or Cayley numbers, despite the multiplicativity of the norm; associativity, as well as commutativity, was necessary for us to prove our theorem.

REFERENCES

1. J. A. Baker, *The stability of the sine equation*, Proc. Amer. Math. Soc. **80** (1980), 411–416.
2. J. A. Baker, J. Lawrence and F. Zorzitto, *The stability of the equation $f(x+y) = f(x)f(y)$* , Proc. Amer. Math. Soc. **74** (1979), 242–246.
3. D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
4. S. M. Ulam, *A collection of mathematical problems*, Interscience, New York, 1960.

INSTITUTE OF MATHEMATICS, SILESIA UNIVERSITY, BANKOWA 14, 40-007 KATOWICE, POLAND