ON THE ALEXANDER POLYNOMIAL OF A CYCLICALLY PERIODIC KNOT

JONATHAN A. HILLMAN

ABSTRACT. We show that a theorem of Burde on the Alexander polynomial of a cyclically periodic knot K , such that $\Delta_2(K) = 1$, may be extended to all cyclically periodic knots.

A knot K in a homology 3-sphere Σ has cyclic period q if it is invariant under an automorphism of Σ of order q which has fixed point set a circle disjoint from K . In [3] we gave new proofs of theorems of Burde [1] and Murasugi [5] on the Alexander polynomial $\Delta_1(K)$ of such a knot.

A refinement of the second lemma of [3] enables us to remove the restriction $\Delta_2(K) = 1$ from Burde's theorem. We shall adopt the notation of [3].

THEOREM. *Let K be a knot in an homology 3-sphere Σ which has cyclic period q , where $q = p^r$ for some prime p . Then either*

- (i) $\Delta_1(K) \equiv 1 \pmod{p}$ or
- (ii) *the degree of the q th roots of unity over the splitting fields of $\Delta_1(K)$ is at most n , where $\Delta_n(K) \neq 1$ but $\Delta_{n+1}(K) = 1$.*

PROOF. The argument given in [3] shows that if (i) fails then the $\mathbb{Q}[t, t^{-1}]$ -module $C = (G'/G'') \otimes_{\mathbb{Z}} \mathbb{Q}$ (the rational homology of the infinite cyclic cover of the knot complement) has an automorphism of order exactly q . (For otherwise the (q/p) th power of the given periodic self-homeomorphism τ of Σ induces the identity on G'/G'' . Applying Milnor duality and the Wang sequence with coefficients $\mathbb{Z}/p\mathbb{Z}$ to the p -fold branched cyclic covering map from Z to $\bar{Z}$, the results of 0-framed surgery on K in Σ and on the image of K in $\Sigma/\tau^{(q/p)}$, respectively, we conclude that $(G'/G'') \otimes (\mathbb{Z}/p\mathbb{Z})$ must be 0, and so $\Delta_1(K) \equiv 1 \pmod{p}$.) Then for some irreducible factor δ of $\Delta_1(K)$, the δ -primary submodule of C has an automorphism τ of order exactly q , so we may assume $\Delta_0(C) = \delta^l$. Let $L = \mathbb{Q}[t, t^{-1}]/(\delta)$. If $m \geq 1$ the kernel of the natural map from $\text{Aut}(C/\delta^{m+1}C)$ to $\text{Aut}(C/\delta^m C)$ is isomorphic to $\text{Hom}(C/\delta^{m+1}C, \delta^m C/\delta^{m+1}C)$ and so is an L -vector space. In particular, it is torsion free. Therefore the image of τ in $\text{Aut}(C/\delta C) \approx \text{GL}(n, L)$ has order exactly q . Let $A = L[\tau]$ be the subalgebra of $M(n, L)$ generated by this element. Then $\dim_L A \leq n$ by the Cayley-Hamilton theorem. On the other hand, the image of τ in the semisimple algebra $A/\text{rad } A$ again has order exactly q , as the kernel of the map on

Received by the editors December 15, 1981.

1980 *Mathematics Subject Classification.* Primary 57M25.

Key words and phrases. Alexander polynomial, cyclic period, knot, root of unity, splitting field.

©1983 American Mathematical Society
0002-9939/82/0000-2055/\$01.50

units is again an iterated extension of L -vector spaces. The algebra $A/\text{rad } A$ is a product of fields containing L , so some factor is an extension L' of L which contains the q th roots of unity and for which $[L': L] = \dim_L L' \leq \dim_L A \leq n$. Since L is contained in the splitting field for $\Delta_1(K)$, the theorem follows.

REMARKS. Suppose (ii) holds. If $q = p^r$, with $r > 1$, then either $L = L'$, and so contains the q th roots of unity, or $[L': L] \geq p$, and so $p \leq n$. If $q = p$ and L is linearly disjoint from the field $Q(\xi)$ of p th roots of unity, that is if $L \cap Q(\xi) = Q$, then $[L': L] = p - 1$, so $p \leq n + 1$.

If $\Delta_2(K) = 1$, so that $n = 1$, we obtain Burde's theorem [1] (itself an extension of a result of Trotter [8]). Burde's version applies to all but four of the prime knots with ten crossings or less (excepting 8_{18} , 9_{40} , 10_{99} and 10_{123}), but does not usefully apply to the n -fold connected sum of a knot K with itself, for $\Delta_2(\#^n K)$ is nontrivial unless $\Delta_1(K) = 1$ or $n = 1$.

For other recent work on knots with cyclic periods see [2, 3, 4, 6, 7]. (Note that there are some discrepancies between the tables of [4] and [6].)

REFERENCES

1. G. Burde, *Über periodische knoten*, Arch. Math. (Basel) **30** (1978), 487–492.
2. C. McA. Gordon, R. A. Litherland and K. Murasugi, *Signatures of covering links*, Canad. J. Math. **33** (1981), 381–394.
3. J. A. Hillman, *New proofs of two theorems on periodic knots*, Arch. Math. (Basel) **37** (1981), 457–461.
4. U. Lüdicke, *Zyklische knoten*, Arch. Math. (Basel) **32** (1979), 588–599.
5. K. Murasugi, *On periodic knots*, Comment. Math. Helv. **46** (1971), 162–174.
6. ———, *On symmetry of knots*, Tsukuba J. Math. **4** (1980), 331–347.
7. M. Sakuma, *On the polynomials of periodic links*, Math. Ann. **257** (1981), 487–494.
8. H. F. Trotter, *Periodic automorphisms of groups and knots*, Duke Math. J. **28** (1961), 553–558.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, AUSTIN, TEXAS 78712

Current address: Department of Mathematics, Faculty of Science, The Australian National University, Canberra 2600, Australia