CARDINAL FUNCTIONS ON PIXLEY-ROY HYPERSPACES

SHOZO SAKAI

ABSTRACT. Some cardinal functions on the Pixley-Roy hyperspace $\mathcal{F}[X]$ of a space X are determined by those on X , and conditions of X by cardinal functions from which $\mathcal{F}[X]$ has or has not some properties, e.g. cosmic, paracompact, ccc, etc. are given.

All spaces considered in this paper will be assumed to be infinite T_1 -spaces. Cardinals are the initial ordinals, τ will always denote an infinite cardinal and ω is the smallest infinite cardinal. The cardinality of a set X is denoted by $|X|$ and c is the cardinality of the continuum. The successor cardinal of τ is denoted by τ^+ .

The *Pixley-Roy hyperspace* $\mathcal{F}[X]$ of a space X has as its underlying set the collection of all finite nonempty subsets of X . If $A \in \mathcal{F}[X]$, then the basic nbds (= neighbourhoods) of A are $[A, U] = \{S \in \mathcal{F}[X]: A \subset S \subset U\}$, where U is an open set of X containing A .

We refer to [J] for the following cardinal functions— w (*weight*), π (π -*weight*), nw (*net weight*), d (*density*), c (*cellularity*), L (*Lindelöf-degree*), p (= e (*extent*) in [E]), χ (*character*), ψ (*pseudo character*), Ψ ($\Psi(X) = \omega$ iff X is perfect), ψ_Δ (*diagonal degree*), $\pi\chi$ (π -*character*), t (*tightness*) and psw (*point separating weight*). However, following Engelking [E], the *hereditary density* and the *hereditary Lindelöf-degree* are denoted by hd and hL , respectively.

In the following, Theorem 2 is the main result of this paper. Some results in it were given by van Douwen [vD], Lutzer [L] and Przymusiński [P]. Though Proposition 1 is an auxiliary one for Theorem 2, it is interesting in itself.

PROPOSITION 1. *If $A \in \mathcal{F}[X]$, then the following hold:*

- (1) $\psi(A, \mathcal{F}[X]) = \max\{\psi(x, X): x \in A\}$.
- (2) $\pi\chi(A, \mathcal{F}[X]) = \chi(A, \mathcal{F}[X]) \leq \max\{\chi(x, X): x \in A\}$.

If $x \in X$, then the following hold:

- (3) $\chi(\{x\}, \mathcal{F}[X]) = \chi(x, X)$.
- (4) $t(x, X) \leq t(\{x\}, \mathcal{F}[X])$.

If X is Hausdorff and $A \in \mathcal{F}[X]$, then the following holds:

- (5) $\chi(A, \mathcal{F}[X]) = \max\{\chi(x, X): x \in A\}$.

Received by the editors November 5, 1982 and, in revised form, February 16, 1983.

1980 *Mathematics Subject Classification*. Primary 54A25, 54B20; Secondary 54D18, 54D20.

©1983 American Mathematical Society
0002-9939/83 \$1.00 + \$.25 per page

PROOF. (1) Let $A = \{x_1, \dots, x_n\} \in \mathcal{F}[X]$. For each $i \in \{1, \dots, n\}$, there is a local ψ -base $\mathcal{U}_i$ of x_i in X such that $|\mathcal{U}_i| \leq \psi(x_i, X)$. Let

$$\mathcal{U} = \left\{ \bigcup \{U_i : i = 1, \dots, n\} : U_i \in \mathcal{U}_i \right\} \quad \text{and} \quad \mathcal{B} = \{[A, U] : U \in \mathcal{U}\}.$$

Then $\mathcal{B}$ is a local ψ -base of A in $\mathcal{F}[X]$ and $|\mathcal{B}| \leq \max\{\psi(x_i, X) : i = 1, \dots, n\}$. Thus, we have $\psi(A, \mathcal{F}[X]) \leq \max\{\psi(x, X) : x \in A\}$. Conversely, let $\mathcal{B}$ be a local ψ -base of A in $\mathcal{F}[X]$ and $x \in A$. Then $\mathcal{U} = \{B \setminus (A \setminus \{x\}) : [A, B] \in \mathcal{B}\}$ is a local ψ -base of x in X . Therefore, we have $\psi(x, X) \leq \psi(A, \mathcal{F}[X])$ and $\max\{\psi(x, X) : x \in A\} \leq \psi(A, \mathcal{F}[X])$.

(2) Let $\mathcal{B}$ be a local π -base of A in $\mathcal{F}[X]$. For each $B \in \mathcal{B}$, we can take an element $A_B \in B$ and an open nbd U_B of A_B in X such that $[A_B, U_B] \subset B$. Then, $\mathcal{U} = \{[A, U_B] : B \in \mathcal{B}, A \subset U_B\}$ is a nbd base of A in $\mathcal{F}[X]$. For, let V be an open nbd of A in X . Then, there is $B \in \mathcal{B}$ such that $B \subset [A, V]$. Therefore, $[A_B, U_B] \subset B \subset [A, V]$, i.e. $A \subset A_B \subset U_B \subset V$. Thus, $[A, U_B] \subset [A, V]$ and $\mathcal{U}$ is a nbd base of A in $\mathcal{F}[X]$. Since $|\mathcal{U}| \leq |\mathcal{B}|$, we have $\chi(A, \mathcal{F}[X]) \leq \pi\chi(A, \mathcal{F}[X])$. $\pi\chi(A, \mathcal{F}[X]) \leq \chi(A, \mathcal{F}[X])$ holds always and the other part is proved similar to (1).

(3) The proof is clear.

(4) Let $H \subset X$, $x \in X \setminus H$ and $x \in \text{cl}_X H$. Let us put $\mathcal{K} = \mathcal{F}[H \cup \{x\}] \setminus \{\{x\}\}$. Then $\{x\} \in \text{cl}_{\mathcal{F}[X]} \mathcal{K}$. Therefore, there is a subset $\mathcal{M}$ of $\mathcal{K}$ such that $|\mathcal{M}| \leq t(\{x\}, \mathcal{F}[X])$ and $\{x\} \in \text{cl}_{\mathcal{F}[X]} \mathcal{M}$. Let $M = (\bigcup \mathcal{M}) \setminus \{x\}$. Then $x \in \text{cl}_X M$. For, let V be a nbd of x in X . Since $\{x\} \in \text{cl}_{\mathcal{F}[X]} \mathcal{M}$, there is an element $A \in [\{x\}, V] \cap \mathcal{M}$. Then, $x \in A \subset V$ and $A \in \mathcal{M}$. Since $A \neq \{x\}$, we have $V \cap M \neq \emptyset$ and $x \in \text{cl}_X M$. Since $M \subset H$ and $|M| \leq |\mathcal{M}| \leq t(\{x\}, \mathcal{F}[X])$, we have $t(x, X) \leq t(\{x\}, \mathcal{F}[X])$.

(5) Let us assume that there is $x \in A$ with $\chi(A, \mathcal{F}[X]) < \chi(x, X)$. Let $\tau = \chi(A, \mathcal{F}[X])$ and $\mathcal{U}$ a nbd base of A in $\mathcal{F}[X]$ with $|\mathcal{U}| \leq \tau$. We can assume $\mathcal{U} = \{[A, U_\alpha] : \alpha < \tau\}$, where U_α is an open set of X containing A . Since X is Hausdorff, there is an open nbd U of x with $(A \setminus \{x\}) \cap \text{cl}_X U = \emptyset$. Since $\chi(x, X) > \tau$, there is an open nbd V of x with $U \cap U_\alpha \not\subset V$ for each $\alpha < \tau$. Let $W = V \cup (X \setminus \text{cl}_X U)$. Then $[A, W]$ is an open nbd of A in $\mathcal{F}[X]$. Since $U_\alpha \not\subset W$, $[A, U_\alpha] \not\subset [A, W]$ for each $\alpha < \tau$. Therefore, $\mathcal{U}$ is not a nbd base of A , a contradiction. Thus, we have $\chi(x, X) \leq \chi(A, \mathcal{F}[X])$ for each $x \in A$. This completes the proof of Proposition 1.

THEOREM 2. For each space X , the following hold:

- (1) $|\mathcal{F}[X]| = nw(\mathcal{F}[X]) = d(\mathcal{F}[X]) = L(\mathcal{F}[X]) = p(\mathcal{F}[X]) = |X|$.
- (2) $\chi(\mathcal{F}[X]) = \pi\chi(\mathcal{F}[X]) = \chi(X)$.
- (3) $w(\mathcal{F}[X]) = \pi(\mathcal{F}[X]) = \chi(X) |X|$.
- (4) $psw(\mathcal{F}[X]) = \psi(\mathcal{F}[X]) = \psi(X)$.
- (5) $\psi(X) = \omega$ iff $\Psi(\mathcal{F}[X]) = \omega$ iff $\psi_\Delta(\mathcal{F}[X]) = \omega$.
- (6) $hd(X)hL(X) \leq c(\mathcal{F}[X]) \leq nw(X)$.

PROOF. (1) Since $\{\{x\} : x \in X\}$ is a closed discrete subspace of $\mathcal{F}[X]$, $|X| \leq p(\mathcal{F}[X])$. $|\mathcal{F}[X]| = d(\mathcal{F}[X]) = |X|$ appears in [vD]. Therefore, (1) is clear.

(2) By Proposition 1, this is trivial.

(3) From

$$\chi(X) = \pi\chi(\mathcal{F}[X]) \leq \pi(\mathcal{F}[X]) \quad \text{and} \quad |X| = d(\mathcal{F}[X]) \leq \pi(\mathcal{F}[X]),$$

we have $\chi(X)|X| \leq \pi(\mathcal{F}[X]) \leq w(\mathcal{F}[X])$. On the other hand,

$$w(\mathcal{F}[X]) \leq \chi(\mathcal{F}[X])|\mathcal{F}[X]|$$

is always true. Therefore, by (1) and (2), (3) is proved.

(4) Since $\psi(\mathcal{F}[X]) \leq psw(\mathcal{F}[X])$ is well known, we will prove $psw(\mathcal{F}[X]) \leq \psi(X)$ only. For each $A \in \mathcal{F}[X]$, there is a local ψ -base $\mathcal{U}(A)$ of A with $|\mathcal{U}(A)| \leq \psi(\mathcal{F}[X]) = \psi(X)$. Then, $\mathcal{B} = \{[A, U]: A \in \mathcal{F}[X], U \in \mathcal{U}(A)\}$ is a pseudo base of $\mathcal{F}[X]$. Let $S \in \mathcal{F}[X]$. $S \in [A, U]$ iff $A \subset S \subset U$. Since $|S| < \omega$, $\{A: \emptyset \neq A \subset S\}$ is finite. Therefore, $\text{ord}(S, \mathcal{B}) = |\{[A, U] \in \mathcal{B}: S \in [A, U]\}| \leq \psi(X)$. Thus we have $\text{ord}(\mathcal{B}) = \sup\{\text{ord}(S, \mathcal{B}): S \in \mathcal{F}[X]\} \leq \psi(X)$. Therefore, $psw(\mathcal{F}[X]) \leq \psi(X)$.

(5) $\psi(X) = \omega$ iff $\Psi(\mathcal{F}[X]) = \omega$ appears in [L] and [P], and $\psi(\mathcal{F}[X]) \leq \psi_\Delta(\mathcal{F}[X])$ is well known. Therefore, only $\psi_\Delta(\mathcal{F}[X]) \leq \psi(\mathcal{F}[X]) = \omega$ needs proof. Let $\psi(\mathcal{F}[X]) = \omega$ and $\mathcal{U}(A) = \{U_n(A): n < \omega\}$ be a decreasing countable local ψ -base of each $A \in \mathcal{F}[X]$ in $\mathcal{F}[X]$. We can assume that $U_n(A)$ is of the form $[A, V]$, where V is an open nbd of A in X . Since $B_n = \bigcup \{U_n(A) \times U_n(A): A \in \mathcal{F}[X]\}$ is an open nbd of the diagonal $\Delta(\mathcal{F}[X]) = \{(A, A): A \in \mathcal{F}[X]\}$ in $\mathcal{F}[X]^2 = \mathcal{F}[X] \times \mathcal{F}[X]$, $\mathcal{B} = \{B_n: n < \omega\}$ is a countable local ψ -base of $\Delta(\mathcal{F}[X])$ in $\mathcal{F}[X]^2$. For, let $(S, T) \in \mathcal{F}[X]^2 \setminus \Delta(\mathcal{F}[X])$. We may assume that there is an element $t \in T \setminus S$. For each nonempty $A \subset S$, there exists $n(A) < \omega$ such that $\{t\} \notin U_{n(A)}(A)$. If we put $n_0 = \max\{n(A): \emptyset \neq A \subset S\}$, then $T \notin \bigcup \{U_n(A): \emptyset \neq A \subset S\}$ for each $n \geq n_0$. Therefore, $(S, T) \notin B_{n_0}$ and $\mathcal{B}$ is a local ψ -base of $\Delta(\mathcal{F}[X])$ in $\mathcal{F}[X]^2$. Therefore, we have $\psi_\Delta(\mathcal{F}[X]) = \omega$.

(6) Since $c(\mathcal{F}[X]) \leq nw(X)$ appears in [L], it suffices to show $hd(X)hL(X) \leq c(\mathcal{F}[X])$.

Let us assume $hL(X) > \tau$. Then, there are a subspace Y of X and an open covering $\mathcal{G}$ of Y such that $Y \not\subset \bigcup \mathcal{G}'$ for each subfamily $\mathcal{G}'$ of $\mathcal{G}$ with $|\mathcal{G}'| \leq \tau$. We can assume that each element of $\mathcal{G}$ is open in X . By transfinite induction, we can easily construct $\{x_\alpha: \alpha < \tau^+\} \subset Y$ and $\{G_\alpha: \alpha < \tau^+\} \subset \mathcal{G}$ with $x_\alpha \in G_\alpha \setminus (\bigcup \{G_\beta: \beta < \alpha\})$ for each $\alpha < \tau^+$. If $\beta < \alpha < \tau^+$, since $x_\alpha \notin G_\beta$, we have $[\{x_\beta\}, G_\beta] \cap [\{x_\alpha\}, G_\alpha] = \emptyset$. Thus, $\{[\{x_\alpha\}, G_\alpha]: \alpha < \tau^+\}$ is a pairwise disjoint family of nonempty open sets of $\mathcal{F}[X]$. Thus, we have $c(\mathcal{F}[X]) \geq \tau^+$ and $hL(X) \leq c(\mathcal{F}[X])$.

Let Y be a subspace of X . Consider the system $\mathcal{U}$ of all sets of the form $[A, U]$ where $A \in \mathcal{F}[Y]$ and U is an open set of X containing A . Let $\mathcal{B}$ be a pairwise disjoint maximal family of sets in $\mathcal{U}$. Then $D = \bigcup \{A: [A, U] \in \mathcal{B}\}$ is dense in Y . Because, assume there is $x \in Y \setminus \text{cl}_X D$. Then, there is an open nbd V of x in X such that $V \cap D = \emptyset$. Since $[\{x\}, V] \cap [A, U] = \emptyset$ for each $[A, U] \in \mathcal{B}$, $\mathcal{B} \cup \{[\{x\}, V]\}$ is a pairwise disjoint family of sets in $\mathcal{U}$. This contradicts the maximality of $\mathcal{B}$. Since $|D| \leq |\mathcal{B}| \leq c(\mathcal{F}[X])$, we have $d(Y) \leq c(\mathcal{F}[X])$. Thus, $hd(X) \leq c(\mathcal{F}[X])$ is proved. This completes the proof of Theorem 2.

Hajnal and Juhász studied conditions when $\mathcal{F}[X]$ has countable cellularity. Readers, who take interest in this case, refer to [HJ].

Since $\mathcal{F}[X]$ is regular [vD], we have the following:

COROLLARY 3. *In the following, (1)–(5) are equivalent, and each of them implies (6).*

(1) X is countable.

(2) $\mathcal{F}[X]$ is countable.

- (3) $\mathfrak{F}[X]$ is a cosmic space.
- (4) $\mathfrak{F}[X]$ is (hereditarily) separable.
- (5) $\mathfrak{F}[X]$ is (hereditarily) Lindelöf.
- (6) $\mathfrak{F}[X]$ is paracompact.

LEMMA 4. If X is a paracompact Hausdorff space, then $L(X) \leq c(X)$.

PROOF. Since each open covering of X has a σ -discrete open refinement, the lemma is clear.

COROLLARY 5. If $nw(X) < |X|$ holds, then $\mathfrak{F}[X]$ is not paracompact. In particular, if X is an uncountable separable metric space, then $\mathfrak{F}[X]$ is not metrizable.

The following corollaries are also clear.

COROLLARY 6. If $\mathfrak{F}[X]$ has G_δ -diagonal, then the pseudo character of X is ω .

COROLLARY 7. If $\mathfrak{F}[X]$ has countable cellularity, then X is both hereditarily separable and hereditarily Lindelöf.

THEOREM 8. Let ϕ be a cardinal function listed above except psw , ψ_Δ and Ψ . Then, we have $\phi(X) \leq \phi(\mathfrak{F}[X])$ for each space X .

PROOF. $\iota(X) \leq \iota(\mathfrak{F}[X])$ follows from Proposition 1. Proofs for other cardinal functions follow from Theorem 2.

EXAMPLE 1. By (4) in Theorem 2, we have always $psw(\mathfrak{F}[X]) \leq psw(X)$. There is a compact Hausdorff GO-space X for which $psw(\mathfrak{F}[X]) < psw(X)$, $\psi_\Delta(\mathfrak{F}[X]) < \psi_\Delta(X)$ and $\Psi(\mathfrak{F}[X]) < \Psi(X)$ hold. For example, let X be the lexicographic ordered square [E, 3.12.3]. Then $psw(X) = \psi_\Delta(X) = \Psi(X) = c > \omega = \psi(X)$. Therefore, we have $psw(\mathfrak{F}[X]) = \psi_\Delta(\mathfrak{F}[X]) = \Psi(\mathfrak{F}[X]) = \omega$ by Theorem 2.

EXAMPLE 2. There is a compact Hausdorff GO-space X for which $hd(X)hL(X) < c(\mathfrak{F}[X]) = nw(X)$ holds. In fact, let X be the two arrows space [E, 3.10.C]. That is, $X = \{(x, 0) : 0 < x \leq 1\} \cup \{(x, 1) : 0 \leq x < 1\}$ considered as a subspace of the lexicographic ordered square. Then, $hd(X) = hL(X) = \omega < c = nw(X)$. Let us take a point $P(x) = (x, 0)$ in X . Since $U(x) = \{(x, 0), (0, 1)\} \cup \{(y, i) : 0 < y < x, i = 0, 1\}$ is an open nbd of $P(x)$, $\{\{P(x)\}, U(x)\} : 0 < x \leq 1\}$ is a disjoint family of open sets of $\mathfrak{F}[X]$. Therefore, we have $c(\mathfrak{F}[X]) = c$.

EXAMPLE 3. There is a T_1 -space X which has a finite subset A such that $\chi(A, \mathfrak{F}[X]) < \max\{\chi(x, X) : x \in A\}$ holds. Let X be the set of all lattice points (i, j) of positive integers with two ideal points p and q . The topology of X is defined by declaring each lattice point to be open, and by taking as open nbds of p sets of the form $X \setminus (B \cup \{q\})$ where B is any set of lattice points with at most finitely many points on each row, and as open nbds of q sets of the form $X \setminus (C \cup \{p\})$ where C is any set of lattice points selected from at most finitely many rows. Then X is a compact T_1 -space and $\chi(p, X) > \omega$ holds [SS, 99]. Let $A = \{p, q\}$. If U is a nbd of A , then $|X \setminus U| < \omega$. Therefore, $\mathfrak{B} = \{[A, U] : A \subset U \subset X, |X \setminus U| < \omega\}$ is a nbd base of A in $\mathfrak{F}[X]$. Since $|\mathfrak{B}| = \omega$, we have $\chi(A, \mathfrak{F}[X]) < \chi(p, X)$.

Concerning theorems and examples, we ask the following.

Question 1. Let $\phi \in \{\Psi, \psi_\Delta\}$. Does $\phi(\mathfrak{F}[X]) \leq \phi(X)$ hold for each space X ?

Question 2. Determine exactly t , Ψ and ψ_Δ on $\mathcal{F}[X]$ in terms of those on X . Lutzer asked this question for the cellularity c [L].

REFERENCES

- [vD] E. K. van Douwen, *The Pixley-Roy topology on spaces of subsets*, Set-theoretic Topology, Academic Press, New York, 1977, pp. 111–134.
- [E] R. Engelking, *General topology*, PWN, Warsaw, 1977.
- [HJ] A. Hajnal and I. Juhász, *When is a Pixley-Roy hyperspace CCC?*, Topology Appl. **13** (1982), 33–41.
- [J] I. Juhász, *Cardinal functions in topology—ten years later*, Math. Centre Tracts, No. 123, North-Holland, Amsterdam, 1980.
- [L] D. J. Lutzer, *Pixley-Roy topology*, Topology Proc. **3** (1978), 139–158.
- [P] T. C. Przymusiński, *Normality and paracompactness of Pixley-Roy hyperspaces*, Fund. Math. **113** (1981), 201–219.
- [SS] L. A. Steen and J. A. Seebach, Jr., *Counterexamples in topology*, Springer-Verlag, Berlin and New York, 1978.

FACULTY OF LIBERAL ARTS, SHIZUOKA UNIVERSITY, OHYA, SHIZUOKA, 422, JAPAN