A COMPOSITION THEOREM FOR δ -CODES

C. H. YANG

ABSTRACT. If Golay complementary sequences (or equivalently a two-symbol δ -code) of length n and a Turyn δ -code of length t exist then four-symbol δ -codes of length $(2n + 1)t$ can be composed. Therefore new families of Hadamard matrices of orders $4uw$ and $20uw$ can be constructed, where $u = (2^{a+1}10^b26^c + 1)t$ for odd $t \leq 59$ or $t = 2^d10^e26^f + 1$ (all a, b, c, d, e , and $f \geq 0$), and w is the order of Williamson matrices.

The following result has been obtained: If Golay complementary sequences of length s and Turyn base sequences for length t (see [1]) exist, then four-symbol δ -codes of length $(2s + 1)t$ can be composed. Consequently four-symbol δ -codes of length $u = (2^{a+1}10^b26^c + 1)t$ can be constructed for odd $t \leq 59$ or $t = 2^d10^e26^f + 1$, where a, b, c, d, e and f are nonnegative integers. Therefore new families of Hadamard matrices of orders $4uw$ and $20uw$ can be constructed, where w is the order of Williamson matrices.

In this paper, many notations and definitions are the same as in [1]. *Turyn base sequences for length $t = 2m + p$* (abbreviated as $TBS(t)$) are four $(1, -1)$ -sequences $(A, B; C, D)$, respectively, of lengths $m + p$ and m pairwise such that

$$(1) \quad A = (a_k)_{m+p}, \quad B = (b_k)_{m+p}, \quad C = (c_k)_m \quad \text{and} \quad D = (d_k)_m$$

having zero auto-correlation sum, i.e. $a(j) + b(j) + c(j) + d(j) = 0$ for $j \neq 0$, where $p(j) = \sum_{k=1}^{n-j} p_k p_{k+j}$ for $P = (p_k)_n$. Another characterization of $TBS(t)$ is that the associated polynomials satisfy

$$(2) \quad |A|^2 + |B|^2 + |C|^2 + |D|^2 = 2t, \quad \text{for any } z \in \mathbf{K} = \{z \in \mathbf{C} : |z| = 1\},$$

where we use the same notation P to represent a given sequence $(p_k)_n$ and its associated polynomial $P(z) = \sum_{k=1}^n p_k z^{k-1}$; $\mathbf{K}$ is the unit circle and $\mathbf{C}$ is the complex field.

Turyn sequences of length t (abbreviated as $TS(t)$) (or a *four-symbol δ -code of length t*) are four $(0, \pm 1)$ -sequences (I, J, K, L) of length t , where $I = (i_h)_t$, $J = (j_h)_t$, $K = (k_h)_t$ and $L = (l_h)_t$, satisfying $i(h) + j(h) + k(h) + l(h) = 0$ for $h \neq 0$ and $|i_h| + |j_h| + |k_h| + |l_h| = 1$ for each h . It is known that the associated polynomials $TS(t)$ satisfy $|I|^2 + |J|^2 + |K|^2 + |L|^2 = t$ for any z in $\mathbf{K}$.

A *regular δ -code of length u* (abbreviated as $RD(u)$) is a quad (Q, R, S, T) of $(0, \pm 1)$ -sequences of length u such that

$$(3) \quad Q = I + J, \quad R = I - J, \quad S = K + L \quad \text{and} \quad T = K - L,$$

Received by the editors August 5, 1982 and, in revised form, August 23, 1982.

1980 *Mathematics Subject Classification*. Primary 05B20, 62K05, 05A19.

©1983 American Mathematical Society
0002-9939/83 \$1.00 + \$.25 per page

which is derived from $TS(u)$: (I, J, K, L) . The existence of $RD(u)$ and that of $TS(u)$ are equivalent since, from (3), we also have $I = (Q + R)/2$, $J = (Q - R)/2$, $K = (S + T)/2$ and $L = (S - T)/2$. We note here that the associated polynomials of $RD(u)$ satisfy

$$(4) \quad |Q|^2 + |R|^2 + |S|^2 + |T|^2 = 2u \quad \text{for any } z \text{ in } \mathbf{K},$$

and

(5) either $q_k r_k = \pm 1$ and $s_k = t_k = 0$, or $q_k = r_k = 0$ and $s_k t_k = \pm 1$, for each k , where q_k , r_k , s_k , and t_k are the k th term of the sequences Q , R , S , and T , respectively.

We first prove the following *Lagrange identity theorem for polynomials*, which is the key to our constructions for $RD(u)$.

THEOREM 1. *Let A, B, C, D, E, F, G and H be polynomials in $z \in \mathbf{K}$ with real coefficients. Also let*

$$(6) \quad \begin{aligned} W &= -B'E + AF' + CG + DH, & Y &= -D'E - CF + AG' - BH, \\ X &= A'E + BF' + DG' - CH', & Z &= C'E - DF + BG + AH', \end{aligned}$$

where $P' = P(z^{-1})$ for $P = P(z)$. Then

$$(7) \quad |W|^2 + |X|^2 + |Y|^2 + |Z|^2 = (|A|^2 + |B|^2 + |C|^2 + |D|^2)(|E|^2 + |F|^2 + |G|^2 + |H|^2).$$

PROOF. Since

$$(8) \quad |W|^2 + |X|^2 + |Y|^2 + |Z|^2 = WW' + XX' + YY' + ZZ' \quad \text{and}$$

$$(9) \quad \begin{aligned} W' &= -BE' + A'F + C'G' + D'H', \\ X' &= AE' + B'F + D'G - C'H, \\ Y' &= -DE' - C'F' + A'G - B'H', \\ Z' &= CE' - D'F' + B'G' + A'H, \end{aligned}$$

by substituting the right-hand sides of (6) and (9) into that of (8), and by expansions, simplifications and regrouping, (7) can be derived.

Golay complementary sequences of length s (abbreviated as $GCL(s)$) are two $(1, -1)$ -sequences: $\mathbf{F} = (f_k)_s$ and $\mathbf{G} = (g_k)_s$ having zero auto-correlation sum, $f(j) + g(j) = 0$ for $j \neq 0$. Another characterization of $GCL(s)$ (see [2]) is that the associated polynomials satisfy $|\mathbf{F}|^2 + |\mathbf{G}|^2 = 2s$ for any z in $\mathbf{K}$. $GCL(s)$ are known to exist for $s = 2^a 10^b 26^c$, where a, b , and c are nonnegative integers (see [3]).

THEOREM 2. *Let $(A, B; C, D)$ be $TBS(t)$ of (1) and $\mathbf{F} = (f_k)_s$ and $\mathbf{G} = (g_k)_s$ be $GCL(s)$, where $s = 2r$. Then the following (Q, R, S, T) are the associated polynomials of $RD((2s + 1)t)$.*

$$(*) \quad \begin{aligned} Q &= (-B'E + AF' + CG)U, & S &= (-D'E - CF + AG')V, \\ R &= (A'E + BF' + DG')U, & T &= (C'E - DF + BG)V, \end{aligned}$$

where $P = P(z)$, for $P = A, B, C, D$, and $E = z^{(s+1)t-1}$,

$$F = \sum_{k=-r}^{r-1} f_{r+k+1} z^{(2k+1)t+M}, \quad G = \sum_{k=-r}^{r-1} g_{r+k+1} z^{(2k+1)t};$$

$M = m + p = t - m$; $U = z^{(s-1)t+M}$, and $V = z^{st}$.

PROOF. By applying Theorem 1 to (Q, R, S, T) and observing that $UU' = 1 = VV'$, $|F|^2 + |G|^2 = 2s$, $|E|^2 = 1$ and $H = 0$, we obtain, from (2),

$$\begin{aligned} |Q|^2 + |R|^2 + |S|^2 + |T|^2 &= (|A|^2 + |B|^2 + |C|^2 + |D|^2)(|E|^2 + |F|^2 + |G|^2) \\ &= 2t(2s + 1), \end{aligned}$$

for any z in $\mathbf{K}$. Therefore, (4) is satisfied. And since

$$F' = \sum_{k=-r}^{r-1} f_{r+k+1} z^{-(2k+1)t-M} = \sum_{h=-r}^{r-1} f_{r-h} z^{(2h+1)t-M}$$

and

$$G' = \sum_{k=-r}^{r-1} g_{r+k+1} z^{-(2k+1)t} = \sum_{h=-r}^{r-1} g_{r-h} z^{(2h+1)t},$$

we have from (*)

$$\begin{aligned} (**) \quad Q &= \sum_{h=-r}^{r-1} (Af_{r-h} z^{(2h+1)t} + Cg_{r+h+1} z^{(2h+1)t+M}) z^{(s-1)t} - B' z^{2st+M-1}, \\ R &= \sum_{h=-r}^{r-1} (Bf_{r-h} z^{(2h+1)t} + Dg_{r-h} z^{(2h+1)t+M}) z^{(s-1)t} + A' z^{2st+M-1}, \\ S &= \sum_{h=-r}^{r-1} (Ag_{r-h} z^{(2h+1)t} - Cf_{r+h+1} z^{(2h+1)t+M}) z^{st} - D' z^{(2s+1)t-1}, \\ T &= \sum_{h=-r}^{r-1} (Bg_{r+h+1} z^{(2h+1)t} - Df_{r+h+1} z^{(2h+1)t+M}) z^{st} + C' z^{(2s+1)t-1}. \end{aligned}$$

Consequently, (5) is also satisfied. When $s = 1$, Theorem 2 takes the following form: i.e. $r = \frac{1}{2}$, thus $F = z^M$ and $G = 1$, if $f_1 = g_1 = 1$.

COROLLARY. When $s = 1$, we let $F = z^M$, $G = 1$, and E, U and V be as in Theorem 2. Then (Q, R, S, T) is $RD(3t)$.

For example, from the corollary, we obtain $RD(3t)$ as follows:

$$\begin{aligned} Q &= (A, C; 0, 0; -B', 0), & S &= (0, 0; A, -C; 0, -D'), \\ R &= (B, D; 0, 0; A', 0), & T &= (0, 0; B, -D; 0, C'). \end{aligned}$$

We note here that the above (Q, R, S, T) corresponds to $(f, g^*, e, -h^*)$ in the proof of the theorem of [1]. By setting $\mathbf{F} = (-1, 1)$ and $\mathbf{G} = (1, 1)$ as $GCL(2)$, we obtain the following $RD(5t)$ from (**).

$$\begin{aligned} Q &= (A, C; 0, 0; -A, C; 0, 0; -B', 0), & S &= (0, 0; A, C; 0, 0; A, -C; 0, -D'), \\ R &= (B, D; 0, 0; -B, D; 0, 0; A', 0), & T &= (0, 0; B, D; 0, 0; B, -D; 0, C'). \end{aligned}$$

In general, we obtain the following $RD((2s + 1)t)$ from Theorem 2, (**).

$$\begin{aligned} & (Af_s, Cg_1; 0, 0; AF_{s-1}, Cg_2; 0, 0; \dots; Af_1, Cg_s; 0, 0; -B', 0), \\ & (Bf_s, Dg_s; 0, 0; Bf_{s-1}, Dg_{s-1}; 0, 0; \dots; Bf_1, Dg_1; 0, 0; A', 0), \\ & (0, 0; Ag_s, -Cf_1; 0, 0; Ag_{s-1}, -Cf_2; 0, 0; \dots; 0, 0; Ag_1, -Cf_s; 0, -D'), \\ & (0, 0; Bg_1, -Df_1; 0, 0; Bg_2, -Df_2; 0, 0; \dots; 0, 0; Bg_s, -Df_s; 0, C'). \end{aligned}$$

REFERENCES

1. C. H. Yang, *Hadamard matrices and δ -codes of length $3n$* , Proc. Amer. Math. Soc. **85** (1982), 480–482.
2. ———, *Hadamard matrices, finite sequences, and polynomials defined on the unit circle*, Math. Comp. **33** (1979), 688–693.
3. R. J. Turyn, *Hadamard matrices, Baumert-Hall units, four symbol sequences, pulse compression, and surface wave encodings*, J. Combin. Theory Ser. A **16** (1974), 313–333.
4. ———, *Computation of certain Hadamard matrices*, Notices Amer. Math. Soc. **20** (1973), A–1.
5. A. C. Mukhopadhyay, *Some infinite classes of Hadamard matrices*, J. Combin. Theory Ser. A **25** (1978), 128–141.
6. A. V. Geramita and J. Seberry, *Orthogonal designs*, Dekker, New York, 1979.

DEPARTMENT OF MATHEMATICAL SCIENCES, STATE UNIVERSITY OF NEW YORK, ONEONTA, NEW YORK
13820