EXTENDING FAMILIES OF DISJOINT ZERO SETS

C. E. AULL

ABSTRACT. The z -cellularity $z(X)$ of a space X is defined as

$$z(X) = \sup\{|Z| : Z \subset Z(X)\}$$

where $Z(X)$ is the family of zero sets of X . It is proved using CH that a Tychonoff space S is TC^* -embedded in every Tychonoff space it is C -embedded in iff $z(S) \leq c$. A space S is defined to be TC^* -embedded in a space X if any disjoint family of zero sets of S can be extended to a family of disjoint zero sets of X . Similar theorems are proved for C^* -embedding when S is a P -space or the zero sets have the Isiwata property.

1. Introduction. If a set S is dense and C -embedded in a Tychonoff space X the closure of a zero set of S in X is a zero set of X [6]. Hence any family of disjoint zero sets of S may be extended to a family of disjoint zero sets of X . A set A of S is said to be extended to a set $E(A)$ of X if $E(A) \cap S = A$.

In this paper, we investigate the disjoint extension of families of disjoint zero sets of a set to a space, some families with restricted cardinals and some with restrictions to regular zero sets. Unless noted, definitions and terminology will be found in [6].

THEOREM 1. *A space S is C^* -embedded in a space X iff any denumerable family of disjoint zero sets may be extended disjointly to a family of zero sets of X .*

PROOF. Let $\{Z_n\}$ be a disjoint denumerable family of zero sets of X . Let H_{nk} be a zero set extension of Z_n to X disjoint from H_{kn} , $k \neq n$, guaranteed by the C^* -embedding of S in X . Then the family $\{H_n : H_n = \bigcap \{H_{nk} : k \neq n\}\}$ is the desired family of disjoint extensions of $\{Z_n\}$.

DEFINITION 1. A set S is TC^* -embedded (T_z -embedded) in a space X if any disjoint family of zero (cozero) sets of S can be extended to a disjoint family of zero (cozero) sets of X .

In [1], T_z -embeddings were studied.

It is clear that any C^* -embedded zero set is TC^* -embedded and if every subset of a space is TC^* -embedded, then the space is perfectly normal and extremally disconnected.

2. C -embeddings and TC^* -embeddings. We can improve Theorem 1 with respect to TC^* -embeddings when the embedded set is C -embedded and dense as mentioned earlier. More generally we have the following theorem.

THEOREM 2. *Let S be C -embedded in a Tychonoff space X . Then ω_1 disjoint zero sets in S may be extended to disjoint zero sets of X .*

Received by the editors August 9, 1982 and, in revised form, February 7, 1983.

1980 *Mathematics Subject Classification*. Primary 54C50, 54C45; Secondary 54D60, 54G05, 54G10, 54G20.

© 1983 American Mathematical Society
0002-9939/83 \$1.00 + \$.25 per page

PROOF. We order the ω_1 zero sets $\{Z_\alpha\}$ by putting them into one-to-one correspondence with the countable ordinals. Let $E(Z_\alpha)$ be a zero set extension of Z_α for $\alpha < \omega_1$. By means of Lemma 2, we construct a zero set Z_α^β of X such that $Z_\alpha \subset Z_\alpha^\beta$ and $Z_\alpha^\beta \cap E(Z_\beta) = \emptyset$ for each $\beta < \alpha$. The family $\{H_\alpha\}$, $H_\alpha = \bigcap_{\beta} \{Z_\alpha^\beta : \beta < \alpha\} \cap E(Z_\alpha)$ is a family of disjoint zero set extensions of $\{Z_\alpha\}$.

LEMMA 2. *A z -embedded set S is C -embedded in a space X iff for H a zero set of S and Z a zero set of X , $H \cap Z = \emptyset$, H and Z are completely separated in X .*

PROOF. *Sufficiency.* Since S is z -embedded and completely separated from any disjoint zero set of X , S is C -embedded in X [3].

Necessity. Let $f \in C(X)$ and $h \in C(S)$ such that $Z = f^{-1}(0)$ and $H = h^{-1}(0)$. Then $|f| + |h|$ is continuous on S and can be extended to a positive continuous function g on X . Set $h' = g - |f|$. Then $h' = |h|$ on S . Since $g > 0$, $h' > 0$ on Z . So $h^{-1}(0)$ and Z are disjoint zero sets of X containing H and Z respectively.

REMARK 1. It is clear from the above proof that C -embedding of a set $S \subset X$ is equivalent to C^* -embedding and every positive $f \in C^*(X)$ having a positive continuous extension. This is probably known.

DEFINITION 2. (z -cellularity) $z(X) = \sup\{|Z| : Z \subset Z(X), Z \text{ a disjoint family}\}$ where $Z(X)$ is the family of all zero sets of X .

COROLLARY 2. *If X is almost compact [6] with $z(X) = \omega_1$, then X is TC^* -embedded in every Tychonoff space X is embedded in.*

The necessity of the above conditions follows from the next theorem.

THEOREM 3 (CH). *A Tychonoff space X is TC^* -embedded in every Tychonoff space it is C -embedded (C^* -embedded) in iff (only if) $z(X) \leq c$.*

PROOF. Sufficiency of the C -embedding result follows from Theorem 2. Necessity is obtained by embedding νX in the product of real lines, in which X will be C -embedded in and by Engelking's [4, p. 295] result that $z(X) \leq c$ for the product of regular separable spaces. Thus if $z(X) > c$, there is a Tychonoff space in which X is C -embedded in but not TC^* -embedded in. The C^* -embedding result follows.

REMARK 2. We cannot replace C -embedding by C^* -embedding in the sufficiency argument as Ralph Fox has shown that R is not TC^* -embedded in βR .

COROLLARY 3 (CH). *A Tychonoff space X is TC^* -embedded in every Tychonoff space it is embedded in iff X is almost compact and $z(X) \leq c$.*

2. C^* -embeddings and TC^* -embeddings.

THEOREM 4. *If a family of ω_1 disjoint zero sets of a Tychonoff space S may be extended disjointly to βS , then this family of zero sets of S may be disjointly extended to any Tychonoff space S is C^* -embedded in.*

PROOF. If S is C^* -embedded in X , $\beta S \subset \beta X$ and βS is C -embedded in βX . By Theorem 2 ω_1 disjoint zero sets of βS may be extended to βX . The result follows.

COROLLARY 4 (CH). *Let S be Tychonoff and TC^* -embedded in βS . Then S is TC^* -embedded in every space it is C^* -embedded in iff $z(S) \leq c$.*

It is not unusual to have a set S TC^* -embedded in βS even when there are families of disjoint zero sets of S of large cardinality. This is always true when S is pseudocompact or discrete. In these cases, by Theorem 4, any family of ω_1 disjoint zero sets may be extended disjointly to a family of zero sets of any Tychonoff space in which S is C^* -embedded in. The next two theorems are modifications of these results.

THEOREM 5. *Let $\{z_\alpha\}$ be a family of disjoint zero sets of cardinality ω_1 of a set S such that for $f \in C^*(S)$ such that $Z(f) \cap Z_\alpha = \emptyset$, $|f(z)| \geq \epsilon > 0$ for $z \in Z_\alpha$. Then $\{Z_\alpha\}$ may be extended disjointly to any space in which S is C^* -embedded.*

PROOF. Let $E(Z_\alpha)$ be a zero set extension of Z_α to X . Thus as in Lemma 2, $E(Z_\alpha)$ is completely separated in X from Z_β for $\beta \neq \alpha$. The proof is completed as in Theorem 2.

The property referred to at the end of the first sentence of the statement of the theorem was discovered by Isiwata [7] to be satisfied by all zero sets of a pseudocompact space. We will refer to the above property as the Isiwata property. By modifying the proof of Theorem 5, one may prove the following result.

THEOREM 5A. *let $S \subset X$, X Tychonoff, have at most ω_1 disjoint zero sets and let every disjoint family of zero sets, none of which has the Isiwata property be denumerable. Then S is TC^* -embedded in X if S is C^* -embedded in X .*

DEFINITION 3 (COMPARE [2]). A weak 0_z -space is a space where the closure of a cozero set is a zero set.

We will call a zero set that is also a regular closed set a regular zero set.

THEOREM 6. *Let βS be weak 0_z . If $\{Z_\alpha\}$ is a disjoint family of regular zero sets (of cardinality ω_1) then $\{Z_\alpha\}$ may be extended disjointly to a family of zero sets of βS (of X Tychonoff where S is C^* -embedded in X).*

PROOF. Since βS is weak 0_z , S is weak 0_z . So the interior of Z_α is a cozero set. From [1], $\{\text{int } Z_\alpha\}$ has a family of disjoint cozero extensions $\{C_\alpha\}$ in βS . It is immediate that in βS , $\overline{C_\alpha} = \overline{Z_\alpha}$ so that $\overline{Z_\alpha}$ is a zero set of βS . By the C^* -embedding of S in βS , these zero sets are necessarily disjoint. An application of Theorem 4 completes the proof.

An important class of spaces that satisfy the conditions of the theorem are basically disconnected spaces.

COROLLARY 6. *Let S be a P -space. Then S is TC^* -embedded in βS and any family of ω_1 disjoint zero sets of S may be extended disjointly to any Tychonoff space S is C^* -embedded in.*

PROOF. Since S is a P -space, βS is basically disconnected and each zero set of S is a regular closed set.

4. Some examples. The author is indebted to E. van Douwen for pointing out the following theorem of Juhasz which will be useful in analyzing examples.

THEOREM A (JUHASZ [8]). *If X is compact (even if X is the union of G_δ -sets of some compactification, i.e. X is of point-countable type) and S has cellularity m , m infinite, then X has at most 2^m disjoint G_δ -sets.*

In the above theorem, we can replace G_δ -sets by zero sets and compact by pseudocompact sets based on Theorem 2.

We have already noted R. Fox's result that R is not TC^* -embedded in βR . However as all disjoint families of regular zero sets are countable, these families are disjointly extendable. By Theorem 5, using CH, all disjoint families of compact zero sets of R are disjointly extendable.

The following example of R. Pol, pointed out to the author by R. Hodel, shows that even in the case of dense C^* -embedding we may be limited in the cardinality of the number of disjoint zero sets that are extended in case S is not pseudocompact or a P -space.

EXAMPLE 1 (R. POL [9]). The space S is Tychonoff, satisfies c.c.c., has countable pseudocharacter and has cardinality 2^c . Thus S has 2^c disjoint zero sets. Furthermore βS satisfies c.c.c.; so by Theorem A the z -cellularity of βS is at most c . So S is not TC^* -embedded in βS .

EXAMPLE 2. Let X be the absolute of S ; then βX is the absolute of βS [11] and since c.c.c. is preserved under taking absolutes [11], X satisfies c.c.c. and since there is a continuous map from X to S , $z(x) \geq 2^c$. By Theorem A and the fact that βX satisfies c.c.c., the z -cellularity of βX is at most c . Thus X is not TC^* -embedded in βX and X is extremally disconnected.

EXAMPLE 3. Since $\beta N \sim N$ is a zero set of βN , $\beta N \sim N$ is TC^* -embedded in βN and is not T_z -embedded in βN . Even though C^* -embedding implies z -embedding TC^* -embedding does not imply T_z -embedding.

Blair [2] has shown that if a weakly perfectly normal space is not realcompact one can add a point of $\nu X \sim X$ and obtain a weakly perfectly normal space that is not perfectly normal (the additional point is not a G_δ in the space). We will call this space $W(X)$. It is clear that if X is normal and hereditarily extremally disconnected then so will $W(X)$ be.

EXAMPLE 4 (BLAIR [2]). If D is a discrete space of measurable cardinality $W(D)$ is a hereditarily extremally disconnected normal space that is not perfectly normal. Furthermore it is easily shown that every subset of $W(D)$ is TC^* -embedded so that every subset of a space may be TC^* -embedded without the space being perfectly normal.

Below we have another example of this where we use the set-theoretic axiom club.

EXAMPLE 5 (WAGE [10]). The space X is perfectly normal and extremally disconnected and constructed using club. Then $W(X)$ is normal and hereditarily extremally disconnected and is not perfectly normal.

The question then arises as to whether normal hereditarily extremally disconnected spaces have the property that every subset is TC^* -embedded, particularly under $MA + \sim CH$.

Finally, the author would like to thank the referee for his many helpful suggestions.

REFERENCES

1. C. E. Aull, *Extendability and expandability* (submitted for publication).
2. R. L. Blair, *Spaces in which special sets are z -embedded*, *Canad. J. Math.* **28** (1976), 673–690.
3. R. L. Blair and A. W. Hager, *Extensions of zero-sets and of real-valued functions*, *Math. Z.* **136** (1974), 41–52.

4. R. Engelking, *Cartesian products and dyadic spaces*, Fund. Math. **57** (1965), 287–306.
5. ———, *General topology*, PWN, Warsaw, 1977.
6. L. Gillman and M. Jerison, *Rings of continuous functions*, Van Nostrand, Princeton, N. J., 1966.
7. T. Isiwata, *Mappings and spaces*, Pacific J. Math. **20** (1967), 455–480.
8. I. Juhász, *Cardinal functions in topology*, Math. Centre Tracts, No. 34, Mathematisch Centrum, Amsterdam, 1971.
9. R. Pol, *Short proofs of two theorems of cardinality of topological spaces*, Bull. Acad. Polon. Sci. Sér. Math. **22** (1974), 1245–1249.
10. M. Wage, *Extremally disconnected S -spaces*, Topology Proc. (Conf. Auburn Univ. Auburn, Ala, 1976), Math. Dept., Auburn Univ., Auburn, 1977, pp. 181–186.
11. R. G. Woods, *Absolutes of topological spaces*, Math. Centre Tracts, No. 116, Mathematisch Centrum, Amsterdam, 1979, pp. 323–362.

DEPARTMENT OF MATHEMATICS, VIRGINIA POLYTECHNIC INSTITUTE AND STATE UNIVERSITY, BLACKSBURG, VIRGINIA 24061