PRESENTATION OF ASSOCIATED GRADED RINGS OF COHEN-MACAULAY LOCAL RINGS

YOUNG-HYUN CHO¹

ABSTRACT. Let $(R, \mathfrak{m})$ be a local ring and I be an $\mathfrak{m}$ -primary ideal such that $\dim_k(I/I\mathfrak{m}) = l$, where $k = R/\mathfrak{m}$. Denote the associated graded ring with respect to I , $\bigoplus_{n=0}^{\infty} I^n/I^{n+1}$, by $G_I(R)$. Then $G_I(R) \cong R/I[X_1, \dots, X_l]/\mathcal{L}$, for some homogeneous ideal $\mathcal{L}$. Set $M = \max \deg_{1 \leq i \leq l} f_i$, where $\{f_1, \dots, f_l\}$ is a set of homogeneous elements which form a minimal basis of $\mathcal{L}$. The main result in this note is that if R is a Cohen-Macaulay local ring of dimension 1 and if $G_I(R)$ is free over R/I , then $M \leq r(I) + 1$, where $r(I)$ is the reduction number of I . It follows that $M \leq e(R)$ where $e(R)$ is the multiplicity of R .

1. Introduction. First of all, we refer the reader to [5] for notations. Let $(R, \mathfrak{m})$ be a local ring and I be an $\mathfrak{m}$ -primary ideal such that $\dim_k(I/I\mathfrak{m}) = l$, where $k = R/\mathfrak{m}$. Choose a minimal basis of I , $x_1, \dots, x_l$, and give a map $\phi: R/I[X_1, \dots, X_l] \rightarrow G_I(R)$ defined by $\phi(X_i) = \bar{x}_i$, for all $i = 1, \dots, l$. Here, $\bar{x}_i$ denotes the image of x_i in I/I^2 . Then we can see that $G_I(R)$ is the homomorphic image of $R/I[X_1, \dots, X_l]$, i.e., $G_I(R) \cong R/I[X_1, \dots, X_l]/\mathcal{L}$, for some homogeneous ideal $\mathcal{L}$. When $G_I(R)$ is free over R/I , $\mathcal{L}$ does not depend on the choice of the basis of I , because if we have another minimal basis of I , $y_1, \dots, y_l$, and give a map $\phi': R/I[X_1, \dots, X_l] \rightarrow G_I(R)$ defined by $\phi'(X_i) = \bar{y}_i$, for all $i = 1, \dots, l$, then we get the following commutative diagram

$$\begin{array}{ccccccc}
 0 & \rightarrow & \text{Ker } \phi & \rightarrow & R/I[X_1, \dots, X_l] & \xrightarrow{\phi} & G_I(R) \rightarrow 0 \\
 & & \downarrow i & & \downarrow = & & \downarrow \psi \\
 0 & \rightarrow & \text{Ker } \phi' & \rightarrow & R/I[X_1, \dots, X_l] & \xrightarrow{\phi'} & G_I(R) \rightarrow 0
 \end{array}$$

$\psi(\bar{x}_i) = \bar{y}_i, \text{ for } i = 1, \dots, l,$

and this induces an isomorphism $i: \text{Ker } \phi \xrightarrow{\cong} \text{Ker } \phi'$. Let $\{f_1, \dots, f_l\}$ be a set of homogeneous elements which form a minimal basis of $\mathcal{L}$. We will study the upper bound $\max \deg_{1 \leq i \leq l} f_i = M$.

REMARK 1. We can see that the number of elements and the set of degrees of elements in the minimal basis of $\mathcal{L}$ are invariant of $\mathcal{L}$. This fact comes from [1, III-§1, Definition 1 and Lemma 1] with a slight modification.

Received by the editors September 15, 1982.

1980 *Mathematics Subject Classification*. Primary 13H10, 13H15.

¹The author was partially supported by the Korea Science and Engineering Foundation research grant.

©1983 American Mathematical Society
 0002-9939/83 \$1.00 + \$.25 per page

2. The case when $\dim R = 1$. We consider the case when R is a Cohen-Macaulay local ring of dimension 1. Before we prove the main theorem, we will mention the flat change of local rings $R \rightarrow R(u) = R[u]_{\mathfrak{m}R[u]}$, where u is an indeterminate.

LEMMA 2. *Let $(R, \mathfrak{m})$ be a local ring. Consider $R \rightarrow R(u)$. Then*

- (a) $R(u)$ is faithfully flat over R .
- (a') $IR(u) \cap R = I$, for any ideal I in R .
- (b) $\dim R = \dim R(u)$.
- (c) $\text{depth } R = \text{depth } R(u)$.
- (d) $v(I) = v(IR(u))$, for any ideal I in R .
- (e) $\lambda(R/I) = \lambda(R(u)/IR(u))$, for any $\mathfrak{m}$ -primary ideal I .
- (f) $e(I) = e(IR(u))$, for any $\mathfrak{m}$ -primary ideal I .

PROOF. (a) and (c) See [3, p. 27, Corollary and (21c)].

(b) By [3, Theorem 19(2)], $\text{ht}(\mathfrak{m}R(u)) = \text{ht}(\mathfrak{m}R(u) \cap R) = \text{ht}(\mathfrak{m})$.

(d) $v(I) = \dim_k(I/I\mathfrak{m})$, where

$$k = R/\mathfrak{m} = \dim_{k(u)}(I/I\mathfrak{m} \otimes R(u)) = \dim_{k(u)}(IR(u)/I\mathfrak{m}R(u)) = v(IR(u)).$$

(e) If $0 = I/I \subset I_1/I \subset \cdots \subset R/I$ is a composition series of R/I , then $0 = IR(u)/IR(u) \subset I_1R(u)/IR(u) \subset \cdots \subset R(u)/IR(u)$ is also.

(f) From (e), $\lambda(I^n/I^{n+1}) = \lambda(I^nR(u)/I^{n+1}R(u))$, for any $n > 0$. Thus $e(I) = e(IR(u))$ follows.

LEMMA 3. *For any ideal I in a local ring $(R, \mathfrak{m})$,*

- (a) $\dim G_I(R) = \dim G_{IR(u)}(R(u))$.
- (b) $\text{depth } G_I(R) = \text{depth } G_{IR(u)}(R(u))$. Here, $\text{depth } G_I(R)$ (or $\text{depth } G_{IR(u)}(R(u))$) means a maximal $G_I(R)$ -sequence (or $G_{IR(u)}(R(u))$ -sequence) in $\mathfrak{m}/I \oplus I/I^2 \oplus \cdots$ (or $\mathfrak{m}R(u)/IR(u) \oplus IR(u)/I^2R(u) \oplus \cdots$).

PROOF. (a) Consider the Rees ring $\mathfrak{R} = R[[t]]$ of R with respect to I . Then t^{-1} is not a zero divisor in $\mathfrak{R}$ and $\dim \mathfrak{R} = \dim R + 1$. (See [4, Remark 3.7].) Since $G_I(R) = \mathfrak{R}/t^{-1}\mathfrak{R}$, $\dim G_I(R) = \dim R$. By the same way, $\dim G_{IR(u)}(R(u)) = \dim R(u)$. By Lemma 2(b), the assertion follows.

(b) Put $\mathfrak{N} = \mathfrak{m}/I \oplus I/I^2 \oplus \cdots$ and $\mathfrak{N}(u) = \mathfrak{m}R(u)/IR(u) \oplus IR(u)/I^2R(u) \oplus \cdots$. We only need to show that $\text{depth } G_I(R)_{\mathfrak{N}} = \text{depth } G_{IR(u)}(R(u))_{\mathfrak{N}(u)}$. Since $R(u)$ is faithfully flat over R , by [3, (4a)], $R(u) \otimes_R G_I(R) = G_{IR(u)}(R(u))$ is faithfully flat over $G_I(R)$. Then by [3, p. 154, Corollary],

$$\text{depth } G_I(R)_{\mathfrak{N}} = \text{depth } G_{IR(u)}(R(u))_{\mathfrak{N}(u)}.$$

Now we are going to prove the main theorem. The reduction number of I , denoted by $r(I)$, means the least integer r such that $xI^r = I^{r+1}$ for some $x \in I \setminus I^2$.

THEOREM 1. *Let $(R, \mathfrak{m})$ be a 1-dimensional Cohen-Macaulay local ring and I be an $\mathfrak{m}$ -primary ideal such that $G_I(R)$ is free over R/I . Then $M \leq r(I) + 1$.*

PROOF. Set $\dim_k(I/I\mathfrak{m}) = l$, where $R/\mathfrak{m} = k$, and let ϕ map $R/I[X_1, \dots, X_l]$ onto $G_I(R)$ by $\phi(X_i) = \bar{x}_i$, where $x_1, \dots, x_l$ is a minimal basis of I . Let $\mathfrak{L} = \text{Kernel } \phi$

and let $f_1, \dots, f_t$ be any minimal basis of $\mathcal{L}$ consisting of homogeneous elements. Then

$$G_{IR(u)}(R(u)) \simeq R(u)/IR(u)[X_1, \dots, X_t]/(f_1, \dots, f_t) \otimes R(u)$$

and $f_1, \dots, f_t$ is a minimal set of generators for $(\text{Kernel } \phi) \otimes R(u)$. The degree of f_i in $R/I[X_1, \dots, X_t]$ is same as the degree of f_i in $R(u)/IR(u)[X_1, \dots, X_t]$ by Lemma 2. Hence, by Remark 1 and Lemmas 2 and 3 we can show that theorem just in the case of the ring $R(u)$.

Therefore we can assume $R/\mathfrak{m}$ is infinite and we can pick up x_1 which is a superficial element of I such that $x_1 I^r = I^{r+1}$, where $r = r(I)$. (See [6, Lemma 5, p. 286] for the existence of x_1 .) We can check that $(0: \bar{x}_1 G_I(R)) \cap I^n/I^{n+1} = 0$ for all $n \geq r-1$. Choose $x_2, \dots, x_t$ such that $\{\bar{x}_i\}_{i=1, \dots, t}$ is a minimal basis for I/I^2 , and define $\phi: R/I[X_1, \dots, X_t] \rightarrow G_I(R)$ by $\phi(X_i) = \bar{x}_i$ as before. Let $\mathcal{L} = \text{Kernel } \phi$. Then $G_I(R) \simeq R/I[X_1, \dots, X_t]/\mathcal{L}$. Since $x_1 I^r = I^{r+1}$,

$$\bar{x}_1(I/I^2 \oplus I^2/I^3 \oplus \dots)^r = (I/I^2 \oplus I^2/I^3 \oplus \dots)^{r+1},$$

i.e., $X_1(X_1, \dots, X_t)^r + \mathcal{L}/\mathcal{L} = (X_1, \dots, X_t)^{r+1} + \mathcal{L}/\mathcal{L}$. Hence $(X_1, \dots, X_t)^{r+1} + \mathcal{L} = X_1(X_1, \dots, X_t)^r + \mathcal{L}$, therefore $(X_1, \dots, X_t)^{r+1} \subset X_1(X_1, \dots, X_t)^r + \mathcal{L}$. For convenience, let P denote $(X_1, \dots, X_t)$ in following Claims 1 and 2.

Claim 1. $\mathcal{L} \cap X_1^{j+1}P^r \subset P^j\mathcal{L}$, for any $j \geq 0$.

PROOF. We will prove by induction on j . When $j = 0$, $\mathcal{L} \cap X_1 P^r \subset \mathcal{L}$ is obviously true. Suppose $j > 0$ and the claim is true for $j-1$. Let $X_1^{j+1}g \in \mathcal{L}$, where $g \in P^r$. We write $g = \sum_{i=r}^p g_i$ with g_i homogeneous and degree of $g_i = i$, for any $i = r, \dots, p$. Since $\sum_{i=r}^p X_1^{j+1}g_i \in \mathcal{L}$ and $\mathcal{L}$ is a homogeneous ideal, $X_1^{j+1}g_i \in \mathcal{L}$ for $i = r, \dots, p$. Therefore, $(X_1 + \mathcal{L})(X_1^j g_i + \mathcal{L}) = 0$ in $R/I[X_1, \dots, X_t]/\mathcal{L}$. We know that $X_1 + \mathcal{L}$ corresponds to $\bar{x}_1$ and $X_1^j g_i + \mathcal{L}$ corresponds to some elements in I^n/I^{n+1} , where $n \geq r+j$. Hence $X_1^j g_i + \mathcal{L} = 0$ since $(0: \bar{x}_1 G_I(R)) \cap I^n/I^{n+1} = 0$ for all $n \geq r-1$. This means $X_1^j g_i \in \mathcal{L}$ for $i = r, \dots, p$. By induction hypothesis, $X_1^j g_i \in P^{j-1}\mathcal{L}$, for $i = r, \dots, p$. Now, $X_1^{j+1}g = X_1 \sum_{i=r}^p X_1^j g_i \in P^j\mathcal{L}$. So we get $\mathcal{L} \cap X_1^{j+1}P^r \subset P^j\mathcal{L}$.

Claim 2. $\mathcal{L} \cap P^{r+1+j} \subset P^j\mathcal{L}$, for any $j \geq 0$.

PROOF. Before Claim 1, we had $P^{r+1} \subset X_1 P^r + \mathcal{L}$. Now

$$\begin{aligned} P^{r+2} &= P \cdot P^{r+1} \subset P(X_1 P^r + \mathcal{L}) \\ &= X_1 P^{r+1} + P\mathcal{L} \subset X_1(X_1 P^r + \mathcal{L}) + P\mathcal{L} = X_1^2 P^r + P\mathcal{L}. \end{aligned}$$

By the same way we get $P^{r+1+j} \subset X_1^{j+1}P^r + P^j\mathcal{L}$, for any $j \geq 0$. Next,

$$P^{r+1+j} \cap \mathcal{L} \subset (X_1^{j+1}P^r + P^j\mathcal{L}) \cap \mathcal{L} = X_1^{j+1}P^r \cap \mathcal{L} + P^j\mathcal{L} \subset P^j\mathcal{L},$$

since Claim 1 holds.

To finish the proof, let $\{f_1, \dots, f_t\}$ be a set of homogeneous elements which form a minimal basis of $\mathcal{L}$. Assume some f_i 's have degree greater than $r+1$, say $\deg f_1 = r+1+j$, for some $j > 0$. Then $f_1 \in \mathcal{L} \cap (X_1, \dots, X_t)^{r+1+j} \subset (X_1, \dots, X_t)^j\mathcal{L}$. Therefore $f_1 = \sum_{i=1}^n h_i g_i$ with $h_i \in (X_1, \dots, X_t)^j$, $g_i \in \mathcal{L}$ and h_i, g_i are homogeneous elements for any i . Since f_1 is a homogeneous element, we can assume $\deg f_1 = \deg h_i g_i$ for any i . Now $\deg h_i g_i = r+1+j$ and $\deg h_i \geq j$ imply $\deg g_i \leq r+1$ so that $g_i \in \mathcal{L}$ is in $(f_2, \dots, f_t)$. So, $f_1 \in (f_2, \dots, f_t)$. A contradiction follows from the minimality of the f_i 's.

COROLLARY 1. *Let $R, \mathfrak{m}, I$ be the same as in Theorem 1. Then $M \leq e(R)$.*

PROOF. If R is a discrete valuation ring, $G_I(R) = R/I[X]$, since $I = (x)$ for some regular element x . Otherwise $r(I) \leq e(R) - 1$ (cf. Remark 1.5 and Corollary 2.2 of [2]). Hence $M \leq r(I) + 1 \leq e(R)$. Of course the hypothesis that $G_I(R)$ is free over R/I is satisfied by $I = \mathfrak{m}$.

3. The general case. For the proof of Theorem 2, we need the following lemma and corollary.

LEMMA 4. *Let A be any ring and $\mathfrak{L}$ any homogeneous ideal in $A[X_1, \dots, X_n]$. Then $f_1, \dots, f_l$ generate $\mathfrak{L}$ if and only if $f_1 + \mathfrak{J}\mathfrak{L}, \dots, f_l + \mathfrak{J}\mathfrak{L}$ generate $\mathfrak{L}/\mathfrak{J}\mathfrak{L}$ over $A[X_1, \dots, X_n]/\mathfrak{L}$, where $\mathfrak{J}$ is a homogeneous ideal contained in $(X_1, \dots, X_n)$.*

PROOF. ($\rightarrow$) It is obvious.

($\leftarrow$) Suppose $f_1, \dots, f_l$ are elements in $\mathfrak{L}$ such that $(f_1, \dots, f_l) + \mathfrak{J}\mathfrak{L} = \mathfrak{L}$. Then we get $(f_1, \dots, f_l) + \mathfrak{J}^n\mathfrak{L} = \mathfrak{L}$ for any n . If we choose any homogeneous element g in $\mathfrak{L}$ with $\deg g = m$, then $g \in (f_1, \dots, f_l) + \mathfrak{J}^{m+1}\mathfrak{L}$. By comparing degrees, g must belong to $(f_1, \dots, f_l)$. Therefore $\mathfrak{L} = (f_1, \dots, f_l)$.

COROLLARY 2. *Let $\mathfrak{L}$ be a homogeneous ideal in $A[X_1, \dots, X_n]$ and $X_1 + \mathfrak{L}$ is a regular element in $A[X_1, \dots, X_n]/\mathfrak{L}$. Then the number of minimal generators of $\mathfrak{L} =$ number of minimal generators of $(\mathfrak{L}, X_1)/(X_1)$.*

PROOF. ($\geq$) It is obvious.

($\leq$) Suppose $\{f_1, \dots, f_t\}$ is a subset of $\mathfrak{L}$ such that

$$(f_1, \dots, f_t) + (X_1)/(X_1) = \mathfrak{L} + (X_1)/(X_1) \simeq \mathfrak{L}/(X_1) \cap \mathfrak{L} = \mathfrak{L}/X_1\mathfrak{L},$$

since $X_1 + \mathfrak{L}$ is a regular element in $A[X_1, \dots, X_n]/\mathfrak{L}$. Now, images of f_i in $\mathfrak{L}/X_1\mathfrak{L}$ generate $\mathfrak{L}/X_1\mathfrak{L}$. By Lemma 4, $f_1, \dots, f_t$ generate $\mathfrak{L}$.

THEOREM 2. *Let $(R, \mathfrak{m})$ be a d -dimensional Cohen-Macaulay local ring and I be an $\mathfrak{m}$ -primary ideal such that $\text{depth } G_I(R) \geq d - 1$ and $G_I(R)$ is free over R/I . Then $M \leq e(R)$.*

PROOF. By Lemmas 2 and 3, we just need to prove the theorem in the ring $R(u)$. Therefore we can assume $R/\mathfrak{m}$ is infinite. We will prove the theorem by induction on d . When $d = 1$, we showed it in Corollary 1. Suppose $d > 1$. First, we can choose $x_1 \in I \setminus I^2$ such that $\bar{x}_1$ is a regular element in $G_I(R)$. (Sketch of proof: Let $\text{Ass } G_I(R) = \{p_1, \dots, p_t\}$. Then

$$p_i \cap I/I^2 \subsetneq I/I^2, \quad \text{for any } i = 1, \dots, t.$$

Since $R/\mathfrak{m}$ is infinite, $I/I^2 \supsetneq \bigcup_{i=1}^t (p_i \cap I/I^2)$. Choose $x_1 \in I \setminus I^2$ such that $\bar{x}_1 \in I/I^2 \setminus \bigcup_{i=1}^t (p_i \cap I/I^2)$. Note that such x_1 is a nonzero divisor in R . We can check that $G_{I/x_1R}(R/x_1R) \simeq G_I(R)/\bar{x}_1 G_I(R)$, and R/x_1R is a Cohen-Macaulay local ring of dimension $d - 1$ and $\text{depth } G_{I/x_1R}(R/x_1R) \geq d - 2$, and $\dim_k(I/I\mathfrak{m} + x_1R) = \dim_k(I/I\mathfrak{m}) - 1$, and $e(R/x_1R) = e(R)$. (See [5] for the reference.) Therefore we can assume that the theorem is true for R/x_1R by the

induction hypothesis. Next, choose $x_2, \dots, x_l$ ($l = \dim_k(I/I\mathfrak{m})$) such that $\{\bar{x}_i\}_{i=1, \dots, l}$ is a minimal basis for I/I^2 . Define $\phi: R/I[X_1, \dots, X_l] \rightarrow G_l(R)$ by $\phi(X_i) = \bar{x}_i$ for any $i = 1, \dots, l$. Let $\mathfrak{L} = \text{Kernel } \phi$ and $f_1, \dots, f_t$ be homogeneous elements which form a minimal basis of $\mathfrak{L}$. Then

$$\begin{aligned} G_{I/x_1 R}(R/x_1 R) &\simeq G_l(R)/\bar{x}_1 G_l(R) \simeq \frac{R/I[X_1, \dots, X_l]/\mathfrak{L}}{X_1(R/I[X_1, \dots, X_l]/\mathfrak{L})} \\ &\simeq \frac{R/I[X_1, \dots, X_l]}{(X_1) + \mathfrak{L}} \simeq \frac{R/I[X_2, \dots, X_l]}{(f_1, \dots, f_t) + (X_1)/(X_1)}. \end{aligned}$$

By assumption, there exists a minimal set $g_1, \dots, g_s$ such that g_i 's are homogeneous elements with respect to $X_2, \dots, X_l$ and $\deg g_i \leq e(R)$ for any i , and

$$(f_1, \dots, f_t) + (X_1)/(X_1) = (g_1, \dots, g_s) + (X_1)/(X_1).$$

We will show that $\max \deg_{1 \leq i \leq t} f_i \leq e(R)$. Suppose not, i.e. $\deg f_1 > e(R)$. Then $g_i \in (f_2, \dots, f_t) + (X_1)$ for any $i = 1, \dots, s$, since $\deg g_i \leq e(R)$. Therefore $(g_1, \dots, g_s) + (X_1) = (f_2, \dots, f_t) + (X_1)$ so that $(f_1, \dots, f_t) + (X_1)/(X_1) = (f_2, \dots, f_t) + (X_1)/(X_1)$. This contradicts that $\{\bar{f}_1, \dots, \bar{f}_t\}$ is a minimal basis of $\mathfrak{L} + (X_1)/(X_1)$, by Corollary 2.

COROLLARY 3. *Let $(R, \mathfrak{m})$ be a d -dimensional Cohen-Macaulay local ring such that $\text{depth } G_{\mathfrak{m}}(R) \geq d - 1$. Assume that $R = S/J$ with $(S, \mathfrak{m})$ an l -dimensional regular local ring and $J \subset \mathfrak{m}^2$. Then $G_{\mathfrak{m}}(R) \simeq k[X_1, \dots, X_l]/\mathfrak{L}$, $k = S/\mathfrak{m}$, where the degrees of elements in a homogeneous minimal basis of $\mathfrak{L}$ are at most $e(R)$.*

REMARK 2. In many cases $\max \deg_{1 \leq i \leq t} f_i$ may be strictly less than $r(I) + 1$. (Cf. Theorem 1.) Consider $R = k[[t^6 t^7 t^8]]$, k is any field. Then $e(R) = 6$, $r(\mathfrak{m}) = 3$, $G_{\mathfrak{m}}(R) \simeq k[X, Y, Z]/(Z^3, Y^2 - XZ)$. Hence the maximal degree of generators $< r(\mathfrak{m}) + 1$. But, consider $R = k[[t^6, t^7, t^{15}]]$. Then $e(R) = 6$, $r(\mathfrak{m}) = 5$, $G_{\mathfrak{m}}(R) \simeq k[X, Y, Z]/(XZ, Z^2, ZY^3, Y^6)$. Therefore, in this case the maximal degree of generators $= r(\mathfrak{m}) + 1$.

This note forms a part of my doctoral thesis. I gratefully acknowledge my appreciation to Professor Judith Sally.

REFERENCES

1. H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero*. II, Ann. of Math. (2) **79** (1964), 205–326.
2. J. Herzog and R. Waldi, *A note on the Hilbert function of a 1-dimensional Cohen-Macaulay ring*, Manuscripta Math. **16** (1975), 251–260.
3. H. Matsumura, *Commutative algebra*, 2nd ed., Benjamin Cummings, New York, 1980.
4. L. J. Ratliff, Jr., *On quasi-unmixed local domains, the altitude formula, and the chain condition for prime ideals*. II, Amer. J. Math. **92** (1970), 99–144.
5. J. D. Sally, *Numbers of generators of ideals in local rings*, Lecture Notes in Pure and Appl. Math., Dekker, New York and Basel, 1978.
6. O. Zariski and P. Samuel, *Commutative algebra*, Lecture Notes in Math., vol. 2, Springer-Verlag, Berlin and New York, 1975.

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL, KOREA