ON A FAMILY OF SPECIAL LINEAR SYSTEMS ON ALGEBRAIC CURVES¹

EDMOND E. GRIFFIN II

ABSTRACT. $\mathcal{W}_6^2$ is a scheme parametrizing pairs $L \rightarrow C$ of smooth algebraic curves C of genus 10 together with line bundles L of degree 6 such that $H^0(C, L) \geq 3$. It is shown that one of the irreducible components of this scheme is nonreduced at every point.

Introduction. In [ACGH] a general method is outlined for computing the dimension of the tangent space $T_l(\mathcal{W}_{d,g}^r)$. $\mathcal{W}_{d,g}^r$ is the scheme parametrizing pairs $L \rightarrow C$ in which C is a smooth curve of genus g , L is a line bundle such that

$$(1) \deg_C(L) = d,$$

$$(2) \dim H^0(C, L) = h^0(C, L) \geq r + 1,$$

and $l = L \rightarrow C$ is a "generic" point on $\mathcal{W}_{d,g}^r$. Specifically, $l \in \mathcal{W}_{d,g}^r - \mathcal{W}_{d,g}^{r+1}$.

The dimension of this tangent space is determined by the kernels of two naturally defined maps on the cohomology of C in L and K_C . These maps, μ_0 and μ_1 , together with their kernels will be computed explicitly in the case $l \in \mathcal{W}_{6,10}^2 - \mathcal{W}_{6,10}^3$ and C a hyper-elliptic curve. This calculation will then be used to show that $\mathcal{W}_{6,10}^2$ has a *nonreduced* component.

It will be clear that these computations generalize to higher genus and degree hyper-elliptic curves, but the actual calculations become fairly intractable. For this reason only the case of $\mathcal{W}_{6,10}^2$ will be considered here.

1. We begin with a review of the basic set up and terminology of [ACGH]. For a fixed, smooth curve C of genus g , define

$$\text{Pic}^d(C) = \{L \mid \deg_C(L) = d\}$$

$$\cup$$

$$W_d^r(C) = \{L \mid \deg_C(L) = d \text{ and } h^0(C, L) \geq r + 1\}.$$

It is well known that

$$T_L(\text{Pic}^d(C)) \simeq H^1(C, \mathcal{O}_C).$$

Received by the editors April 25, 1983.

1980 *Mathematics Subject Classification*. Primary 14H15, 14D15, 14H45.

¹This work forms part of the author's doctoral thesis, written at Harvard University in 1982 under the advice of Phillip Griffiths.

©1984 American Mathematical Society
0002-9939/84 \$1.00 + \$.25 per page

A first order variation of a line bundle L on C is a commuting diagram,

$$\begin{array}{ccc} L & \rightarrow & \mathbb{L} \\ \downarrow & & \downarrow \\ C & \rightarrow & C \times S_1 \\ \downarrow & & \downarrow \\ S_0 & \hookrightarrow & S_1 \end{array}$$

where $S_n = \text{Spec}(\mathbb{C}[t]/(t^{n+1}))$. The set of isomorphism classes of such diagrams can be canonically identified with $H^1(C, \mathcal{O}_C)$.

The obstruction to extending a section $s \in H^0(C, L)$ to a section of $\mathbb{L}$ can be seen to be $\varphi \cdot s \in H^1(C, L)$, where

$$“\cdot” H^1(C, \mathcal{O}_C) \otimes H^0(C, L) \rightarrow H^1(C, L)$$

is the usual cup product, and $\varphi \in H^1(C, \mathcal{O}_C)$ is the element corresponding to $\mathbb{L}$. Thus, one sees that, at a point $L \in W_d^r(C) - W_d^{r+1}(C)$, a tangent vector φ in $H^1(C, \mathcal{O}_C) = T_L(\text{Pic}^d(C))$ is also in $T_L(W_d^r(C))$ if and only if $\varphi: H^0(C, L) \rightarrow H^1(C, L)$ is the zero map. In other words, φ is in the kernel of the map

$$H^1(C, \mathcal{O}_C) \rightarrow \text{Hom}(H^0(C, L), H^1(C, L)).$$

Dualizing yields the map

$$\mu_0: H^0(C, L) \otimes H^0(C, KL^{-1}) \rightarrow H^0(C, K)$$

such that

$$T_L(W_d^r(C)) = (\text{Image } \mu_0)^\perp$$

in $h^0(C, L) = r + 1$.

Now the Riemann-Roch theorem implies $h^0(C, KL^{-1}) = g - d + r$. So,

$$\begin{aligned} \dim(\text{Im } \mu_0)^\perp &= g - \dim(\text{Im } \mu_0) = g - ((r + 1)(g - d + r) - \dim(\ker \mu_0)) \\ &= (g - (r + 1)(g - d + r)) + \dim(\ker \mu_0) =_{\text{def}} \rho + \dim(\ker \mu_0). \end{aligned}$$

The number ρ is called the Brill-Noether number. So, at a generic L ,

$$\dim T_L(W_d^r(C)) = \rho + \dim(\ker \mu_0).$$

Next we allow C to vary and define (with g fixed)

$$\begin{aligned} \text{Pic}^d &= \{L \rightarrow C \mid \deg_C(L) = d, g(C) = g\} \\ \cup \\ \mathcal{W}_d^r &= \{L \rightarrow C \mid \deg_C(L) = d, g(C) = g, \text{ and } h^0(C, L) \geq r + 1\}. \end{aligned}$$

The fact that these are well-defined schemes with good properties is nontrivial, but is studied carefully in [ACGH].

The tangent space $T_L(\text{Pic}^d)$ can be identified with $H^1(C, \Sigma_L)$, where Σ_L is the extension of $\mathcal{O}_C$ by Θ_C ,

$$0 \rightarrow \mathcal{O}_C \rightarrow \Sigma_L \rightarrow \Theta_C \rightarrow 0,$$

with extension class

$$\omega = c_1(L) \in H^1(C, \underline{\text{Hom}}(\Theta, \mathcal{O})) \simeq H^1(C, K).$$

This leads (nontrivially) to the definition of $\mu_1: \ker(\mu_0) \rightarrow H^0(C, K^2)$ given by

$$\mu_1\left(\sum_i s_i \otimes r_i\right) = \frac{\partial s_{i\alpha}}{\partial z_\alpha} \otimes r_{i\alpha} \in H^0(C, K^2),$$

where $\sum_i s_i \otimes r_i$ is in $\ker(\mu_0)$.

Again, by studying obstructions, it can be seen that when $h^0(C, L) = r + 1$, we have

$$(*) \quad \dim T_l(\mathcal{W}_{d,g}^r) = 3g - 3 + \rho + \dim(\ker \mu_1).$$

Thus in order to compute the dimension of $T_l(\mathcal{W}_{6,10}^2)$, one must first find $\ker(\mu_0)$ and then the dimension of $\ker(\mu_1)$.

2. Begin by fixing $l = L \rightarrow C \in \mathcal{W}_{6,10}^2 - \mathcal{W}_{6,10}^3$, where C is a smooth, genus 10, hyper-elliptic curve. A simple investigation of the map associated to the linear system $|L|$, $\varphi_{L_1}: C \rightarrow \mathbf{P}^2$ (see [Griffin 2]) shows that, in this case, $|L|$ has two basepoints and is of the form $|L| = 2g_2^1 + Q + R$, with $Q \neq iR$, where $i: C \rightarrow C$ is the involution on C . We may assume $|L| = |4P| + Q + R$, where P is a Weierstrass point on C .

In order to compute μ_0 , choose a basis for $H^0(C, L) = H^0(C, 4P + Q + R)$, say x_0, x_1, x_2 , where $\text{ord}_P(x_i) = 2i$. Then a basis for

$$H^0(C, KL^{-1}) = H^0(C, \mathcal{O}_C(10P + iQ + iR))$$

is

$$\{x_0^2, x_0x_1, x_0x_2, x_1x_2, x_2^2, y_4\} \quad \text{where } \text{ord}_P(y_4) = 10.$$

On the other hand, $H^0(C, K) = H^0(C, \mathcal{O}_C(18P))$ has basis

$$\{x_0^3, x_0^2x_1, x_0^2x_2, x_0x_1x_2, x_0x_2^2, x_1x_2^2, x_2^3, x_2y_4, z_5, z_6\} \quad \text{where } \text{ord}_P(z_j) = 2j + 6.$$

From this it is clear that μ_0 is neither injective nor surjective (μ_0 just “multiplies” $x_i \otimes x_j x_k \rightarrow x_i x_j x_k$). Its kernel has dimension 10 and basis given in

TABLE 1

$$\begin{aligned} v_1: & x_0 \otimes x_0x_1 - x_1 \otimes x_0^2 \\ v_2: & x_0 \otimes x_0x_2 - x_2 \otimes x_0^2 \\ v_3: & x_0 \otimes x_1x_2 - x_1 \otimes x_0x_2 \\ v_4: & x_0 \otimes x_1x_2 - x_2 \otimes x_0x_1 \\ v_5: & x_0 \otimes x_2^2 - x_2 \otimes x_0x_2 \\ v_6: & x_0 \otimes y_4 - x_2 \otimes x_1x_2 \\ v_7: & x_1 \otimes x_1x_2 - x_2 \otimes x_0x_2 \\ v_8: & x_1 \otimes x_2^2 - x_2 \otimes x_1x_2 \\ v_9: & x_1 \otimes y_4 - x_2 \otimes x_2^2 \\ v_{10}: & x_0 \otimes x_0x_2 - x_1 \otimes x_0x_1 \end{aligned}$$

The elements v_6, v_7, v_9, v_{10} come from relations in $H^0(C, K)$, namely $x_0x_2 - x_1^2 = 0$, $x_0y_4 - x_1x_2^2 = 0$ and $x_1y_4 - x_2^2 = 0$. The other elements are Koszul relations.

In order to compute $\ker(\mu_1)$ we will need

$$\begin{aligned}\frac{\partial x_{2\alpha}}{\partial z_\alpha} &= \frac{\partial}{\partial z_\alpha} \left(\frac{x_{1\alpha}^2}{x_{0\alpha}} \right) = \frac{2x_{1\alpha}x_{0\alpha}(\partial x_{1\alpha}/\partial z_\alpha) - x_{1\alpha}^2(\partial x_{0\alpha}/\partial z_\alpha)}{x_{0\alpha}^2} \\ &= \frac{2x_{1\alpha}(\partial x_{1\alpha}/\partial z_\alpha) - x_{2\alpha}(\partial x_{0\alpha}/\partial z_\alpha)}{x_{0\alpha}},\end{aligned}$$

which comes from the relation $x_0x_2 - x_1^2$. The α subscript will be dropped (for obvious reasons) and the dependence will be denoted by $\{ \}$'s.

We now give a table of $w_i = \mu_1(v_i)$.

TABLE 2

w_1 :	$\{x_0x_1(\partial x_0/\partial z) - x_0^2(\partial x_1/\partial z)\}$
w_2 :	$\{2x_0x_2(\partial x_0/\partial z) - 2x_0x_1(\partial x_1/\partial z)\}$
w_3 :	$\{x_1x_2(\partial x_0/\partial z) - x_0x_2(\partial x_1/\partial z)\}$
w_4 :	$\{2x_1x_2(\partial x_0/\partial z) - 2x_0x_2(\partial x_1/\partial z)\}$
w_5 :	$\{2x_2^2(\partial x_0/\partial z) - 2x_1x_2(\partial x_1/\partial z)\}$
w_6 :	$\{2y_4(\partial x_0/\partial z) - 2x_2^2(\partial x_1/\partial z)\}$
w_7 :	$\{x_2^2(\partial x_0/\partial z) - x_1x_2(\partial x_1/\partial z)\}$
w_8 :	$\{y_4(\partial x_0/\partial z) - x_2^2(\partial x_1/\partial z)\}$
w_9 :	$\{(x_1y_4/x_0)(\partial x_0/\partial z) - y_4(\partial x_1/\partial z)\}$
w_{10} :	$\{x_0x_2(\partial x_0/\partial z) - x_0x_1(\partial x_1/\partial z)\}$

Trivially, $w_2 = 2w_{10}$, $w_4 = 2w_3$, $w_5 = 2w_7$, $w_6 = 2w_8$, and in $H^0(C, K^2L)$ we have

$$\begin{aligned}x_0w_2 &= 2x_1w_1, & x_1w_2 &= 2x_0w_3, & x_0w_5 &= 2x_1w_3, \\ x_0w_6 &= x_1w_5 & \text{and} & & x_1w_6 &= 2x_0w_9.\end{aligned}$$

This clearly implies there can be *no* linear relation among $\{w_1, w_2, w_3, w_5, w_6, w_9\}$ in $H^0(C, K^2)$. Consequently, the kernel of μ_1 has basis $\{w_2 - 2w_{10}, w_4 - 2w_3, w_5 - 2w_7, w_6 - 2w_8\}$. By equation (*) we finally have

$$\begin{aligned}\dim T_l(\mathcal{W}_{6,10}^2) &= 3g - 3 + \rho + \dim(\ker \mu_1) \\ &= 3(10) - 3 + 10 - 3(10 - 6 + 2) + 4 = 23.\end{aligned}$$

It is easy to compute the dimension of the hyper-elliptic component W of $\mathcal{W}_{6,10}^2$. Since the g_2^1 on a hyper-elliptic curve is unique and $l = L \rightarrow C$, where $|L| = 2g_2^1 + P + Q$, one has immediately that

$$\dim W = \dim\{\text{genus } 10, \text{hyper-elliptic curves}\} + 2.$$

By Hurwitz's formula,

$$\dim\{\text{hyper-elliptic curves of genus } g\} = 2g - 1.$$

Therefore,

$$\dim W = 19 + 2 = 21!$$

Thus, since $\dim T_l(\mathcal{W}_{6,10}^2) = 23 > \dim_l(\mathcal{W}_{6,10}^2) = 21$, we conclude

THEOREM. *The “component of hyper-elliptics” in $\mathcal{W}_{6,10}^2$ is nonreduced.*

A final remark: In general the singularities of $\mathcal{W}_{d,g}^r$ are “worst” along $\mathcal{W}_{d,g}^{r+1}$, and in fact one can show, using these methods, that if $l \in \mathcal{W}_{6,10}^3$,

$$\dim T_l(\mathcal{W}_{6,10}^2) > 23.$$

In spite of this, $\mathcal{W}_{6,10}^3$ is *smooth*, i.e.

$$\dim T_l(\mathcal{W}_{6,10}^3) = \dim_l(\mathcal{W}_{6,10}^3) = 19$$

for all $l \in \mathcal{W}_{6,10}^3$. So it is very important to distinguish between $T_l(\mathcal{W}_{d,g}^{r+1})$ and $T_l(\mathcal{W}_{d,g}^r)$ when $l \in \mathcal{W}_{d,g}^{r+1} \subseteq \mathcal{W}_{d,g}^r$.

REFERENCES

[ACGH] E. Arbarello, M. Cornalba, P. A. Griffiths and J. Harris, *Topics in the theory of algebraic curves*, Princeton Univ. Press. (to appear).

[Griffin 1] E. Griffin, *Special fibers in families of plane curves*, Ph. D. Thesis, Harvard Univ., 1982.

[Griffin 2] ———, *The component structure of $\mathcal{W}_6^2$* (to appear).

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, CAMBRIDGE, MASSACHUSETTS 02138

Current address: Department of Mathematics, University of California, Los Angeles, California 90024