HYPERBOLICITY OF A COMPLEX MANIFOLD AND OTHER EQUIVALENT PROPERTIES

KYONG T. HAHN¹ AND KANG T. KIM

ABSTRACT. Defining the notions of Schottky, Landau and Picard properties on a plane domain, the first author [3] proved that a domain in $\mathbb{C}$ having any of these properties is equivalent to the hyperbolicity of the domain.

In this paper the authors extend these notions to higher-dimensional case and obtain other various equivalent conditions for the hyperbolicity of a complex manifold.

1. Introduction. Let M be a connected complex manifold of dimension m with hermitian metric h_M . Let ρ_M denote the distance function associated with h_M . Following S. Kobayashi [6], we define M to be (Kobayashi) *hyperbolic* if the Kobayashi pseudometric k_M is a metric. In [8], H. Royden has constructed the infinitesimal form of the Kobayashi pseudometric

$$(1) \quad K_M(p, \xi) = \inf\{|v| : \exists f \in \mathcal{H}(\Delta, M) \ni f(0) = p, f'(0)v = \xi\}$$

where $(p, \xi) \in T(M)$, the tangent bundle of M , and $\mathcal{H}(\Delta, M)$ denotes the class of all holomorphic maps f of the open disc Δ into M , and he has shown that if M is hyperbolic then k_M is the integrated form of K_M . The notion of hyperbolicity of M can also be defined in terms of K_M : A complex manifold M is hyperbolic if for each $p \in M$ there exists a coordinate neighborhood U of p and a positive constant c such that for $(q, \eta) \in T(U) \simeq U \times \mathbb{C}^m$

$$(2a) \quad K_M(q, \eta) \geq c\|\eta\|,$$

where $\|\cdot\|$ denotes the euclidean distance. This definition of hyperbolicity coincides with that of Kobayashi (see [8]). In the case of a hermitian manifold, the above condition is equivalent to the condition that for each $p \in M$ there is a neighborhood U of p and a positive constant c such that

$$(2b) \quad K_M(q, \eta) \geq ch_M(q, \eta)$$

for all $(q, \eta) \in T(U)$.

DEFINITION 1. A complex manifold M is said to satisfy the *Schottky property* if for each $p \in M$, for each relatively compact open set W in a coordinate neighborhood of p and an $r \in (0, 1)$, there exists a positive constant $S = S(W, r)$ such that every

Received by the editors June 23, 1983.

1980 *Mathematics Subject Classification.* Primary 32H20, 32H25; Secondary 32H15.

Key words and phrases. Schottky property, Landau property, Picard property, Bloch mapping, Kobayashi metric and hyperbolic manifold.

¹Research partially supported by NSF Grant MCS 80-02915 at Pennsylvania State University.

holomorphic map $f \in \mathcal{H}(\Delta, M)$ with $f(0) \in W$ satisfies

$$(3) \quad \rho_M(p, f(z)) \leq S \quad \text{for } |z| \leq r.$$

M satisfies the *Landau property* if for each $p \in M$ and each relatively compact open set W in a coordinate neighborhood of p , there exists a positive constant $R = R(W)$ such that

$$(4) \quad \sup\{h_M(f(0), f'(0)): f \in \mathcal{H}(\Delta, M) \text{ with } f(0) \in W\} \leq R.$$

Then we can prove the following.

THEOREM 1. *The following statements are equivalent on any hermitian manifold M .*

- (a) M is hyperbolic.
- (b) M satisfies the Schottky property.
- (c) M satisfies the Landau property.

DEFINITION 2. A map $f \in \mathcal{H}(\Delta, M)$ is said to be *Bloch* (see [4]) if

$$(5a) \quad \sup\{Q_f(z): z \in \Delta\} < \infty,$$

where

$$(5b) \quad Q_f(z) = (1 - |z|^2)h_M(f(z), f'(z)).$$

We remark that the nonnegative function Q_f is *invariant* under the group $\text{Aut}(\Delta)$ of holomorphic automorphisms of Δ in the sense that for all $\varphi \in \text{Aut}(\Delta)$

$$(6) \quad Q_{f \circ \varphi}(z) = Q_f(\varphi(z)), \quad z \in \Delta.$$

It is an easy consequence of the definition of Kobayashi metric that if (5a) holds for all $f \in \mathcal{H}(\Delta, M)$, then M is hyperbolic. The converse is, however, not true in general (see lemma in §3). If M is compact, we obtain the following characterization of the hyperbolicity of M .

THEOREM 2. *Let M be a connected compact hermitian manifold. The following statements are equivalent.*

- (a) M is hyperbolic.
- (b) $\sup\{h_M(f(0), f'(0)): f \in \mathcal{H}(\Delta, M)\} < \infty$.
- (c) $\sup\{Q_f(z): z \in \Delta\} < \infty$ for all $f \in \mathcal{H}(\Delta, M)$, i.e., every $f \in \mathcal{H}(\Delta, M)$ is Bloch.
- (d) M has the Picard property, i.e., there is no nonconstant holomorphic map $f: \mathbb{C} \rightarrow M$.

Unlike in the one complex-dimensional case [3], the notion of the Picard property is a weaker notion than the Schottky property, the Landau property or the hyperbolicity for a *noncompact* manifold [6].

We note that the notion of the Picard property used in [3] and here, somewhat reluctantly, coincides with the term previously used by S. Kobayashi [6] and others.

Finally, we acknowledge our indebtedness to S. Krantz and also to the referee for their helpful comments on the paper. In particular, the example given in the proof of the second half of the lemma was suggested by the referee. This example is much simpler than the one originally given by the authors.

2. Proof of Theorem 1. It will be proved through the following implications:

(a) $\Leftrightarrow$ (b), (a) $\Rightarrow$ (c) $\Rightarrow$ (a).

To prove (a) $\Rightarrow$ (b), let $p \in M$ and U be its local coordinate neighborhood. Let W be a relatively compact neighborhood of p in U . Since k_M is a metric, it induces the standard topology in M [1]. Therefore, we may define $W = \{q: k_M(p, q) < \rho_0\}$ for some $\rho_0 > 0$ and $U = \{q: k_M(p, q) < \rho_0 + \rho_1\}$, where $\rho_1 = \tan h^{-1}r$, for any given $r \in (0, 1)$. If $f \in \mathcal{H}(\Delta, M)$ satisfies $f(0) \in W$, then $k_M(p, f(0)) < \rho_0$ and $k_M(f(0), f(z)) \leq k_\Delta(0, z) \leq \rho_\Delta(0, r) = \tan h^{-1}r = \rho_1$ for $|z| \leq r$. Therefore, it follows from the triangle inequality that $k_M(p, f(z)) \leq \rho_0 + \rho_1$ whenever $|z| \leq r$. Since k_M is a metric, $k_M \geq c\rho_M$ for some $c > 0$ which proves (b) with $S = (\rho_0 + \rho_1)/c$.

For the proof of (b) $\Rightarrow$ (a), we follow the method of [5]: Let p and q be two distinct points of M . We may take a coordinate neighborhood U of p which does not contain q . Let $\varphi: U \rightarrow B = \{w \in \mathbb{C}^m: \|w\| < 1\}$ be a biholomorphic map with $\varphi(p) = 0$, $\varphi(U) = B$. Without loss of generality, we may assume that $\varphi(W) = \{w \in \mathbb{C}^m: \|w\| < \rho\} = B_\rho$ for some $\rho \in (0, 1)$ and $\varphi(V) \subset B$, $V = \{m \in M: \rho_M(p, m) < S\}$. Let $r \in (0, 1)$ be given. If $|z| < r/2$, then there exists a constant $c > 0$ such that $\tan h^{-1}|z| \geq c \tan h^{-1}|z|/r$ or, equivalently, $\rho_\Delta(0, z) \geq c\rho_\Delta(0, z)$. Let $\alpha = \{p = p_0, p_1, \dots, p_l = q; a_1, \dots, a_l; f_1, \dots, f_l\}$ be any chain connecting p and q , used in the construction of the Kobayashi metric $k_M(p, q)$ [6]. Since $\varphi(q)$ lies outside B , there must be an index, say l_0 , $0 \leq l_0 < l$, such that $\varphi(p_0), \varphi(p_1), \dots, \varphi(p_{l_0-1}) \in B_\rho$ and $\varphi(p_{l_0}) \notin B_\rho$. By the homogeneity of Δ , we may assume that $a_1, \dots, a_{l_0}$ lie in $\Delta_{r/2}$. Then

$$\begin{aligned} (1) \quad |\alpha| &\geq \sum_{i=1}^{l_0} \rho_\Delta(0, a_i) \geq c \sum_{i=1}^{l_0} \rho_{\Delta_r}(0, a_i) \geq c \sum_{i=1}^{l_0} k_B(\varphi(p_{i-1}), \varphi(p_i)) \\ &\geq ck_B(0, \varphi(p_{l_0})) = c \tan h^{-1} \|\varphi(p_{l_0})\| \geq c \tan h^{-1} \rho \equiv c'. \end{aligned}$$

Thus, $k_M(p, q) \geq c' > 0$. Here we used the fact that on B the Kobayashi metric agrees with the standard Kähler metric. It is given by

$$(2) \quad k_B(0, z) = \frac{1}{2} \log \frac{1 + \|z\|}{1 - \|z\|} = \tan h^{-1} \|z\| \quad \text{for } z \in B.$$

See [4].

(a) $\Rightarrow$ (c). Suppose that M fails to satisfy the Landau property. Then there exist a point p_0 in an open neighborhood W which is relatively compact in a local coordinate neighborhood U of p_0 and a sequence $f_k \in \mathcal{H}(\Delta, M)$ such that $f_k(0) \in W$ and $h_M(f_k(0), f'_k(0)) \rightarrow \infty$ as $k \rightarrow \infty$. We claim that if M is hyperbolic then there exists an $r \in (0, 1)$ such that $\{f_k\}$ contains a subsequence which converges uniformly to a holomorphic map $f: \Delta_r \rightarrow M$. First we observe that $\overline{W} \cap (M \setminus U) = \emptyset$ where $\overline{W}$ is compact and $M \setminus U$ is closed. Therefore, there exists a number $\rho' > 0$ such that

$$(3) \quad Q = \{p \in M: k_M(\overline{W}, p) < \rho'\} \subset \subset U.$$

Since $k_M(f_k(0), f_k(z)) \leq \rho_\Delta(0, z) = \tan h^{-1}|z|$ for all $f_k \in \mathcal{H}(\Delta, M)$ with $f_k(0) \in W$, $f_k(z) \in Q$ whenever $|z| \leq r' = \tan h \rho'$. Since Q is relatively compact, it is bounded in $\mathbb{C}^m$. Therefore, by Montel's theorem there is a subsequence of $\{f_k\}$ which

converges uniformly on Δ_r to a holomorphic map $f: \Delta_r \rightarrow M$ for $r < r'$. Denote again this convergent subsequence by $\{f_k\}$. By Weierstrass' theorem, $f'_k(z)$ converges to $f'(z)$ uniformly on Δ_r . In particular, $f'_k(0)$ converges to $f'(0)$. Thus, $h_M(f_k(0), f'_k(0)) \rightarrow h_M(f(0), f'(0)) < \infty$ as $k \rightarrow \infty$, which is a contradiction.

(c) $\Rightarrow$ (a). Given $(q, \xi) \in T(W)$ and $v \in \mathbb{C}$, let $f \in \mathcal{K}(\Delta, M)$ satisfy $f(0) = q$ and $f'(0)v = \xi$. Then by (4) of §1

$$R|v| \geq h_M(f(0), f'(0)|v|) = h_M(f(0), f'(0)v) = h_M(q, \xi)$$

or

$$(4) \quad |v| \geq \frac{1}{R} h_M(q, \xi),$$

where R is the upper bound given in (4) of §1. Thus,

$$(5) \quad K_M(q, \xi) \geq ch_M(q, \xi),$$

$c = 1/R$, as desired.

3. Proof of Theorem 2. By a result of R. Brody [2], (a) $\Leftrightarrow$ (d) holds on a connected compact manifold. Therefore, it is enough to prove the implications (a) $\Rightarrow$ (b) $\Rightarrow$ (c) $\Rightarrow$ (a). First we prove the following

LEMMA. *A complex manifold M is hyperbolic if $\sup\{Q_f(z): z \in \Delta\} < \infty$ for all $f \in \mathcal{K}(\Delta, M)$. The converse is not true in general.*

PROOF. The above hypothesis implies that there exists a constant $Q > 0$ such that for all $f \in \mathcal{K}(\Delta, M)$

$$(1) \quad h_M(f(z), f'(z)v) \leq QK_\Delta(z, v) \quad (z \in \Delta, v \in \mathbb{C}).$$

Let $\tilde{\rho}_M$ be the distance function associated with the differential metric $\tilde{h}_M = Q^{-1}h_M$. By integrating both sides of (1) along the geodesic curve between two points z and w in Δ and using the definition of integrated distance $\tilde{\rho}_M$, we have

$$(2) \quad \tilde{\rho}_M(f(z), f(w)) \leq k_\Delta(z, w)$$

for all $f \in \mathcal{K}(\Delta, M)$. Since k_M is the largest among those pseudometrics which satisfy (2), $\tilde{\rho}_M \leq k_M$ (see [6]). Thus, M is hyperbolic.

To show the second half, let M be the right half-plane of $\mathbb{C}$ and h_M the euclidean metric. Then M is clearly hyperbolic. Set

$$(3) \quad f(z) = (1+z)/(1-z) \quad \text{and} \quad z_n = 1 - 1/n \in \Delta.$$

Then $f \in \mathcal{K}(\Delta, M)$ and

$$Q_f(z_n) = (1 - |z_n|^2)|f'(z_n)| = 2(2n - 1) \rightarrow \infty$$

as $n \rightarrow \infty$. This completes the lemma.

From this lemma, (c) $\Rightarrow$ (a) trivially follows. To prove (a) $\Rightarrow$ (b), let M be hyperbolic. By (c) of Theorem 1, for each $p \in M$ there exists a relatively compact open neighborhood W_p of p such that

$$(4) \quad \sup\{\|f'(0)\|: f \in \mathcal{K}(\Delta, M), f(0) \in W_p\} < \infty.$$

Since M is compact, it can be covered by a finite number of such W_p 's. Thus, (b) follows.

(b) $\Rightarrow$ (c). Suppose (c) fails to hold. Then there exist sequences $\{z_n\}$ in Δ and $\{f_n\}$ in $\mathcal{H}(\Delta, M)$ such that $Q_{f_n}(z_n) > n$ for all n . Since Δ is homogeneous, for each z_n there exists $\varphi_n \in \text{Aut}(\Delta)$ such that $\varphi_n(0) = z_n$. Therefore, by the invariant property of Q_f (see (6) of §1),

$$(5) \quad Q_{f_n}(z_n) = Q_{f_n}(\varphi_n(0)) = Q_{g_n}(0) = h_M(g_n(0), g'_n(0)),$$

where $g_n = f_n \circ \varphi_n$. Since h_M is hermitian, it follows from the compactness of M , condition (b) and (5) that $Q_{f_n}(z_n)$ is bounded for all n , which is a contradiction.

REFERENCES

1. T. J. Barth, *The Kobayashi distance induces the standard topology*, Proc. Amer. Math. Soc. **35** (1972), 439–441. MR **46** #5668.
2. R. Brody, *Compact manifolds and hyperbolicity*, Bull. Amer. Math. Soc. **235** (1978), 213–219.
3. K. T. Hahn, *Equivalence of the classical theorems of Schottky, Landau, Picard and hyperbolicity*, Proc. Amer. Math. Soc. **89** (1983), 628–632.
4. ———, *Asymptotic behavior of normal mappings of several complex variables*, Canad. J. Math. (to appear).
5. P. Kiernan, *On the relation between taut, tight and hyperbolic manifolds*, Bull. Amer. Math. Soc. **76** (1970), 49–51. MR **40** #5896.
6. S. Kobayashi, *Hyperbolic manifolds and holomorphic mappings*, Pure and Appl. Math., vol. 2, Dekker, New York, 1970. MR **43** #3503.
7. M. Kwack, *Some classical theorems for holomorphic mappings into hyperbolic manifolds*, Proc. Sympos. Pure Math., vol. 27, Amer. Math. Soc., Providence, R. I., 1975, pp. 99–104.
8. H. Royden, *Remarks on the Kobayashi metric* (Proc. Maryland Conf. Several Complex Variables), Lecture Notes in Math., vol. 185, Springer-Verlag, Berlin, 1971, pp. 125–137. MR **46** #3826.

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PENNSYLVANIA 16802

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA 90024