APPROXIMATE UNITARY EQUIVALENCE OF POWER PARTIAL ISOMETRIES

KENNETH R. DAVIDSON¹

ABSTRACT. Every power partial isometry (p.p.i.) in the Calkin algebra lifts to a p.p.i. in $\mathcal{B}(\mathcal{H})$.

An element u in a C^* algebra is a power partial isometry (p.p.i.) if u^n is a partial isometry for every integer $n \geq 1$. This notion was introduced by Halmos and Wallen [5], and they characterized all p.p.i.'s in $\mathcal{B}(\mathcal{H})$. In a penetrating study [6], Herrero classifies p.p.i.'s up to their unitary orbits and up to unitary equivalence modulo the compacts. In this note, we show that every p.p.i. in the Calkin algebra lifts to a p.p.i. in $\mathcal{B}(\mathcal{H})$. This is done by using a few straightforward computations involving the theory of C^* extensions. Some of the results of [6] follow from this method as well.

In this paper, Hilbert spaces are always separable. The ideal of compact operators will be denoted by $\mathcal{K}$. The canonical quotient map of $\mathcal{B}(\mathcal{H})$ onto the Calkin algebra $\mathcal{B}(\mathcal{H})/\mathcal{K}$ is denoted by π . Two elements s and t of the Calkin algebra are (strongly) compalant ($s \cong t$) if there is a unitary U in $\mathcal{B}(\mathcal{H})$ so that $t = \pi(U)s\pi(U^*)$. They are weakly compalant ($s \cong_w t$) if there is a unitary u in $\mathcal{B}(\mathcal{H})/\mathcal{K}$ so that $t = usu^*$. For p.p.i.'s, these two notions may differ. Given a nuclear C^* algebra $\mathcal{A}$, $\text{Ext } \mathcal{A}$ denotes the group of extensions of $\mathcal{K}$ by $\mathcal{A}$ modulo compalance [1, 3, 8] and $\text{Ext}_w \mathcal{A}$ denotes the quotient of $\text{Ext } \mathcal{A}$ modulo weak compalance. We use little more than the fact that when

$$0 \rightarrow \mathcal{J} \rightarrow \mathcal{A} \rightarrow \mathcal{A}/\mathcal{J} \rightarrow 0$$

is an exact sequence, and $\text{Ext } \mathcal{J} = 0 = \text{Ext } \mathcal{A}/\mathcal{J}$, then $\text{Ext } \mathcal{A} = 0$.

The first step is to classify p.p.i.'s up to algebraic equivalence by computing $C^*(T)$ for all p.p.i.'s. That is, given two p.p.i.'s S and T , when is there a C^* isomorphism ϕ of $C^*(S)$ onto $C^*(T)$ such that $\phi(S) = T$? For convenience, this will be written $S \sim T$. Let N_k be the nilpotent Jordan cell of order k acting on a k -dimensional Hilbert space, and let S be the unilateral shift. If α belongs to $\mathbb{N}_0 \cup \{\omega\}$, let $A^{(\alpha)}$ be the direct sum of α copies of A acting on $\mathcal{H}^{(\alpha)}$, the direct sum of α copies of $\mathcal{H}$. By [5], every p.p.i. T represented on a Hilbert space can be decomposed as

$$T \cong \bigoplus_{k=1}^{\infty} N_k^{(\alpha_k)} \oplus S^{(\alpha)} \oplus S^{*(\beta)} \oplus V$$

where V is a unitary. (If $\alpha_k = 0$, the summand is deemed to be absent.) The existence of the finite nilpotent blocks is easy to determine because there is a

Received by the editors July 9, 1983.

1980 *Mathematics Subject Classification*. Primary 47B99, 47C15; Secondary 46M20.

¹Research partially supported by NSERC grant #A3488. I would also like to thank the people at Indiana University for their hospitality during the period when this work was completed.

polynomial $p_k(x, y)$ in two noncommuting variables such that $P_k = p_k(T, T^*)$ is a (selfadjoint) projection onto the summand $\mathcal{H}_k^{(\alpha_k)}$ on which $N_k^{(\alpha_k)}$ acts. The classification in [5] proceeds by pulling off all the nilpotent blocks, then one removes the isometric and coisometric parts, and what remains is the unitary.

This procedure is not possible in an arbitrary C^* algebra because one can pull off only finitely many pieces. Also, if T has no nilpotent blocks and is not unitary, then $C^*(T)$ has only a few possibilities up to algebraic equivalence. If Σ is a nonempty compact subset of the unit circle S^1 , let V_Σ be a fixed unitary with spectrum Σ .

PROPOSITION 1. (a) *If T is a p.p.i. with no nilpotent blocks, then there are only the following possibilities:*

(a1) *T is unitary ($TT^* = I = T^*T$), in which case the spectrum $\Sigma = \sigma(T)$ determines $C^*(T)$ up to algebraic equivalence: $T \sim V_\Sigma$.*

(a2) *T is a nonunitary isometry ($T^*T = I \neq TT^*$), in which case $T \sim S$.*

(a3) *T is a nonunitary coisometry ($T^*T \neq I = TT^*$), in which case $T \sim S^*$.*

(a4) *T is neither isometric nor coisometric, in which case $T \sim S \oplus S^*$.*

(b) *If T is a p.p.i. with finitely many nilpotent blocks of orders $k_1, \dots, k_n$, then $P_0 = I - \sum_{i=1}^n P_{k_i}$ is a projection in $C^*(T)$ and $T_1 = P_0 T|_{P_0}$ is determined up to algebraic equivalence by (a). So $T \sim \bigoplus_{i=1}^n N_{k_i} \oplus T_1$.*

(c) *If T is a p.p.i. with nilpotent blocks of orders $k_1 < k_2 < k_3 < \dots$, then $T \sim \bigoplus_{i=1}^\infty N_{k_i}$.*

PROOF. If T is a p.p.i. in a C^* algebra $\mathcal{A}$, represent $\mathcal{A}$ on a Hilbert space and apply the theorem of [5]. Most of this proposition is straightforward, based on the simple fact that for any representation π of $C^*(T)$, one has $T \sim T \oplus \pi(T)$.

Case (a1) follows from the spectral theorem. Since there is a $*$ homomorphism of $C^*(S)$ onto $C(S^1)$ taking S to the identity function z , one has $S \sim S^{(\alpha)} \oplus V$ for any $\alpha \geq 1$ and any unitary V . Case (a3) is similar. Finally, the same remarks show that $S \oplus S^* \sim S^{(\alpha)} \oplus S^{*(\beta)} \oplus V$ for any $\alpha \geq 1, \beta \geq 1$ and any unitary V . The projections P_k onto the nilpotent part of order k belong to $C^*(T)$. Thus if there are only finitely many P_{k_i} 's which are not zero, one can define $P_0 = I - \sum_{i=1}^n P_{k_i}$. This pulls out a direct summand T_1 which is analyzed by part (a).

Given an infinite sequence $k_1 < k_2 < k_3 < \dots$, consider the p.p.i. $W = \bigoplus_{i=1}^\infty N_{k_i}$ acting on $\mathcal{H} = \bigoplus_{i=1}^\infty \mathcal{H}_i$. Let J be the ideal $C^*(W) \cap \mathcal{K}$, where $\mathcal{K}$ is the compact operators. Clearly J contains the projection P_{k_i} onto $\mathcal{H}_i$, and thus contains all of $\mathcal{B}(\mathcal{H}_i) \cong M_{k_i}(\mathbb{C})$ for each $i \geq 1$. It follows that J is the c_0 direct sum $\bigoplus_{i=1}^\infty \mathcal{B}(\mathcal{H}_i)$. The quotient $C^*(W)/J$ is isomorphic to $C^*(\mathfrak{w})$ where $\mathfrak{w}$ is the image of W in the Calkin algebra. Because P_k is compact, $p_k(\mathfrak{w}) = \pi P_k = 0$ so $\mathfrak{w}$ is a p.p.i. with no nilpotent summands. Also,

$$\mathfrak{w}\mathfrak{w}^* = \pi WW^* \neq 1 \quad \text{and} \quad \mathfrak{w}^*\mathfrak{w} = \pi W^*W \neq 1,$$

so $\mathfrak{w}$ is neither isometric nor coisometric. By (a), $\mathfrak{w} \sim S \oplus S^*$. Thus by (a) and (b),

$$W \sim \bigoplus_{i=1}^\infty N_{k_i}^{(\alpha_i)} \oplus S^{(\alpha)} \oplus S^{*(\beta)} \oplus V$$

for any $\alpha_i \geq 1$, any α, β and any unitary V . $\square$

THEOREM 2. *Every p.p.i. $\mathfrak{t}$ in the Calkin algebra lifts to a p.p.i. in $\mathcal{B}(\mathcal{H})$. These p.p.i.'s are classified up to unitary equivalence in the Calkin algebra as follows.*

(a1) *If $\mathfrak{t}$ is unitary and $\sigma(\mathfrak{t}) = \Sigma$ is a proper subset of S^1 , then $\mathfrak{t} \cong \pi(V_\Sigma^{(\infty)})$.*

- (a2) If t is unitary and $\sigma(t) = S^1$, and $\text{ind } t = n$, then $t \cong \pi(S^{(-n)})$ if $n < 0$, $t \cong \pi(V_{S_1})$ if $n = 0$, and $t \cong \pi(S^{*(n)})$ if $n > 0$.
- (b) If t is isometric but not unitary, $t \cong \pi(S^{(\infty)})$.
- (c) If t is coisometric but not unitary, $t \cong \pi(S^{*(\infty)})$.
- (d) If t has no nilpotent part and is neither isometric nor coisometric, then $t \cong \pi(S^{(\infty)} \oplus S^{*(\infty)})$.
- (e) If t has finitely many nilpotent parts $k_1, \dots, k_n$ with corresponding projections $p_1, \dots, p_n$, then
- (e1-4) if $p_0 = 1 - \sum_{i=1}^n p_i$ is not zero, then $t \cong \pi(\bigoplus_{i=1}^n N_{k_i}^{(\infty)} \oplus T_1)$ where T_1 is determined by (a)-(d),
- (e5) if $p = 0$, then there is an integer d with $0 \leq d < \text{g.c.d.}\{k_1, \dots, k_n\}$ such that $t \cong \pi(\bigoplus_{i=1}^n N_{k_i}^{(\infty)} \oplus \mathcal{O}_d)$ where $\mathcal{O}_d = N_1^{(d)}$ is the zero operator on a d -dimensional space.
- (f) If t has nilpotent parts for the infinite sequence $k_1 < k_2 < k_3 < \dots$, then $t \cong \pi(\bigoplus_{i=1}^{\infty} N_{k_i}^{(\infty)})$.

PROOF. Cases (a1) and (a2) follow from early results of [2]. Cases (b) and (c) are well known [4]. This is also immediate from the fact that $\text{Ext}(C^*(S)) = 0$. For case (d), we have $t \sim S \oplus S^* \sim \pi(S^{(\infty)} \oplus S^{*(\infty)})$. It suffices to show that $\text{Ext}(C^*(S \oplus S^*)) = 0$. Let $T = S \oplus S^*$, and let J be the ideal in $C^*(T)$ generated by $I - T^*T$. It is easy to see that J is isomorphic to the compact operators $\mathcal{K}$, and that $C^*(T)/J$ is isomorphic to $C^*(S)$. Thus

$$0 \rightarrow \mathcal{K} \rightarrow C^*(T) \rightarrow C^*(S) \rightarrow 0$$

is exact. But $\text{Ext } \mathcal{K} = 0 = \text{Ext } C^*(S)$, and hence $\text{Ext } C^*(T) = 0$. So t is unitarily equivalent to $\pi(S^{(\infty)} \oplus S^{*(\infty)})$.

In case (e5), $C^*(t)$ is isomorphic to $\bigoplus_{i=1}^n M_{k_i}(\mathbb{C})$, and thus $\text{Ext } C^*(t)$ is isomorphic to $\mathbb{Z}/p\mathbb{Z}$ where $p = \text{g.c.d.}\{k_1, \dots, k_n\}$ and $\text{Ext}_w C^*(t) = 0$ [7]. It follows readily that if d is the integer with $0 \leq d < p$ such that $d + p\mathbb{Z}$ is the element of $\text{Ext } C^*(t)$ determined by t , then $t \cong \pi(\bigoplus_{i=1}^n N_{k_i}^{(\infty)} \oplus \mathcal{O}_d)$, and $t \cong_w \pi(\bigoplus_{i=1}^n N_{k_i}^{(\infty)})$.

When $p_0 \neq 0$, let $t = t_0 \oplus t_1$ where $p_0 = 0 \oplus 1$. The above analysis shows that

$$t \cong \pi\left(\bigoplus_{i=1}^n N_{k_i}^{(\infty)} \oplus \mathcal{O}_d \oplus T_1\right) \cong \pi\left(\bigoplus_{i=1}^n N_{k_i}^{(\infty)}\right) \oplus \pi(\mathcal{O}_d \oplus T_1).$$

The latter term has no nilpotent part, so it is analyzed by parts (a) and (b).

Finally, if t has nilpotent parts for the infinite sequence $k_1 < k_2 < k_3 < \dots$, we refer back to the analysis of Proposition 1 to obtain an ideal J isomorphic to $\bigoplus_{i=1}^{\infty} M_{k_i}(\mathbb{C})$ in $C^*(t)$ so that

$$0 \rightarrow J \rightarrow C^*(t) \rightarrow C^*(S \oplus S^*) \rightarrow 0.$$

By [7], $\text{Ext } J = 0$. Since $\text{Ext } C^*(S \oplus S^*) = 0$, it follows that $\text{Ext } C^*(t) = 0$. Thus t is unitarily equivalent to $\pi(\bigoplus_{i=1}^{\infty} N_{k_i}^{(\infty)})$. $\square$

REMARKS. (1) One of the most surprising facts, noted first in [5], is that $\bigoplus_{k=1}^{\infty} N_k$ is approximately unitarily equivalent to $S^{(\infty)} \oplus S^{*(\infty)}$. This is now "explained" by noting that, in the Calkin algebra, both have nonisometric, non-coisometric images without nilpotent parts, and hence the images are algebraically equivalent to $S \oplus S^*$. This can be represented in the Calkin algebra in only one way.

(2) Except in case (e5) in which $C^*(t)$ is finite dimensional, weak and strong equivalence coincide for p.p.i.'s.

(3) The classification of p.p.i.'s in $\mathcal{B}(\mathcal{H})$ up to approximate unitary equivalence is carried out in [6]. These results are also immediate from Theorem 2. In fact, it would be sufficient to show every p.p.i. in the Calkin algebra lifts to a p.p.i. without classifying them as one goes along. But the nature of our method forces us to do it all.

REFERENCES

1. W. B. Arveson, *Notes on extensions of C^* algebras*, Duke Math. J. **44** (1977), 329–355.
2. L. G. Brown, R. G. Douglas and P. A. Fillmore, *Unitary equivalence modulo the compact operators and extensions of C^* algebras*, Lecture Notes in Math., Vol. 345, Springer-Verlag, Berlin and New York, 1973.
3. M. D. Choi and E. G. Effros, *The completely positive lifting problem*, Ann. of Math. (2) **104** (1976), 265–324.
4. P. R. Halmos, *Limits of shifts*, Acta Sci. Math. (Szeged) **34** (1973), 131–139.
5. P. R. Halmos and L. J. Wallen, *Powers of partial isometries*, J. Math. Mech. **19** (1970), 657–663.
6. D. A. Herrero, *Unitary orbits of power partial isometries and approximation by block diagonal nilpotents*, Topics in Modern Operator Theory, Birkhäuser, Basel, 1981, pp. 171–210.
7. M. Pimsner and S. Popa, *On the Ext-group of an AF-algebra*, Rev. Roumaine Math. Pures Appl. **23** (1978), 251–267.
8. D. Voiculescu, *A non-commutative Weyl-von Neumann theorem*, Rev. Roumaine Math. Pures Appl. **21** (1976), 97–113.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO
N2L 3G1, CANADA