A SHORT PROOF OF THE CIMA-WOGEN $L(f) = \text{CIRCLE}$ THEOREM

JOSÉ A. ANTONINO AND SALVADOR ROMAGUERA

ABSTRACT. In [1] Cima and Wogen showed that if $f \in \text{ball } \tilde{B}_0$ and $L(f)$ contains a circle γ , then $\gamma = L(f)$. This note presents a new and straightforward proof of Cima and Wogen's theorem.

Let $D = \{z: |z| < 1\}$ and let f be a holomorphic function in D . Following the notation of [1] let

$$M(f) = \sup\{|f'(z)|(1 - |z|^2): z \in D\}$$

and

$$B = \{f: f \text{ holomorphic in } D \text{ and } M(f) < \infty\},$$

B is the Bloch space with the norm $\|f\| = M(f) + |f(0)|$.

Now let

$$B_0 = \left\{ f \in B: \lim_{|z| \rightarrow 1} |f'(z)|(1 - |z|^2) = 0 \right\},$$

$$\tilde{B} = \{f \in B: f(0) = 0\}, \quad \tilde{B}_0 = B_0 \cap \tilde{B}$$

and

$$\text{ball } \tilde{B}_0 = \{f \in \tilde{B}_0: M(f) \leq 1\}.$$

If $R' = \{f: f \text{ holomorphic in } D \text{ and } M(f) \leq 1\}$, then it is obvious that $\text{ball } \tilde{B}_0 \subset R'$.

For each $f \in R'$ let

$$L(f) = \{z \in D: |f'(z)|(1 - |z|^2) = 1\}.$$

Now let $a \in D$ and let $r > 0$ such that $r < 1 - |a|$. The circle of center a and radius r is denoted by $C(a, r)$.

THEOREM 1. *For each $a \in D$ there is a sequence of real numbers $\{\delta_n(a)\}$ such that if $f \in R'$ and $L(f)$ contains a sequence of distinct points $\{z_k\}$ such that $|z_k| = r$ for every k , then $L(f) = C(a, r)$ and $r = \delta_n(a)$ for some n .*

PROOF. The function

$$(1) \quad h(z) = f''(z)/(zf''(z) + 2f'(z))$$

is meromorphic in D and $h(z) = \bar{z}$ on $L(f)$ (see [2]).

Since $\{z_k\} \subset L(f)$ we have

$$h(z_k) = \bar{z}_k = r^2/z_k \quad \text{for every } k.$$

Received by the editors November 11, 1983.

1980 *Mathematics Subject Classification.* Primary 30A10.

Key words and phrases. Holomorphic function, Bloch space, identity principle.

©1984 American Mathematical Society
 0002-9939/84 \$1.00 + \$.25 per page

Hence, by using the identity principle (note that $\{z_k\}$ has an accumulation point s with $|s| = r$) we get $h(z) = r^2/z$.

An integration on (1) yields $f'(z) = cz^\alpha$ with c a complex constant and $\alpha = 2r^2/(1-r^2)$. Condition $f \in R'$ implies $2r^2/(1-r^2) = n$ for some n . Consequently, $r = (n/(n+2))^{1/2} = \delta_n(0)$. Also we have that $c = \lambda/r^n(1-r^2)$ with $|\lambda| = 1$. Finally, it is clear that $|f'(z)|(1-|z|^2) < 1$ if $|z| \neq r$ since the real-valued function $g(x) = x^n(1-x^2)/r^n(1-r^2)$, $0 < x < 1$, satisfies $g(x) < 1$ if $x \neq r$. Then

$$f'(z) = \lambda z^n / r^n (1 - r^2)$$

with $L(f) = C(0, r)$ and $r = \delta_n(0) = \delta_n$ (shortening).

Now let $a \neq 0$. By using a suitable bilinear transformation

$$\psi(z) = e^{i\theta}(z-p)/(1-z\bar{p}), \quad |p| < 1,$$

and the first part of the proof, we obtain

$$f'(z) = \lambda \frac{[\psi(z)]^n}{\delta_n^n (1 - \delta_n^2)} \psi'(z) \quad \text{for some } n, \quad |\lambda| = 1.$$

A routine calculation gives

$$f'(z) = \lambda e^{i(n+1)\theta} \frac{(z-p)^n (1-|p|^2)}{\delta_n^n (1 - \delta_n^2) (1 - z\bar{p})^{n+2}}$$

and $L(f) = C(a, r)$.

From Theorem 1 and its proof we deduce

THEOREM 2 (CIMA AND WOGEN). *For each $a \in D$ there is a countable family of circles $\gamma_n(a)$ in D such that:*

(i) *If f is in the ball $\tilde{B}_0$ and $L(f)$ contains a circle γ , then $\gamma = \gamma_n(a)$ for some (a, n) .*

(ii) *Conversely, given $\gamma_n(a)$ there is a function f_{na} in the ball $\tilde{B}_0$ so that $L(f_{na}) = \gamma_n(a)$.*

The results in this paper were presented by the authors on September 29, 1983 at the VI Congreso de Ecuaciones Diferenciales y Aplicaciones at the University of Jaca-Zaragoza (Spain).

The authors are grateful to the referee for his suggestions.

REFERENCES

1. J. A. Cima and W. R. Wogen, *Extreme points of the unit ball of the Bloch space B_0* , Michigan Math. J. **25** (1978), 213-221.
2. K. J. Wirths, *On holomorphic functions satisfying $|f(z)|(1-|z|^2) \leq 1$ in the unit disc*, Proc. Amer. Math. Soc. **85** (1982), 19-23.

DEPARTAMENTO DE MATEMÁTICA PURA, ETSICCP, UNIVERSIDAD POLITÉCNICA, VALENCIA-22, SPAIN