DOUBLE COMMUTANTS OF OPERATORS QUASI-SIMILAR TO NORMAL OPERATORS

KATSUTOSHI TAKAHASHI

ABSTRACT. It is shown that double commutants of operators quasi-similar to normal operators are reflexive.

An algebra $\mathcal{A}$ of bounded linear operators on a Hilbert space is said to be *reflexive* if $\mathcal{A} = \text{Alg Lat } \mathcal{A}$, where $\text{Lat } \mathcal{A}$ denotes the family of all subspaces invariant under all elements of $\mathcal{A}$ and $\text{Alg Lat } \mathcal{A}$ is the algebra of all operators X for which $X\mathcal{M} \subseteq \mathcal{M}$ for every $\mathcal{M} \in \text{Lat } \mathcal{A}$. If $\mathcal{A}$ is a von Neumann algebra, then $\mathcal{M} \in \text{Lat } \mathcal{A}$ means that the projection to $\mathcal{M}$ belongs to the commutant $\mathcal{A}'$, therefore $\text{Alg Lat } \mathcal{A}$ coincides with the double commutant $\mathcal{A}''$. Thus the von Neumann double commutant theorem can be rephrased as follows (cf. [4, Theorem 9.17]): every von Neumann algebra is reflexive.

Recall some definitions. For operators T_1 and T_2 , let us write $T_1 \prec_i T_2$ (resp. $T_1 \prec T_2$) to denote that there exists a *quasi-affinity*, i.e. injection with dense range (resp. an injection) X such that $XT_1 = T_2X$. T_1 and T_2 are *quasi-similar* by definition if $T_1 \prec_i T_2$ and $T_2 \prec_i T_1$.

If N is a normal operator, the commutant $\{N\}'$ becomes a von Neumann algebra by the Fuglede theorem, so that both the commutant and the double commutant $\{N\}''$ are reflexive. If an operator T is quasi-similar to a normal operator N , the commutant $\{T\}'$ is reflexive because the reflexivity of commutant is preserved under quasi-similarity [1] and $\{N\}'$ is reflexive. In this note we shall show that the double commutant $\{T\}''$ is reflexive too.

THEOREM. *If an operator T is quasi-similar to a normal operator N , then the double commutant $\{T\}''$ is reflexive.*

Before going into the proof, let us present an immediate consequence.

COROLLARY. *If T is a contraction of class C_{11} , that is, $\lim_{n \rightarrow \infty} \|T^n x\| \neq 0$ and $\lim_{n \rightarrow \infty} \|T^{*n} x\| \neq 0$ for every nonzero x , then the double commutant $\{T\}''$ is reflexive.*

In fact, it is well known that a contraction of class C_{11} is quasi-similar to a unitary operator (see [5, Proposition II.3.5]).

Received by the editors November 11, 1983.

1980 *Mathematics Subject Classification.* Primary 47C05, 47B15; Secondary 47A45.

Key words and phrases. Double commutant, reflexive, quasi-similar, normal operator, C_{11} -contraction.

©1984 American Mathematical Society

0002-9939/84 \$1.00 + \$.25 per page

This result for a C_{11} -contraction T was proved in [3, Theorem 3] under the condition that every injection in $\{T\}'$ has dense range.

For the proof of the theorem we need a lemma, which is an analog of [3, Lemma 5] for C_{11} -contractions.

LEMMA. *Let T be an operator quasi-similar to a normal operator. Then, for $\mathcal{M} \in \text{Lat } T$ (= the family of all T -invariant subspaces), the following conditions are mutually equivalent.*

- (i) $T|_{\mathcal{M}}$ is quasi-similar to a normal operator.
- (ii) $T|_{\mathcal{M}} \succ N_1$ for some normal operator N_1 .
- (iii) $\mathcal{M} = (\text{ran } Z)^{\perp}$ for some $Z \in \{T\}'$.

If one of these conditions holds, then $\mathcal{M} \in \text{Lat}\{T\}''$.

PROOF. Let X and Y be quasi-affinities such that $XT = NX$ and $TY = YN$ for some normal operator N on $\mathcal{G}$. (i) $\Rightarrow$ (ii) is trivial.

(ii) $\Rightarrow$ (iii): Since $N_1 \prec T|_{\mathcal{M}} \prec N|(X\mathcal{M})^{\perp}$, it follows from [2, Lemma 4.1] that $(X\mathcal{M})^{\perp}$ is N -reducing and $N|(X\mathcal{M})^{\perp}$ is unitarily equivalent to N_1 . Therefore we have $(T|_{\mathcal{M}})^* \prec N_1^* \prec N^*|(X\mathcal{M})^{\perp} \prec N^* \prec T^*$. Denote by W the injection from $\mathcal{M}$ to $\mathcal{H}$ (= the space on which T acts) such that $W(T|_{\mathcal{M}})^* = T^*W$, and by J the inclusion mapping of $\mathcal{M}$ into $\mathcal{H}$. Then $Z = JW^*$ is an operator required in (iii).

(iii) $\Rightarrow$ (i): Define an operator K from $\mathcal{G}$ to $\mathcal{M}$ by $Kx = ZYx$ for $x \in \mathcal{G}$. Then K has dense range and $(T|_{\mathcal{M}})K = KN$. Also $(X|_{\mathcal{M}})(T|_{\mathcal{M}}) = N(X|_{\mathcal{M}})$ and $X|_{\mathcal{M}}$ is injective. Therefore it follows from [2, Lemma 4.1] that $N|(\ker K)^{\perp}$ and $N|(X\mathcal{M})^{\perp}$ are unitarily equivalent normal operators, and so $T|_{\mathcal{M}}$ is quasi-similar to a normal operator $N|(X\mathcal{M})^{\perp}$. Finally it is clear that $\mathcal{M} \in \text{Lat}\{T\}''$ for any subspace $\mathcal{M}$ satisfying (iii).

PROOF OF THEOREM. By assumption there are quasi-affinities X and Y satisfying $XT = NX$ and $TY = YN$. Without loss of generality, the product XY can be assumed to be nonnegative. Indeed, since $XY \in \{N\}'$, XY admits the polar factorization $XY = UP$ in the von Neumann algebra $\{N\}'$, where U is unitary and P is nonnegative. The quasi-affinities U^*X and Y satisfy the required condition.

The inclusion $\{T\}'' \subseteq \text{Alg Lat}\{T\}''$ is obvious. To see the converse inclusion, take $A \in \text{Alg Lat}\{T\}''$. In order to show $A \in \{T\}''$, it suffices to prove that A commutes with YCX for all $C \in \{N\}'$. In fact, then A commutes with YX , and also with $YXBYX$ for any $B \in \{T\}'$, because XBY belongs to $\{N\}'$. Therefore we have

$$YXBAYX = YXBYXA = AYXBYX = YXABYX.$$

Then the quasi-affinity properties of X and Y imply $BA = AB$, hence $A \in \{T\}''$.

Let us show that A commutes with YCX for all $C \in \{N\}'$. Since $\{N\}'$ is a von Neumann algebra, we have only to show that $A(YHX) = (YHX)A$ for any selfadjoint injection $H \in \{N\}'$. Take $\mathcal{M} \in \text{Lat}\{XYHXY\}'$. Since $\{N\}'' \subseteq \{XYHXY\}'$, we have $\mathcal{M} \in \text{Lat}\{N\}''$, hence $\mathcal{M}$ is N -reducing and $N|_{\mathcal{M}}$ is normal. Then, since $(Y\mathcal{M})^{\perp} \in \text{Lat } T$ and $T|(Y\mathcal{M})^{\perp} \succ N|_{\mathcal{M}}$, it follows from the Lemma that $(Y\mathcal{M})^{\perp} \in \text{Lat}\{T\}'' \subseteq \text{Lat } A$, and so $XYHXY\mathcal{M} \subseteq (XYHXY\mathcal{M})^{\perp} \subseteq \mathcal{M}$. We can conclude that $XYHXY\mathcal{M} \in \text{Alg Lat}\{XYHXY\}'$. Then, since the commutant $\{XYHXY\}'$ of the

selfadjoint operator $XYHXY$ is reflexive, $XYHXAY \in \{XYHXY\}'$. Taking $H = I$, we have $(XY)^2XYXAY = XYXAY(XY)^2$ and $(XY)^2XAY = XAY(XY)^2$ by the injectivity of XY . Since XY is assumed to be nonnegative, taking the square root of $(XY)^2$, we obtain $XYXAY = XAYXY$. Then for any selfadjoint injection H we have

$$XYHXYXYHXAY = XYHXAYXYHXY = XYHXYXAYHXY.$$

Finally it follows from the injectivity of $XYHXYX$ and the dense range property of Y that $YHXA = AYHX$.

REFERENCES

1. H. Bercovici, C. Foiaş and B. Sz.-Nagy, *Reflexive and hyper-reflexive operators of class C_0* , Acta Sci. Math. **43** (1981), 5–13.
2. R. G. Douglas, *On the operator equation $S^*XT = X$ and related topics*, Acta Sci. Math. **30** (1969), 19–32.
3. L. Kerchy, *On invariant subspace lattices of C_{11} -contractions*, Acta Sci. Math. **43** (1981), 281–293.
4. H. Radjavi and P. Rosenthal, *Invariant subspaces*, Springer-Verlag, Berlin, Heidelberg and New York, 1973.
5. B. Sz.-Nagy and C. Foiaş, *Harmonic analysis of operators on Hilbert space*, North-Holland, Amsterdam, 1970.

DEPARTMENT OF MATHEMATICS, SAPPORO MEDICAL COLLEGE, MINAMI-1 NISHI-17, SAPPORO, JAPAN